

PAEDIATRICS

Pharmacological prevention of sevoflurane- and desflurane-related emergence agitation in children: a meta-analysis of published studies

S. Dahmani^{1*}, I. Stany¹, C. Brasher¹, C. Lejeune¹, B. Bruneau¹, C. Wood¹, Y. Nivoche¹, I. Constant² and I. Murat²

¹Department of Anesthesiology, Robert Debre University Hospital, Assistance Publique des Hôpitaux de Paris, 48 Bd Serurier, 75019 Paris, France. ²Department of Anesthesiology, Trousseau University Hospital, Assistance Publique des Hôpitaux de Paris, 26 avenue du Docteur Arnold-Netter, Paris, France

*Corresponding author. E-mail: souhayl.dahmani@rdb.aphp.fr

Background. Emergence agitation (EA) in children is increased after sevoflurane anaesthesia. The efficacy of prophylactic treatment is controversial. The aim of this study was to provide a meta-analysis of the studies of the pharmacological prevention of EA in children.

Methods. A comprehensive literature search was conducted to identify clinical trials that focused on the prevention of EA in children anaesthetized with sevoflurane, desflurane, or both. The data from each trial were combined using the Mantel–Haenszel model to calculate the pooled odds ratio (OR) and 95% confidence interval. I^2 statistics were used to assess statistics heterogeneity and the funnel plot and the Begg–Mazumdar test to assess bias.

Results. Thirty-seven articles were found which included a total of 1695 patients in the intervention groups and 1477 in the control ones. Midazolam and $5HT_3$ inhibitors were not found to have a protective effect against EA [OR=0.88 (0.44, 1.76); OR=0.39 (0.12, 1.31), respectively], whereas propofol [OR=0.21 (0.16, 0.28)], ketamine [OR=0.28 (0.13, 0.60)], α_2 -adrenoceptors [OR=0.23 (0.17, 0.33)], fentanyl [OR=0.31 (0.18, 0.56)], and peroperative analgesia [OR=0.15 (0.07, 0.34)] were all found to have a preventive effect. Subgroup analysis according to the peroperative analgesia given does not affect the results.

Conclusions. This meta-analysis found that propofol, ketamine, fentanyl, and preoperative analgesia had a prophylactic effect in preventing EA. The analgesic properties of these drugs do not seem to have a role in this effect.

Br | Anaesth 2010; 104: 216-23

Keywords: anaesthetic techniques, regional, caudal; anaesthetics i.v., clonidine; anaesthetics i.v., fentanyl; anaesthetics i.v., propofol; emergence agitation

Accepted for publication: November 23, 2009

The increased use of sevoflurane and desflurane in developing countries has been accompanied by an increase in emergence agitation (EA), a postoperative behavioural disorder. First described in the early 1960s, EA is characterized by a variety of presentations including crying, excitation, agitation, and delirium occurring during the early stage of emergence from anaesthesia in children.²

With sevoflurane or desflurane anaesthesia, the incidence of EA varies widely between 2% and 80% depending on the scoring system and the anaesthetic technique

used, and is more frequently observed in preschool children.^{3–5} Despite its spontaneous resolution, EA is still considered as a potentially serious complication because of the risks of self-injury, and because of the stress caused to both caregivers and families.

The incidence of EA has led many authors to propose prophylactic treatment to reduce its incidence. These have included propofol, α_2 -adrenoceptor (AR) agonists, midazolam, and ketamine. However, their efficacy remains the subject of debate.

Meta-analysis is a statistical method allowing aggregation and quantification of the therapeutic effects from multiple studies. The aim of this study was to perform a meta-analysis of the efficacy of pharmacological prophylactic interventions currently proposed to reduce the incidence of EA.

Methods

The bibliographic search and analysis for this meta-analysis was conducted according to the guidelines of the Cochrane Handbook for systematic reviews of intervention and the QUORUM statements.⁹

The databases searched included Pubmed, Embase, and the Cochrane Database for Systemic Review. As descriptions of EA including agitation, behavioural problems, and delirium, the following search terms were used: 'agitation, and sevoflurane or desflurane, and children or infant, behavior, and sevoflurane or desflurane, and children or infant, delirium, and sevoflurane or desflurane, and children and infant'. Only articles published in English were considered. Where the full text could not be found, authors were contacted to provide a copy of the original article.

The articles obtained were independently analysed by two senior anaesthetists and those meeting the following criteria were included in the analysis: randomized controlled trial, double-blinded (articles in whom the evaluation was blinded were included), sevoflurane or desflurane anaesthesia, absence of neurological disease, standardized anaesthesia protocols, agitation or delirium or behavioural disturbances as one of the study endpoints, a control group, and a standardized definition of EA between the control and intervention groups in each study.

We excluded from analysis all studies comparing two prophylactic agents or interventions, and those exploring curative treatments for EA. In addition, a manual search of references cited in the selected articles (including reviews and meta-analyses) was performed. The most recent search date was February 2009.

Two independent readers assessed article quality and extracted data. Retained data included the age of patients, type of surgery, sedative premedication (dose, timing, and route of administration), dose, timing, and route of administration of prophylactic agent or technique, hypnotic agents, intraoperative opioids used, peroperative analgesia, and the percentage of patients experiencing agitation in each group based on the criteria chosen by the authors. When conflicting results were found, the article was rechecked by both anaesthetists.

Statistical analysis was performed using the Review Manager 5 software (RevMan 5, The Cochrane Collaboration, Oxford, UK). The analysis of the incidence of EA was performed using the odds ratio (OR) computed using the Mantel-Haenszel method (fixed or random models). The OR represents the odds of EA occurring in

the prophylactic group compared with the control group. A confidence interval for the OR of <1 indicates efficacy in EA prevention. In order to minimize the publication bias, some data transformations were necessary. Secondly, OR computation requires the results to be expressed as percentages. Where outcomes were given as mixed percentages and continuous results, a standardized mean ratio was first computed, then transformed as partial OR using the formula of Chinn: 10 Ln OR=1.814×SMR (Ln: logarithm). The data were then entered as Ln(OR) and sp(LnOR). Overall OR (and 95% confidence interval) was finally computed with the inverse variance method [http://www.cochrane-handbook.org/ (Section 9.4.6)].

To assess the impact of study heterogeneity on the results of the meta-analysis, the differences of study design regarding premedication, anaesthetic agent used (sevoflurane or desflurane), modalities of pain treatment, and parental presence during induction and recovery from anaesthesia, an I^2 test was performed. According to the Cochrane review guidelines, an $I^2 > 40\%$ and a P-value of <0.1 are considered as the threshold for heterogeneity and indicate the use of a random effect in OR calculation.

Each prophylactic agent was tested grouping all administration routes and timing modalities. Then, for each drug, EA prevention was analysed according to timing of administration, route of administration, anaesthesia agent used, concurrent preoperative analgesia, and pain expected as a result of the intervention (anaesthesia for MRI *vs* surgery). In addition, if statistical heterogeneity was present, analysis was carried out by removing studies one by one according to the route and timing of drug administration. Studies displaying statistical heterogeneity were then further analysed for differences in design and results. Results are expressed as OR (95% confidence interval), I^2 , and P-value for heterogeneity.

In the case of studies with more than one intervention group, each group was considered as a study in the meta-analysis and compared with the control group. Finally, to avoid calculation problems related to zero effectives, 1 was added to all groups in these cases. Bias related to unpublished studies was assessed for analysis by aggregating at least 10 studies. The bias was assessed using the funnel plot and the Begg-Mazumdar test.

Results

A total of 324 articles were identified using the search criteria, and 58 relevant articles identified for meta-analysis. The details of the selection process are summarized in Figure 1. Articles examining the efficacy of physostigmine, 11 remifentanil 12 (in addition, the full text for this article was not found), desflurane, 13 delayed emergence, 14 and N₂O administration during sevoflurane washout 15 were discarded after applying selection criteria. When full text could not be found, the authors were contacted twice (without response in all cases).

The meta-analysis included 37 articles (1695 patients in the intervention groups and 1477 in the control groups). The pharmacological treatments studied for prevention of EA were: midazolam $^{16-20}$ (four articles), propofol 16 $^{21-32}$ as single bolus or continuous administration (13), fentanyl $^{33-37}$ (five), ketamine 29 $^{38-40}$ (four), α_2 -AR agonists $^{41-50}$ clonidine or dexmedetomidine, given orally, by i.v. or caudal route (10), peroperative local




Fig 1 Meta analysis flowchart. RCT, randomized controlled trials.

anaesthetics, 39 51 52 caudal analgesia, and peroperative parenteral local anaesthetics (three), and 5 HT $_3$ inhibitors 45 53 either ondansetron or tropisotron (two) (Supplementary Table S1).

Midazolam, given as either premedication 30 min before induction of anaesthesia or after induction, does not have a prophylactic effect against EA [OR=0.88 (0.44, 1.76); I^2 =47%, P=0.11, Supplementary Fig. S1). The heterogeneity identified was explored by analysing studies by the timing of midazolam administration. Removing one study 16 in which midazolam was given after induction did not correct heterogeneity. When subgroup analysis was performed using the preoperative analgesia given, heterogeneity was within acceptable values. Analysis including studies with or without preoperative analgesia decreased heterogeneity, but midazolam was still found to be ineffective in preventing EA [OR=1.88 (0.85, 4.13), I^2 =0%, P=0.79; OR=0.58 (0.31, 1.09), $I^2=9\%$, P=0.33, respectively]. Finally, excluding the study using desflurane 16 had no effect on the results [OR=0.96 (0.46, 2.01), I^2 =55%, P=0.09].

Propofol showed an overall protective effect against EA $[OR=0.21 \ (0.16,\ 0.28),\ I^2=52\%,\ P=0.01,\ Fig.\ 2].$ Subgroup analysis of studies by the timing of administration (continuous, bolus after induction, or bolus at the end of anaesthesia) demonstrated that continuous administration and a bolus dose at the end of anaesthesia were protective $[OR=0.17\ (0.11,\ 0.27),\ I^2=36\%,\ P=0.13;$ $OR=0.21\ (0.09,\ 0.50),\ I^2=0,\ P=0.47,\ respectively],$ whereas bolus at induction was ineffective in preventing EA $[OR=0.46\ (0.20,\ 1.06),\ I^2=40\%,\ P=0.19].$ Subgroup analysis of the studies showing efficacy, that is, those using an end of surgery bolus or continuous administration, by administration of peroperative analgesia found no effect of this on propofol efficacy $[OR=0.15\ (0.06,\ 0.36),\ I^2=0\%,\ P=0.46;\ OR=0.19\ (0.13,\ 0.28),\ I^2=0\%,$

Study or subgroup	log[]	SE	Weight	IV, Fixed, 95% CI	IV, Fixed, 95% CI		
Abou-Shahwan, 2008	-1.97	0.65	5.3%	0.14 (0.04, 0.50)			
Aouad, 2007	-1.31	0.27	30.6%	0.27 (0.16, 0.46)			
Cohen, 2002	-2.04	1.26	1.4%	0.13 (0.01, 1.54)			
Cohen, 2003	0.85	0.85	3.1%	2.34 (0.44, 12.38)			
Grundmann, 1998	-1.61	0.41	13.3%	0.20 (0.09, 0.45)			
Lopez Gil, 1999	-2.41	1.14	1.7%	0.09 (0.01, 0.84)			
Nakayama, 2007_1	-2.21	0.63	5.6%	0.11 (0.03, 0.38)			
Nakayama, 2007_2	-2.21	1.18	1.6%	0.11 (0.01, 1.11)			
Ozdemir Kol, 2008	-2.1	0.44	11.5%	0.12 (0.05, 0.29)			
Picard, 2000	-2.12	0.72	4.3%	0.12 (0.03, 0.49)			
Steinmetz, 2007	0.27	2.11	0.5%	1.31 (0.02, 81.90)			
Tsai, 2008	0.29	0.58	6.6%	1.34 (0.43, 4.17)			
Uezono, 2000	-2.41	1.31	1.3%	0.09 (0.01, 1.17)			
Vitanen, 1999	-2.21	0.41	13.3%	0.11 (0.05, 0.25)			
Total (95% CI)			100.0%	0.21 (0.16, 0.28)	♦		
Heterogeneity: $\chi^2 = 27.35$, d	f - 13 (P - 0	+ + + + + + + + + + + + + + + + + + + +					
Test for overall effect: $Z = 10$		0.002 0.1 1 10 500					
1001101 0401411 011001. 2 = 10	7.40 (1 < 0.00	Favours propofol Favours control					

Fig 2 Forest plot of meta-analysis of propofol prevention in EA (propofol n=442, control n=450). The square shown for each study (first author and year of publication) is the OR for individual trials, and the corresponding horizontal line is the 95% confidence interval (CI). The diamond is the pooled OR with the 95% CI. Studies with more than one intervention group were numbered (author, year of publication_1) and (author, year of publication_2).

P=0.55, respectively]. Analysis of the studies using sevoflurane, by removing the one study that used desflurane,²¹ still found propofol effective for EA [0.21 (0.12, 0.37), I^2 =56%, P=0.007].

Ketamine administration was found to prevent EA $[OR=0.28 \ (0.13,\ 0.60),\ I^2=0\%,\ P=0.68,\ Supplementary Fig. S2].$ Subgroup analysis found no effect of the absence of peroperative analgesia on the preventive effect of ketamine $[OR=0.34\ (0.14,\ 0.86),\ I^2=0\%,\ P=0.84].$ One study³⁸ used preoperative fentanyl and found that ketamine had a preventative effect $[OR=0.17\ (0.04,\ 0.70)].$ After removing the desflurane study³⁸ from the analysis, ketamine still had a preventive effect against EA $[OR=0.34\ (0.14,\ 0.86),\ I^2=0\%,\ P=0.84].$

 α_2 -AR agonists were also found to be protective [OR=0.23 (0.17, 0.33), I^2 =24%, P=0.2, Fig. 3]. Subgroup analysis of the route of administration (i.v. or

caudal), the α_2 -AR agonist used (dexmedetomidine or clonidine), and concurrent peroperative analgesia (none, preoperative analgesia, and locoregional analgesia) found no effect of these factors on the efficacy of α_2 -AR agonists against ER (Supplementary Table S2).

Peroperative fentanyl was also found to protect against EA [OR=0.31 (0.18, 0.56), I^2 =47%, P=0.06, Fig. 4]. Subgroup analysis of the route of administration decreased heterogeneity but found intranasal fentanyl to exhibit a preventive effect against EA [OR=0.23 (0.13, 0.43), I^2 =28%, P=0.24] whereas i.v. fentanyl did not [OR=0.35 (0.12, 1.06), I^2 =20%, P=0.29]. Analysing studies in which sevoflurane was used showed fentanyl to still be preventative against EA [OR=0.29 (0.15, 0.54), I^2 =52%, P=0.05].

Peroperative analgesia was also found to protect against EA [OR=0.15 (0.07, 0.34), I^2 =8%, P=0.36, Supplementary Fig. S3]. For the i.v. route, homogeneity




Fig 3 Forest plot of meta-analysis of α_2 -AR agonists prevention in EA (α_2 -AR n=418, control n=339). The square shown for each study (first author and year of publication) is the OR for individual trials, and the corresponding horizontal line is the 95% CI. The diamond is the pooled OR with the CI. Studies with more than one intervention group were numbered (author, year of publication_1) and (author, year of publication_2).

Otrodo o monte manage	Fentanyl		Control		Odds ratio		Odds ratio	
Study or subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
Binstock, 2004_1	10	50	28	51	17.1%	0.21 (0.08, 0.50)		
Binstock, 2004_2	25	50	28	51	18.8%	0.82 (0.38, 1.80)		_
Cravero, 2003	2	16	9	16	7.7%	0.11 (0.02, 0.66)		
Demirbilek, 2004_1	2	30	4	30	7.7%	0.46 (0.08, 2.75)	-	
Demirbilek, 2004_2	3	30	4	30	9.0%	0.72 (0.15, 3.54)	-	
Finkel, 2001_1	13	51	22	49	17.8%	0.42 (0.18, 0.98)		
Finkel, 2001_2	7	50	22	49	15.7%	0.20 (0.08, 0.53)		
Gallinkin, 2000	1	64	16	69	6.2%	0.05 (0.01, 0.41)		
Total (95% CI)		341		345	100.0%	0.31 (0.18, 0.56)	♦	
Total events	63		133					
Heterogeneity: $\tau^2 = 0.3$	 	 						
Test for overall effect: 2	0.01 0.1 1	10 100						
. ccc. crotan oncon z	Favours fentanyl	Favours control						

Fig 4 Forest plot of meta-analysis of fentanyl prevention in EA (fentanyl n=341, control n=215). The square shown for each study (first author and year of publication) is the OR for individual trials, and the corresponding horizontal line is the 95% CI. The diamond is the pooled OR with the 95% CI. Studies with more than one intervention group were numbered (author, year of publication_1) and (author, year of publication_2).




Fig 5 Funnel plot of the studies included in propofol meta-analysis.




Fig 6 Funnel plot of the studies included in α_2 -AR agonists meta-analysis.

was acceptable [OR=0.19; 95% confidence interval 0.06–0.6; I^2 =30%, P=0.23]. As there was only one study of caudal analgesia, ¹ meta-analysis was not indicated.

Finally, meta-analysis of the two studies examining the preventive effects of 5HT_3 antagonists on EA found these drugs to be ineffective (OR=0.39, 95% confidence interval 0.12-1.31; I^2 =0%, P=0.56, Supplementary Fig. S4).

Bias was assessed for propofol (13 studies) and α_2 -AR agonists (12 studies). The results for propofol were a symmetric funnel plot (Fig. 5) and a Begg-Mazumdar Kendall's τ =0.15 (P=0.47), indicating the absence of bias. However, for α_2 -AR agonists, analysis indicates the possibility of bias (asymmetric funnel plot, Fig. 6; Begg-Mazumdar Kendall's τ =-0.56, P=0.004).

Discussion

The main findings of this meta-analysis are that midazolam, classically considered as the 'ideal' drug for premedication appears to be ineffective in the prevention of EA, propofol seems to be effective, but is dependent upon the timing of administration, and that ketamine, α_2 -AR agonists, fentanyl, and peroperative analgesia were all effective methods of EA prevention.

The aetiology of EA is currently unknown. Recent hypotheses emphasize rapid emergence associated with new anaesthetic agents such as sevoflurane and desflurane. This may create a dissociative state, that is, children awaken with altered cognitive perception. The efficacy of 5HT₃ serotonin receptor antagonists, reported in one study, suggests the involvement of the serotoninergic system processes in the aetiology of EA. Other factors have been proposed as contributing to or exacerbating this problem, including postoperative pain and preoperative anxiety. EA has been shown to be associated with preoperative anxiety, and to be prevented by parental presence and pain prevention.

On this basis, midazolam, an anxiolytic agent widely used as premedication, would appear to be a logical candidate for preventing EA. However, our meta-analysis does not support this. In addition, statistical heterogeneity was decreased when subgroup analysis was performed taking into account peroperative analgesia. Two studies¹⁸ in which patients were given alfentanil and acetaminophen peroperatively found no preventative effect, whereas studies without preoperative analgesia found midazolam to be efficient in preventing EA. Thus, analgesia, sedation, or both produced by opioids may blunt any effect of midazolam.

Propofol delays or modifies emergence and could decrease the incidence of EA. However, the rapid pharmacokinetics of this agent and the relatively low doses given in selected studies (1 mg kg⁻¹) could explain the failure of bolus doses to prevent EA when given during and after induction. This argument is supported by the efficacy of both end-intervention boluses and continuous administration of propofol during maintenance of anaesthesia, as propofol concentration at emergence would be more effective in preventing EA.

Ketamine, α_2 -agonists, fentanyl, and peroperative analgesia were effective in preventing EA. Ketamine, an *N*-methyl-D-aspartate receptor antagonist, produces both analgesic- and opioid-sparing effects when used at low doses. ⁵⁷ ⁵⁸ Dexmedetomidine and clonidine are also considered as potent analgesics and are used both pareneterally and in loco-regional anaesthesia (caudal and epidural). ⁵⁹ ⁶⁰ Finally, fentanyl is a potent opioid receptor agonist routinely used in the peroperative period. Together with the high efficacy of peroperative analgesia in the prevention of EA, these results suggest, as previously stated, a relationship between postoperative pain and agitation. ³⁹

However, subgroup analysis of peroperative analgesia found no influence of this factor on the efficacy of ketamine, propofol, and α_2 -agonists on EA. In addition, i.v. peroperative fentanyl failed to prevent EA. Finally, the greater efficacy of ketamine, α_2 -agonists, and fentanyl in postoperative pain relief was not a constant finding in the analysed articles and was not always a study endpoint. These results suggest that the analgesic properties of these compounds are unlikely to be involved in their preventive effects. Other mechanisms such as potentiation of anaesthesia or sedation might be involved. This is particularly pertinent for α_2 -AR agonists. They have been found to decrease anaesthetic requirements and to induce emergence sedation⁶¹ which could explain their preventive effect on EA. Finally, studies performed during anaesthesia for imaging found EA to occur. 31 35 39 42 despite the non-painful character of this procedure. Excluding them from analysis did not impact upon the efficacy of any of these drugs.

It has been proposed that 5HT₃ antagonists administered for prevention of postoperative nausea and vomiting may also prevent EA. Our results do not support this proposition, but this is based on only two studies and further work is needed before any solid conclusion.

Bias analysis found both funnel plots and Begg–Mazumdar testing suggestive of publication bias in α_2 -AR agonists meta-analysis. Consequently, meta-analysis results for these drugs must be interpreted cautiously, because of the risk of unpublished negative results.

The quality of a meta-analysis relies upon two important factors: the quality of selected studies, and their heterogeneity and bias detection. Concerning article quality, the articles included met strict selection criteria as described above. All studies were randomized and doubleblinded, and anaesthesia protocols and postoperative evaluation were standardized in the intervention and control groups. Concerning heterogeneity and bias, the articles included were heterogeneous because of their anaesthesia and analgesia protocols. Study design varied in terms of: premedication given, administration of peroperative analgesics, the route and timing of administration of the studied agent, and the evaluation tools used to quantify EA. This could explain the wide variability in the incidence of EA observed. However, all statistical analyses were performed after assessment of statistical heterogeneity, and subgroup analyses examined study design variations, resulting in acceptable statistical heterogeneity. The quality of EA evaluation was of great concern, and particular attention was paid to quantification. In all cases, EA quantification was standardized between the control and intervention groups, but the large number of unvalidated evaluation scales, along with individual scale EA threshold definitions may well impact upon EA incidence measurement, and furthermore, postoperative pain can per se induce agitation. However, only one included study used the validated paediatric anaesthesia emergence delirium (PAED) scale.⁶² Ideally, new studies on EA prevention should be based upon the use of a standardized tool. Finally, bias was assessed following the recommendations of the Cochrane collaboration.

In summary, this meta-analysis evaluating EA prevention demonstrates that propofol, pain prevention, ketamine, and α_2 -AR agonists appear to be effective. This is of interest because of the potential use of combinations of these different agents and their various routes of administration. Future studies should focus on associations of these drugs to study their effects in the prevention of EA and the use of the standardized PAED scale.

Supplementary material

Supplementary material is available at *British Journal of Anaesthesia* online.

References

- I Aouad MT, Kanazi GE, Siddik-Sayyid SM, Gerges FJ, Rizk LB, Baraka AS. Preoperative caudal block prevents emergence agitation in children following sevoflurane anesthesia. Acta Anaesthesiol Scand 2005; 49: 300-4
- 2 Yamashita M. Postanaesthetic excitation and agitation. Paediatr Anaesth 2003; 13: 641
- 3 John M. Postanaesthesia excitation. *Paediatr Anaesth* 2002; 12: 293-5
- 4 Johr M, Berger TM. Paediatric anaesthesia and inhalation agents. Best Pract Res Clin Anaesthesiol 2005; 19: 501–22
- 5 Voepel-Lewis T, Malviya S, Tait AR. A prospective cohort study of emergence agitation in the pediatric postanesthesia care unit. Anesth Analg 2003; 96: 1625-30
- 6 Silva LM, Braz LG, Modolo NS. Emergence agitation in pediatric anesthesia: current features. *J Pediatr (Rio J)* 2008; 84: 107–13
- 7 Goa KL, Noble S, Spencer CM. Sevoflurane in paediatric anaesthesia: a review. *Paediatr Drugs* 1999; 1: 127–53
- 8 Veyckemans F. Excitation and delirium during sevoflurane anesthesia in pediatric patients. *Minerva Anestesiol* 2002; **68**: 402–5
- 9 Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. Quality of Reporting of Meta-analyses. *Lancet* 1999; 354: 1896–900
- 10 Chinn S. A simple method for converting an odds ratio to effect size for use in meta-analysis. Stat Med 2000; 19: 3127-31
- 11 Funk W, Hollnberger H, Geroldinger J. Physostigmine and anaesthesia emergence delirium in preschool children: a randomized blinded trial. Eur J Anaesthesiol 2008; 25: 37–42
- 12 Gouda N. The effects of single bolus dose of remifentanil on the emergence characteristics of paediatric patients undergoing myringotomy after sevoflurane anaesthesia. Egypt J Anaesth 2004; 20: 63-6
- 13 Mayer J, Boldt J, Rohm KD, Scheuermann K, Suttner SW. Desflurane anesthesia after sevoflurane inhaled induction reduces severity of emergence agitation in children undergoing minor ear-nose-throat surgery compared with sevoflurane induction and maintenance. Anesth Analg 2006; 102: 400–4
- 14 Oh AY, Seo KS, Kim SD, Kim CS, Kim HS. Delayed emergence process does not result in a lower incidence of emergence

- agitation after sevoflurane anesthesia in children. Acta Anaesthesiol Scand 2005; 49: 297-9
- 15 Shibata S, Shigeomi S, Sato W, Enzan K. Nitrous oxide administration during washout of sevoflurane improves postanesthetic agitation in children. J Anesth 2005; 19: 160-3
- 16 Cohen IT, Drewsen S, Hannallah RS. Propofol or midazolam do not reduce the incidence of emergence agitation associated with desflurane anaesthesia in children undergoing adenotonsillectomy. Paediatr Anaesth 2002; 12: 604–9
- 17 Kain ZN, Caldwell-Andrews AA, Mayes LC, et al. Family-centered preparation for surgery improves perioperative outcomes in children: a randomized controlled trial. Anesthesiology 2007; 106: 65-74
- 18 Viitanen H, Annila P, Viitanen M, Tarkkila P. Premedication with midazolam delays recovery after ambulatory sevoflurane anesthesia in children. Anesth Analg 1999; 89: 75–9
- 19 Viitanen H, Annila P, Viitanen M, Yli-Hankala A. Midazolam premedication delays recovery from propofol-induced sevoflurane anesthesia in children 1–3 yr. Can | Anaesth 1999; 46: 766–71
- 20 Lapin SL, Auden SM, Goldsmith LJ, Reynolds AM. Effects of sevoflurane anaesthesia on recovery in children: a comparison with halothane. *Paediatr Anaesth* 1999: 9: 299–304
- 21 Grundmann U, Uth M, Eichner A, Wilhelm W, Larsen R. Total intravenous anaesthesia with propofol and remifentanil in paediatric patients: a comparison with a desflurane–nitrous oxide inhalation anaesthesia. Acta Anaesthesiol Scand 1998: 42: 845–50
- 22 Steinmetz J, Holm-Knudsen R, Sorensen MK, Eriksen K, Rasmussen LS. Hemodynamic differences between propofol-remifentanil and sevoflurane anesthesia for repair of cleft lip and palate in infants. *Paediatr Anaesth* 2007; 17: 32–7
- 23 Lopez Gil ML, Brimacombe J, Clar B. Sevoflurane versus propofol for induction and maintenance of anaesthesia with the laryngeal mask airway in children. *Paediatr Anaesth* 1999; 9: 485–90
- 24 Picard V, Dumont L, Pellegrini M. Quality of recovery in children: sevoflurane versus propofol. Acta Anaesthesiol Scand 2000; 44: 307–10
- 25 Uezono S, Goto T, Terui K, et al. Emergence agitation after sevoflurane versus propofol in pediatric patients. Anesth Analg 2000; 91: 563-6
- 26 Nakayama S, Furukawa H, Yanai H. Propofol reduces the incidence of emergence agitation in preschool-aged children as well as in school-aged children: a comparison with sevoflurane. J Anesth 2007; 21: 19–23
- 27 Cohen IT, Finkel JC, Hannallah RS, Hummer KA, Patel KM. Rapid emergence does not explain agitation following sevoflurane anaesthesia in infants and children: a comparison with propofol. *Paediatr Anaesth* 2003; 13: 63–7
- 28 Viitanen H, Tarkkila P, Mennander S, Viitanen M, Annila P. Sevoflurane-maintained anesthesia induced with propofol or sevoflurane in small children: induction and recovery characteristics. *Can J Anaesth* 1999; 46: 21–8
- 29 Tsai PS, Hsu YW, Lin CS, Ko YP, Huang CJ. Ketamine but not propofol provides additional effects on attenuating sevofluraneinduced emergence agitation in midazolam premedicated pediatric patients. *Paediatr Anaesth* 2008; 18: 1114–5
- 30 Aouad MT, Yazbeck-Karam VG, Nasr VG, El-Khatib MF, Kanazi GE, Bleik JH. A single dose of propofol at the end of surgery for the prevention of emergence agitation in children undergoing strabismus surgery during sevoflurane anesthesia. *Anesthesiology* 2007; 107: 733–8
- 31 Abu-Shahwan I. Effect of propofol on emergence behavior in children after sevoflurane general anesthesia. *Paediatr Anaesth* 2008; 18: 55–9

- 32 Ozdemir Kol I, Egilmez H, Kaygusuz K, Gursoy S, Mimaroglu C. Open-label, prospective, randomized comparison of propofol and sevoflurane for laryngeal mask anesthesia for magnetic resonance imaging in pediatric patients. *Clin Ther* 2008; 30: 175–81
- 33 Galinkin JL, Fazi LM, Cuy RM, et al. Use of intranasal fentanyl in children undergoing myringotomy and tube placement during halothane and sevoflurane anesthesia. Anesthesiology 2000; 93: 1378–83
- **34** Finkel JC, Cohen IT, Hannallah RS, et al. The effect of intranasal fentanyl on the emergence characteristics after sevoflurane anesthesia in children undergoing surgery for bilateral myringotomy tube placement. *Anesth Analg* 2001; **92**: 1164–8
- 35 Cravero JP, Beach M, Thyr B, Whalen K. The effect of small dose fentanyl on the emergence characteristics of pediatric patients after sevoflurane anesthesia without surgery. Anesth Analg 2003; 97: 364–7
- 36 Demirbilek S, Togal T, Cicek M, Aslan U, Sizanli E, Ersoy MO. Effects of fentanyl on the incidence of emergence agitation in children receiving desflurane or sevoflurane anaesthesia. Eur J Anaesthesiol 2004; 21: 538–42
- 37 Binstock W, Rubin R, Bachman C, Kahana M, McDade W, Lynch JP. The effect of premedication with OTFC, with or without ondansetron, on postoperative agitation, and nausea and vomiting in pediatric ambulatory patients. *Paediatr Anaesth* 2004; 14: 759-67
- 38 Kararmaz A, Kaya S, Turhanoglu S, Ozyilmaz MA. Oral ketamine premedication can prevent emergence agitation in children after desflurane anaesthesia. *Paediatr Anaesth* 2004; 14: 477–82
- 39 Dalens BJ, Pinard AM, Letourneau DR, Albert NT, Truchon RJ. Prevention of emergence agitation after sevoflurane anesthesia for pediatric cerebral magnetic resonance imaging by small doses of ketamine or nalbuphine administered just before discontinuing anesthesia. Anesth Analg 2006; 102: 1056–61
- 40 Abu-Shahwan I, Chowdary K. Ketamine is effective in decreasing the incidence of emergence agitation in children undergoing dental repair under sevoflurane general anesthesia. *Paediatr Angesth* 2007: 17: 846–50
- 41 Shukry M, Clyde MC, Kalarickal PL, Ramadhyani U. Does dexmedetomidine prevent emergence delirium in children after sevoflurane-based general anesthesia? *Paediatr Anaesth* 2005; 15: 1098–104
- 42 Isik B, Arslan M, Tunga AD, Kurtipek O. Dexmedetomidine decreases emergence agitation in pediatric patients after sevoflurane anesthesia without surgery. *Paediatr Anaesth* 2006; 16: 748-53
- 43 Ibacache ME, Munoz HR, Brandes V, Morales AL. Single-dose dexmedetomidine reduces agitation after sevoflurane anesthesia in children. Anesth Analg 2004; 98: 60–3
- 44 Malviya S, Voepel-Lewis T, Ramamurthi RJ, Burke C, Tait AR. Clonidine for the prevention of emergence agitation in young children: efficacy and recovery profile. *Paediatr Anaesth* 2006; 16: 554–9
- 45 Lankinen U, Avela R, Tarkkila P. The prevention of emergence agitation with tropisetron or clonidine after sevoflurane anesthesia in small children undergoing adenoidectomy. *Anesth Analg* 2006; 102: 1383-6
- 46 Tesoro S, Mezzetti D, Marchesini L, Peduto VA. Clonidine treatment for agitation in children after sevoflurane anesthesia. Anesth Analg 2005; 101: 1619–22
- 47 Kulka PJ, Bressem M, Tryba M. Clonidine prevents sevoflurane-induced agitation in children. Anesth Analg 2001; 93: 335–8

- 48 Guler G, Akin A, Tosun Z, Ors S, Esmaoglu A, Boyaci A. Single-dose dexmedetomidine reduces agitation and provides smooth extubation after pediatric adenotonsillectomy. *Paediatr Anaesth* 2005; 15: 762-6
- 49 Bock M, Kunz P, Schreckenberger R, Graf BM, Martin E, Motsch J. Comparison of caudal and intravenous clonidine in the prevention of agitation after sevoflurane in children. Br J Anaesth 2002; 88: 790-6
- **50** Saadawy I, Boker A, Elshahawy MA, et al. Effect of dexmedetomidine on the characteristics of bupivacaine in a caudal block in pediatrics. Acta Anaesthesiol Scand 2009; **53**: 251–6
- **51** Araki H, Fujiwara Y, Shimada Y. Effect of flumazenil on recovery from sevoflurane anesthesia in children premedicated with oral midazolam before undergoing herniorrhaphy with or without caudal analgesia. *J Anesth* 2005; **19**: 204–7
- **52** Davis PJ, Greenberg JA, Gendelman M, Fertal K. Recovery characteristics of sevoflurane and halothane in preschool-aged children undergoing bilateral myringotomy and pressure equalization tube insertion. *Anesth Analg* 1999; **88**: 34–8
- 53 Cohen IT, Joffe D, Hummer K, Soluri A. Ondansetron oral disintegrating tablets: acceptability and efficacy in children undergoing adenotonsillectomy. *Anesth Analg* 2005; 101: 59–63
- 54 Veyckemans F. Excitation phenomena during sevoflurane anaesthesia in children. Curr Opin Anaesthesiol 2001; 14: 339–43

- 55 Kain ZN, Caldwell-Andrews AA, Maranets I, et al. Preoperative anxiety and emergence delirium and postoperative maladaptive behaviors. Anesth Analg 2004; 99: 1648–54
- 56 Arai YC, Ito H, Kandatsu N, Kurokawa S, Kinugasa S, Komatsu T. Parental presence during induction enhances the effect of oral midazolam on emergence behavior of children undergoing general anesthesia. Acta Anaesthesiol Scand 2007; 51: 858-61
- 57 da Conceição MJ, Bruggemann DACD, Carneiro Leao C. Effect of an intravenous single dose of ketamine on postoperative pain in tonsillectomy patients. *Paediatr Anaesth* 2006; 16: 962–7
- 58 Erhan OL, Goksu H, Alpay C, Bestas A. Ketamine in post-tonsillectomy pain. Int J Pediatr Otorhinolaryngol 2007; 71: 735-9
- 59 Reimer EJ, Dunn GS, Montgomery CJ, Sanderson PM, Scheepers LD, Merrick PM. The effectiveness of clonidine as an analgesic in paediatric adenotonsillectomy. Can J Anaesth 1998; 45: 1162-7
- **60** Tripi PA, Palmer JS, Thomas S, Elder JS. Clonidine increases duration of bupivacaine caudal analgesia for ureteroneocystostomy: a double-blind prospective trial. *J Urol* 2005; **174**: 1081–3
- 61 Nishina K, Mikawa K. Clonidine in paediatric anaesthesia. Curr Opin Anaesthesiol 2002; 15: 309-16
- 62 Sikich N, Lerman J. Development and psychometric evaluation of the pediatric anesthesia emergence delirium scale. *Anesthesiology* 2004; 100: 1138–45