

Non-invasive myocardial work normal reference ranges in healthy children

Sabatino J.¹; Leo I.¹; Borrelli N.²; Avesani M.³; La Bella S.³; Josen M.²; Paredes J.²; Sirico D.³; Piccinelli E.²; Delle Donne G.²; Brida M.⁴; Indolfi C.¹; Fraisse A.²; Di Salvo G.³

¹Magna Graecia University of Catanzaro, Medical and Surgical Science, Catanzaro, Italy

²Royal Brompton Hospital, Pediatric Cardiology, London, United Kingdom of Great Britain & Northern Ireland

³University of Padua, Pediatric Cardiology, Padova, Italy

⁴University Hospital Centre Zagreb, Zagreb, Croatia

Funding Acknowledgements: Type of funding sources: None.

BACKGROUND: Myocardial work (MW) estimation by pressure-strain loops (PSL) allows a non-invasive evaluation of myocardial performance, as recently demonstrated in adult patients.

Aim of this study is to provide the reference values for global myocardial work index (MWI), constructive work (MCW), wasted work (MWW), and work efficiency (MWE) in a group of healthy children.

METHODS: Assessment of MW was performed using a commercially available software package (Echopac, GE). MW was measured from PSLs areas, derived from non-invasive LVP curves combined with strain acquired speckle tracking echocardiography (STE).

After calculating GLS, values of brachial blood pressure were inserted and the time of valvular events by echocardiography were indicated, then the software was able to measure non-invasive PSLs.

RESULTS: Two-dimensional (2D) standard and speckle-tracking echocardiography were performed in 90 healthy children (mean age 9.9 ± 4.9 [1-17] years, females: 57%) together with the assessment of MW by means of PSLs.

Mean ± standard deviation, 5° and 95° percentile values for global MWI, MCW, MWW, and MWE in the whole population were 1769 ± 254 mm Hg, (1354-2193); 2201 ± 290 mm Hg, (1657-2658); 78 ± 47 (29-163) mm Hg%; 96 ± 1.8 (92-99)%, respectively.

CONCLUSIONS: The assessment of MW is feasible in healthy children. This study provides useful 2-dimensional echocardiographic reference ranges for novel indices of non-invasive MW.