

The predictive value of sperm chromatin structure assay (SCSA) parameters for the outcome of intrauterine insemination, IVF and ICSI

M.Bungum^{1,4}, P.Humaidan¹, M.Spano², K.Jepson³, L.Bungum¹ and A.Giwercman³

¹Fertility Clinic, Viborg Hospital (Skive), Resenvej 25, DK 7800 Skive, Denmark, ²Section of Toxicology and Biomedical Sciences, BIOTEC-MED, ENEA Casaccia Research Center, Rome, Italy and ³Fertility Centre, Scanian Andrology Centre, Malmö University Hospital, Malmö, Sweden

⁴To whom correspondence should be addressed. E-mail: mona.bungum@sygehusviborg.dk

INTRODUCTION: Sperm chromatin integrity assessment has been suggested as a fertility predictor. The aim of this study was to examine the relationship between the results of sperm chromatin structure assay (SCSA) and the outcome of IVF, ICSI and intrauterine insemination (IUI). **METHODS:** A total of 306 consecutive couples undergoing assisted reproduction were included. IUI was performed in 131, IVF in 109 and ICSI in 66. SCSA results were expressed as DNA fractionation index (DFI) and highly DNA stainable (HDS) cell fractions. Reproductive outcome parameters were biochemical pregnancy (BP), clinical pregnancy (CP) and delivery (D). **RESULTS:** For IUI, the chance of pregnancy/delivery was significantly higher in the group with DFI $\leq 27\%$ and HDS $\leq 10\%$ than in patients with DFI $>27\%$ or HDS $>10\%$. The odds ratios (ORs) (95% confidence intervals) were 20 (2.3–117), 16 (1.9–137) and 14 (1.6–110) for BP, CP and D, respectively. No statistical difference between the outcomes of IVF versus ICSI was observed in the group with DFI $\leq 27\%$. In the DFI $>27\%$ group, however, the results of ICSI were significantly better than those of IVF. Comparing ICSI with IVF, the OR (95% CI) for BP was 26 (1.9–350). **CONCLUSIONS:** SCSA is a useful method for prediction of the outcome of assisted reproduction.

Key words: ART/DNA damage/pregnancy rate/sperm chromatin

Introduction

Sperm nuclear condensation is a complex process that involves a series of critical events including rearrangements of chromosomes, transition of DNA-binding proteins and alteration of transcription (Ward, 1994; Kramer and Krawetz, 1997). Disturbances in this process may have a negative impact on male fertility as the highly organized and compact sperm chromatin is known to have a crucial influence on the fertilizing process (Twigg *et al.*, 1998a,b; Evenson *et al.*, 1980, 1999, 2002; Larson *et al.*, 2000; Spano *et al.*, 2000; Braun, 2001; Agarwal and Said, 2003; Perreault *et al.*, 2003; Sakkas *et al.*, 2003). Infertile men are reported to have a higher fraction of sperm with chromatin defects and DNA breaks than fertile controls (Evenson *et al.*, 1980, 1999; Lopes *et al.*, 1998; Host *et al.*, 1999; Gandini *et al.*, 2000; Irvine *et al.*, 2000; Larson *et al.*, 2000; Carrell and Liu, 2001; Hammadeh *et al.*, 2001; Zini *et al.*, 2001, 2002; Sakkas *et al.*, 2002; Saleh *et al.*, 2002, 2003; Erenpreisa *et al.*, 2003; Muratori *et al.*, 2003). Moreover, sperm DNA defects may have a possible negative impact on the outcome of assisted reproductive techniques (ARTs) (Lopes *et al.*, 1998; Larson *et al.*, 2000; Evenson and Jost, 2000; Duran *et al.*, 2002; Morris *et al.*, 2002; Tomsu *et al.*, 2002; Benchaib *et al.*, 2003; Larson-Cook *et al.*, 2003; Saleh

et al., 2003). Furthermore, it has been questioned whether or not ART is able to compensate for poor DNA quality (Twigg *et al.*, 1998b; Evenson *et al.*, 1999; Larson *et al.*, 2000; Larson-Cook *et al.*, 2003).

The sperm chromatin structure assay (SCSA), first described by Evenson *et al.* (1980), is a flow cytometric (FCM) technique which identifies the spermatozoa with abnormal chromatin packaging defined as susceptibility to acid-induced DNA denaturation *in situ*. Acridine orange (AO) staining is used, after a low pH challenge, to distinguish between denaturated (red fluorescence = single stranded) and native (green fluorescence = double stranded) DNA regions in sperm chromatin, the former being a result of DNA breaks and/or of derailments in protamine quantity and composition and/or of an insufficient level of disulfide groups (Sakkas *et al.*, 1999; Irvine *et al.*, 2000; Aitken and Krausz, 2001). The level of DNA breaks is conveniently expressed by the DNA fragmentation index (DFI) (Larson *et al.*, 2000; Evenson *et al.*, 2002; Larson-Cook *et al.*, 2003), which is the ratio of red to total (red plus green) fluorescence intensities in the flow cytometric analysis. In two independent studies, DFI levels >30 – 40% were incompatible with fertility *in vivo*, whatever sperm concentration, morphology and motility (Evenson *et al.*, 1999; Spano

Table I. Background characteristics for 306 patients undergoing IUI or IVF/ICSI

	IUI		IVF/ICSI	
	DFI <27%	DFI >27%	DFI <27%	DFI >27%
Couples included (<i>n</i>)	108	23	140	35
Female age, median (range) (years)	30.2 (24.4–40.4)	32.1 (25.7–37.6)	31.2 (22.3–40.0)	32.1 (25.2–39.6)
BMI, median, (range) (kg/m ²)	23.8 (16.5–30.0)	23.6 (18.1–30.0)	23.7 (17.3–30.0)	22.7 (18.6–30.0)
Couples undergoing first or second ART treatment (%)	76.0	70.0	71.4	60.0
bFSH, median (range) (IU/l)	6.8 (1.1–12.0)	7.0 (2.4–12.0)	6.6 (1.7–12.0)	6.8 (2.1–12.0)
Male age, median (range) (years)	31.1 (24.3–55.3)	32.4 (27.0–56.7)	33.1 (23.5–47.6)	34.9 (28.0–49.5)

et al., 2000). Furthermore, although based on limited numbers of patients, no pregnancy has been reported after *in vitro* ART procedures, both standard IVF and ICSI, when the DFI in raw semen was >27% (Larson *et al.*, 2000; Larson-Cook *et al.*, 2003; Saleh *et al.*, 2003). As a consequence, the SCSA method has been recommended for routine use in ART programmes in order to predict a couple's probability of obtaining a pregnancy (Evenson and Jost, 2000). Before recommending SCSA as routine prior to ART, however, the predictive value of this method needs to be tested in larger scale studies where other parameters of importance for the success of the treatment (e.g. sperm concentration and motility, age of the female and number of previous unsuccessful ART treatments) are taken into consideration.

The present study was set up to examine the relationship between sperm chromatin defects evaluated by the SCSA and the outcome of IVF, ICSI and intrauterine insemination (IUI), respectively.

Materials and methods

Patients and study design

The study was based on a cohort of consecutive infertile couples undergoing ART treatment at the Fertility Clinic, Skive Hospital during the period April 2002–March 2003. Only cycles with a sperm concentration of at least 1×10^6 /ml in raw semen were included in the study. All patients were to take part in our ART programme. The inclusion criteria for the female partner was: (i) female age ≤ 40 years; (ii) female body mass index (BMI) < 30 kg/m²; and (iii) female baseline FSH < 12 IU/l. For each couple, only one treatment cycle during the study period was included in the analysis. The study was approved by the Ethics Committee of Viborg County.

A total of 306 patients were included in the study: 131 couples underwent IUI, 109 IVF and 66 ICSI. All demographic data including female BMI, pre-treatment levels of FSH and number of previous ART treatments for all the three categories of treatments are given in Table I.

Sperm collection and semen analysis

Semen samples were collected by masturbation at the day of oocyte retrieval or insemination. After semen analysis, within 1 h from the time of ejaculation, 100 μ l of the raw semen sample was frozen at -80°C for later SCSA analysis. Semen analysis was performed according to the World Health Organization guidelines (World Health Organization, 1999).

Sperm chromatin structure assay (SCSA)

Sperm DNA damage was measured by SCSA following the procedure as described elsewhere (Spano *et al.*, 2000; Evenson *et al.*, 2002). On the day of analysis, the samples were quickly thawed and analysed immediately. A cell suspension ($1-2 \times 10^6$) was treated with a low-pH (pH 1.2) detergent solution containing 0.1% Triton X-100, 0.15 mol/l NaCl and 0.08 mol/l HCl for 30 s, followed by staining with 6 mg/l purified AO (Molecular Probes, Eugene, OR) in a phosphate-citrate buffer, pH 6.0. Cells were analysed using a FACSort flow cytometer (Becton Dickinson, San Jose, CA), equipped with an air-cooled argon ion laser. A total of 5000 events were accumulated for each measurement. Under these experimental conditions, when excited with a 488 nm light source, AO intercalated with double-stranded DNA emits green fluorescence and AO associated with single-stranded DNA emits red fluorescence. Thus, sperm chromatin damage can be quantified by the FCM measurements of the metachromatic shift from green (native, double-stranded DNA) to red (denatured, single-stranded DNA) fluorescence and displayed as red (fragmented DNA) versus green (DNA stainability) fluorescence intensity cytogram patterns. For the flow cytometer set-up and calibration, reference samples were used from a normal donor ejaculate sample retrieved from the laboratory repository. Off-line analysis of the flow cytometric data was carried out using dedicated software (List View, Phoenix Flow Systems, San Diego, CA). Computer gates are used to determine the proportion of spermatozoa with increased levels of red fluorescence (denatured single-stranded DNA) and green fluorescence (native double-stranded DNA). We have expressed the extent of DNA denaturation in terms of the DFI, which is the ratio of red to total (red plus green) fluorescence intensity, i.e. the level of denatured DNA over the total DNA (Evenson *et al.*, 2002). The DFI value was calculated for each sperm cell in a sample, and the resulting DFI frequency profile was obtained (Figure 1, upper panel). Most sperm form a unimodal distribution representing the normal population of sperm with no detectable DNA damage. Sperm with higher red fluorescence, falling in the histogram area beyond the curve of normal sperm, represent the population of abnormal sperm with detectable DFI. Additionally, we have also considered the fraction of high DNA stainable (HDS) cells. HDS represents another distinct population in semen that characterizes immature spermatozoa with incomplete chromatin condensation. The percentage of HDS cells was calculated by setting an appropriate gate on the bivariate cytogram (Figure 1, lower panel) and considering as immature spermatozoa those events which exhibit a green fluorescence intensity higher than the upper border of the main cluster of the sperm population with a non-detectable DFI.

All SCSA measurements have been performed on raw semen cryopreserved prior to the preparation for IUI, IVF or ICSI. SCSA studies addressing the outcome of ART were based predominantly on




Figure 1. Upper panel: frequency distribution histogram of the DNA fragmentation index (DFI). The area located to the right of the main peak (which includes normal sperm with non-detectable DFI) represents the region where the sperm with detectable levels of fragmented DNA accumulate. Lower panel: SCSA scattergram of red (fragmented DNA, x-axis) versus green (double-stranded DNA, y-axis) fluorescence intensity of the same semen sample. Cytogram dots represent single spermatozoa with dual-parameter green and red fluorescence values acquired at 10-bit resolution (1024 channels) on the flow cytometer. Debris (bottom left corner) were excluded from the analysis. The region for calculating the fraction of immature sperm with high DNA stainability (HDS) is indicated by the box. The long dashed line indicates the threshold for HDS (channels 550 on the y-axis).

data derived from raw semen (Larson *et al.*, 2000; Larson-Cook *et al.*, 2003; Saleh *et al.*, 2003).

Intrauterine insemination (IUI), hormonal treatment and sperm preparation

All IUI patients had a diagnosis of unexplained infertility. Normal tubal patency was confirmed by either a hysterosalpingography (HSG) or by laparoscopy. IUI was only performed in cases with at least 2.5×10^6 spermatozoa per ml after gradient centrifugation. The IUI patients were stimulated with clomiphene (Pergotim; Serono Nordic, Copenhagen, Denmark) on cycle days 3–7 and 75 IU of recombinant FSH (rec-FSH) (Gonal F; Serono Nordic) on cycle days 7–9. When 1–2 follicles had reached a diameter of ≥ 17 mm, 10 000 IU of HCG (Profasi; Serono Nordic) was administered to induce final follicular maturation. After 36 h, a fresh semen sample was prepared by a standard density gradient centrifugation method. PureSperm, 40 and

80% (Nidacon Ltd, Gothenburg, Sweden) diluted in SpermRinse™ (Vitrolife, Gothenburg, Sweden) was used for sperm preparation. The washing procedure and dilution was performed in IVF-100™ (Vitrolife, Gothenburg, Sweden). A 1 ml aliquot of the prepared semen sample was used for insemination with a Cashmed catheter (catalogue no. IUI18; Cashmed, Surrey, UK). No luteal phase support was given.

IVF and ICSI

Hormonal treatment. IVF/ICSI patients underwent pituitary down-regulation and hormonal treatment as previously described (Bungum *et al.*, 2003). In brief, patients were downregulated with a GnRH agonist (Suprefact, Hoechst, Hørsholm, Denmark), using a long or a short protocol. Ovarian stimulation was performed with rec-FSH (Gonal F; Serono Nordic) (using an individualized dose of between 100 and 375 IU s.c. per day according to age, ovarian volume, baseline FSH and BMI). The ovarian response was monitored by ultrasound examination starting on day 8 of stimulation, and the dose of rec-FSH administered was adjusted if necessary.

Oocyte retrieval and sperm preparation. When at least one follicle had reached a diameter of ≥ 17 mm, 10 000 IU of HCG (Profasi; Serono Nordic) was administered to induce final follicular maturation. Oocyte retrieval was performed 35 h later by vaginal ultrasound-guided follicle aspiration. Gamete™ (Vitrolife, Gothenburg, Sweden), a HEPES-buffered medium, was used to rinse the oocytes.

A standard density gradient centrifugation method, PureSperm, 45 and 90% (Nidacon Ltd) diluted in SpermRinse™ (Vitrolife), was used for sperm preparation. The washing procedure and dilution was performed in IVF-100™ (Vitrolife).

IVF. In cases of conventional IVF, spermatozoa with a final concentration of 150 000/ml were added to the oocytes. After incubation in a humidified incubator with a gas phase of 6% CO₂, 5% O₂ and 89% N₂ for 90 min, the oocytes were washed three times before further culture in IVF-100™ until time of assessment for fertilization.

ICSI. In cases of ICSI, denudation of cumulus cells was performed by exposure of the oocytes to HYASE™ (Vitrolife) for a maximum of 30 s. Denudation of cumulus cells was performed by the use of glass denuding pipettes (SweMed Lab, Billdal, Sweden) immediately before injection. The oocytes were washed four times in Gamete™ after denudation. ICSI was performed in Gamete™ covered by OVOIL™ (Vitrolife) using commercially available ICSI pipettes (Cook, Brisbane, Australia). A 5 µl ultra-microdroplet of poly-vinylpyrrolidone (PVP), ICSI-100™ (Vitrolife) was spread out in a thin layer on the Petri dish. A 1 µl aliquot of the sperm preparation was introduced to the centre of the PVP droplet. After injection, the oocytes were washed twice in G.1.2™ (Vitrolife) medium before further culture in G.1.2™

Culture media and procedure for culturing

IVF-100™ was used in the first step of culture, from oocyte retrieval to fertilization, after which the oocytes were rinsed twice in G1.2™ and cultured in G1.2™ until the morning of day 3. In cases of blastocyst culture, G2.2™ (Vitrolife) was used from the 8-cell stage on day 3 until embryo transfer.

A maximum of six fertilized oocytes were cultured in 20 µl media droplets under oil (OVOIL™). In cases with fewer than six oocytes, 10 µl droplets were used. A gas phase of 6% CO₂, 5% O₂ and 89% N₂ was used in a humidified incubator.

Table II. Comparison of reproductive outcomes in the 306 couples undergoing assisted reproduction, according to the treatment type and the DFI

	IUI			IVF/ICSI		
	DFI <27%	DFI >27%	P-value	DFI <27%	DFI > 27%	P-value
Cycles started, <i>n</i>	108	23	–	140	35	–
Embryo transfer (ET), <i>n</i> (% per started cycle)	–	–	–	123 (87.9%)	34 (97.1%)	NS
Biochemical pregnancies, <i>n</i> (% per ET)	24 (22.2%)	1 (4.3%)	0.07	63 (51.2%)	16 (47.1%)	NS
Clinical pregnancies, % per ET or IUI	22 (20.2%)	1 (4.5%)	0.20	47 (38.2%)	13 (38.2%)	NS
Implantation rate, <i>n</i> (%)	–	–	–	71/217 (32.7%)	18/63 (28.6%)	NS
Deliveries, % per started cycle	19 (17.6%) ^a	1 (4.5%)	NS	44 (31.4%)	12 (34.3%)	NS

^aOne ectopic pregnancy.

NS: $P > 0.05$.

Assessment of fertilization, embryo morphology classification, cryopreservation and embryo transfer

Fertilization was determined 18–20 h after the IVF/ICSI procedure. The oocytes were fertilized when two distinct pronuclei were visible. Cleavage and classification of morphology was assessed 24 h later by criteria set up by Ziebe S, Groendal ML, Agerholm I and Erb K (personal communication). Strict criteria for cryopreservation were used. Only embryos with at least seven blastomeres and <20% intracellular fragments were cryopreserved on day 3. On day 5, embryos were assessed according to scoring criteria for blastocysts (Gardner and Schoolcraft, 1999). Only expanded blastocysts were cryopreserved.

A maximum of two embryos were transferred on day 2, 3 or 5 after retrieval. Embryo transfer was performed in G2.2TM. All embryo transfers were performed with a Cook Soft 5000 catheter (Cook, Brisbane, Australia).

Luteal phase support and pregnancy test

IVF/ICSI patients received luteal phase support in the form of daily vaginal administration of micronized progesterone, either 400 mg twice a day (Cyklogest; Hoechst, Copenhagen, Denmark) or 90 mg once a day (Crinone 8%; Serono Nordic) starting on the day following oocyte retrieval and continuing until the day of the pregnancy test (i.e. day 12 after embryo transfer). A positive pregnancy test was defined by a plasma β HCG concentration >10 IU/l. A clinical pregnancy was defined as an intrauterine gestational sac with a heart beat 3 weeks after a positive HCG test. The implantation rate was calculated as the ratio of gestational sacs determined by ultrasound after 7 weeks in relation to the total number of embryos transferred. Birth or abortion was recorded for all patients.

Statistical methods

Statistical analysis was performed using the SPSS 11.0 software (SPSS Inc., Chicago, IL). To test inter-group differences in sperm concentration, percentage progressive motile sperm, DFI and HDS, the Mann–Whitney test was applied. Binary correlations were assessed by calculating Spearman's rho. The reproductive outcome measures were treated as discrete variables (\pm fertilization; \pm biochemical pregnancy; \pm clinical pregnancy; \pm delivery). Since a DFI of 27% previously was reported to be the cut-off level for achieving a pregnancy by *in vitro* ART (Larson-Cook *et al.*, 2003), the couples were grouped according to whether the DFI in raw semen sample was $\leq 27\%$ or $>27\%$. Furthermore, having assessed the results of HDS analysis in relation to the reproductive outcomes, a cut-off level of 10% was defined for this parameter. By the use of Fischer's exact test, the chance of a positive reproductive outcome was calculated for the different treatment groups (IUI, IVF and ICSI) divided according to the DFI and HDS as described above (DFI: $\leq 27\%$ and $>27\%$; HDS:

$\leq 10\%$ and $>10\%$). To compare the results of IVF and ICSI, in the group with DFI $>27\%$, odds ratios (ORs) and the 95% confidence intervals (95% CIs) were calculated by the use of binary logistic regression analysis. Fischer's exact test and OR calculations were also applied to compare the reproductive outcomes, i.e. biochemical and clinical pregnancy in two groups defined as follows: group 1: DFI $\leq 27\%$ and HDS $\leq 10\%$; and group 2: DFI $>27\%$ or HDS $>10\%$. A P -value ≤ 0.05 was considered as statistically significant.

Results

IUI

The median (range) age of women was 30.2 (24.4–40.4) years in the group with DFI $\leq 27\%$ and 32.1 (25.7–37.6) years in the group with DFI $>27\%$. The median (range) age of men was 31.1 (24.3–55.3) years in the group with DFI $\leq 27\%$, whereas the median (range) age was 32.4 (27.0–56.7) years in the group with DFI $>27\%$ (Table I). When comparing the groups with and without clinical pregnancy, there were no statistically significant differences regarding DFI (medians: 12.1 versus 15.7; $P = 0.14$) or HDS (medians: 6.3 versus 6.7; $P = 0.12$). With biochemical pregnancy as a grouping parameter, lower DFI and HDS values were found in couples with positive reproductive outcome, the difference not being statistically significant for DFI [12.9% (4.3–33.7) versus 15.4% (3.0–48.7); $P = 0.25$] but close to the level of statistical significance for HDS [6.2% (3.6–17.8) versus 6.7% (2.7–20.2); $P = 0.06$].

There was a statistically significant negative correlation between DFI and the percentage of progressively motile spermatozoa ($\rho = -0.25$; $P = 0.004$), but not between DFI and sperm concentration. HDS correlated in a negative manner with both sperm concentration ($\rho = -0.36$; $P < 0.0005$) and percentage progressively motile sperm ($\rho = -0.25$; $P = 0.004$).

Predictive value of DFI >27%. One biochemical pregnancy was observed in the 23 cycles with males having DFI $>27\%$ (4.3%), whereas in the $\leq 27\%$ group, 24 of the 108 cycles resulted in a positive pregnancy test (22.2%) ($P = 0.07$; Table II). The clinical pregnancy and delivery rates were 4.5 versus 20.2% ($P = 0.12$) and 4.5 versus 17.6% ($P = 0.20$), respectively. In the above-mentioned case where the DFI was $>27\%$, the actual value being 33.7%, there was a positive biochemical and subsequently even clinical pregnancy and delivery. When the cut-off level was increased to a DFI of 34%, there were no statistically significant differences with regard to any of the reproductive outcomes.

Predictive value of HDS >10%. Only one biochemical and clinical pregnancy, and birth was seen when the HDS exceeded the value of 10%. An HDS of 17.8% was found in the sample having a DFI of 33.7%, as mentioned above. There were 24 biochemical and 21 clinical pregnancies, respectively, among those 115 having a HDS of $\leq 10\%$. Nineteen of the clinical pregnancies resulted in a delivery. The difference between the two groups defined according to the HDS level was not statistically significant.

Predictive value of combined use of DFI and HDS. When the patients were divided into two groups, group 1, with semen samples with DFI $\leq 27\%$ and HDS $\leq 10\%$, and group 2, with either DFI $>27\%$ or HDS $>10\%$, 24 biochemical pregnancies, 21 clinical pregnancies and 19 deliveries were observed among the 95 couples in group 1, whereas only one of the 36 couples in group 2 experienced a biochemical as well as a clinical pregnancy and, finally, delivery. These differences in reproductive outcomes between group 1 and group 2 were statistically significant, the *P*-values being 0.002 for biochemical pregnancy, 0.008 for clinical pregnancy and 0.01 for delivery rate. The OR for biochemical pregnancy in group 1, with group 2 as reference, was 12 (95% CI 1.5–96). This value increased to 20 (95% CI 2.3–117) when sperm concentration, percentage progressive motile, female age and treatment number were included in the model. For clinical pregnancies, the ORs in group 1 were 9.9 (95% CI 1.3–77) and 16 (95% CI 1.9–137), without and including the above-mentioned covariates, respectively. For delivery as an outcome variable, the ORs were 8.7 (95% CI 1.1–68) and 14 (95% CI 1.6–110), respectively, with and without inclusion of sperm concentration, percentage progressively motile spermatozoa, female age and treatment number in the model.

IVF/ICSI

There was no difference in sperm concentration, percentage of progressively motile sperm, DFI or HDS when couples with positive and negative reproductive outcome were compared, regardless of whether biochemical or clinical pregnancy was used as a criterion. This lack of a statistically significant difference was true for the whole group of IVF/ICSI-treated couples and also when the two treatment types were analysed separately (data not shown).

Similarly to the findings in the IUI group, there was a statistically significant negative correlation between both DFI and HDS and the percentage of progressively motile sperm, and additionally between HDS and sperm concentration (data not shown).

There was a considerable overlap between DFI and HDS values when couples who achieved a pregnancy were compared with those who did not (Figure 2).

The likelihood of obtaining a clinical pregnancy was the same in the group of patients with DFI $>27\%$ (38.2%) and those with DFI $\leq 27\%$ (38.2%). The implantation rates were also comparable, 31.4 versus 34.3%. Forty-four women (31.4%) in the DFI $\leq 27\%$ group and 12 of the women (34.3%) in the group with DFI $>27\%$ (*P* = 0.8) all delivered




Figure 2. Box and whisker plots showing the distribution of DFI (white boxes) and HDS (grey boxes) values among couples undergoing ICSI or standard IVF. The couples are divided into two groups, according to whether or not they achieved pregnancy. The bars indicate median values, the boxes interquartile range and the whiskers 95% confidence intervals.

healthy children. Two pregnancies in the DFI $\leq 27\%$ group and one in the DFI $>27\%$ group were ectopic (Table III).

Predictive value of DFI $>27\%$. In the DFI $>27\%$ group, the results of ICSI were significantly better than those of standard IVF regarding biochemical (70.6 versus 22.2%; *P* = 0.007) and clinical pregnancies (52.9 versus 22.2%; *P* = 0.09). For delivery rates, the percentages were 47.1% in the ICSI and 22.2% in the standard IVF group (*P* = 0.16) (Table III), with an OR of 7.8 (95% CI 1.7–36) for a positive HCG test, 3.9 (95% CI 0.9–17.0) for clinical pregnancy and 3.1 (95% CI 0.72–13) for delivery rate. After inclusion of sperm concentration, percentage progressively motile sperm, female age and treatment number in the model, the ORs for successful reproductive outcome when ICSI was compared with standard IVF increased to 26 (95% CI 1.9–350), 7.7 (95% CI 0.84–70) and 7.0 (95% CI 0.75–65), with biochemical pregnancy, clinical pregnancy and delivery as dependent variables, respectively. Significantly higher DFI (*P* = 0.014) (medians 18.0 versus 16.61) and HDS (medians 7.3 versus 6.81) (*P* = 0.033) were found in the ICSI group compared with IVF.

Discussion

Two major conclusions can be drawn from this, to our knowledge, largest ever reported study on the predictive value of SCSA in relation to the outcome of IUI, IVF and ICSI. First, combining the two SCSA parameters, DFI and HDS gave a high predictive value regarding the outcome of IUI treatments. Secondly, contrary to previous reports (Larson *et al.*, 2000; Larson-Cook *et al.*, 2003; Saleh *et al.*, 2003), we found that a

Table III. Comparison of the reproductive outcomes between couples undergoing IVF and ICSI, allocated according to the DFI

	DFI <27%			DFI >27%		
	IVF	ICSI	<i>P</i> -value	IVF	ICSI	<i>P</i> -value
Cycles started, <i>n</i>	91	49	–	18	17	–
Embryo transfer (ET), <i>n</i> (% e started cycle)	82 (90.1%)	41 (83.7%)	NS	18 (100%)	17 (100%)	NS
Biochemical pregnancies, <i>n</i> IVF/ICSI (% per ET)	42 (51.2%)	21 (51.2%)	NS	4 (22.2%)	12 (70.6%)	0.007
Clinical pregnancies, % per ET	30 (36.6%)	17 (41.5%)	NS	4 (22.2%)	9 (52.9%)	NS
Implantation rate, <i>n</i> (%)	47/141 (33.3%)	24/76 (31.6%)	NS	6/31 (19.4%)	12/32 (37.5%)	NS
Deliveries, % pr started cycle	27 (29.7%) ^a	17 (34.7%)	NS	4 (22.2%)	8 (47.1%) ^b	NS

^aTwo ectopic pregnancies.

^bOne ectopic pregnancy.

NS: *P* > 0.05.

DFI level >27% is compatible with pregnancy and delivery after both IVF and ICSI. Thus it seems that *in vitro* ART is able to compensate for the impairment of sperm chromatin integrity, in particular if ICSI is chosen as a fertilization method.

In couples treated with IUI, clinical pregnancy was seen in only one out of 23 cases when the DFI exceeded 27% and/or a HDS >10% was found in the raw ejaculate prepared and used for insemination. The OR was 16 for clinical pregnancy and 14 for delivery when both these parameters were below the cut-off levels given above. This finding is in accordance with previously published *in vivo* studies in which the time to pregnancy (TTP) for couples with no previous infertility background has been studied (Evenson *et al.*, 1999; Spano *et al.*, 2000). These studies indicated a DFI level of 30–40% as a cut-off for no pregnancy. Two other recently published studies (Duran *et al.*, 2002; Saleh *et al.*, 2003) reached a similar conclusion. Saleh *et al.* (2003), using the SCSA, found significantly higher DFI levels in the couples who failed to obtain a pregnancy after IUI. This study was, however, based on 11 IUI couples only. Using the TUNEL assay indicative for DNA fragmentation, Duran *et al.* (2002) found that men with a DNA fragmentation level >12% had a decreased chance of fathering a pregnancy following IUI.

Apart from slightly lower (close to the level of statistical significance) HDS in men of those IUI-treated couples who achieved a biochemical pregnancy, there was no statistically significant difference in SCSA parameters between the groups with a negative and a positive reproductive outcome. The SCSA parameter, HDS, is believed to correlate to the level of immature spermatozoa and the extent of chromatin packaging (Evenson and Jost, 2000). Although previous studies indicated that high HDS could be a pejorative factor for *in vivo* fertility impairment (Evenson *et al.*, 1999, 2002), in the available materials, the associations did not reach a level of statistical significance. In our study, when a relatively large number of couples were included, an HDS of ≤10% was associated with a higher proportion of clinical pregnancies and births, although the statistical significance was borderline. It is therefore to be considered, as a novel finding, that the predictive value of SCSA increases when both DFI and HDS are used to define groups with high and low chances of obtaining a pregnancy after IUI. Recently, we have shown that both DFI and HDS are

independently and negatively associated with sperm motility (Giwerzman *et al.*, 2003). Motility is essential for fertilization *in vivo*. Impairment of the ability of spermatozoa to reach the ovum might at least partly explain the association between SCSA parameters and the probability of obtaining a pregnancy after IUI. An additional explanation could be the fact that spermatozoa with damaged DNA could be more likely to undergo apoptosis during the transport through the genital tract than spermatozoa with normal DNA integrity (Edwards, 2003).

The present study demonstrates that even men with high levels of DNA damage can become biological fathers, when DFI exceeded 27%; however, significantly higher clinical pregnancy rates (52.9 versus 22.2%) and delivery rates (47.1 versus 22.2%) were seen after ICSI compared with IVF. When DFI exceeded 27%, the OR for a positive reproductive outcome after ICSI compared with standard IVF was 8 for biochemical pregnancy, 4 for clinical pregnancy and 3 for delivery, respectively. These ORs increased after inclusion of sperm concentration, the percentage of progressively motile sperm, number of treatments and the age of the female as covariates. The increase in OR was statistically significant (95% CIs not including the value of 1) for biochemical pregnancy, but not for the two other reproductive outcomes, probably due to the limited number of couples included in this study. None of the previous studies dealing with this issue (Larson *et al.*, 2000; Larson-Cook *et al.*, 2003; Saleh *et al.*, 2003) have taken into consideration the age of the female and the number of previously failed ART treatments, parameters known to be associated with the success rate of the treatment. Therefore, our findings show that DFI as assessed by means of SCSA is an independent predictor of male fertility potential *in vitro*.

This is the first study showing that semen samples with high rates of DNA breaks as assessed by the SCSA method are more likely to result in pregnancy in ICSI procedures, compared with traditional IVF. The finding is novel in relation to the SCSA parameters, but confirms the results of a recent study by Host *et al.* (2000) using the TUNEL assay.

The safety of the ICSI procedure has been questioned (Schultz and Williams, 2002), and the findings from the present study may give further reason for concern. ICSI enables fertilization to take place in spite of severely compromised semen characteristics and damaged DNA. Aitken and Krausz (2001) proposed that sperm DNA damage is promutagenic and

can give rise to mutations after fertilization, as the oocyte attempts to repair DNA damage prior to the initiation of the first cleavage. Mutations occurring at this point will be fixed in the germline and may be responsible for the induction of pathology, such as infertility (Aitken and Krausz, 2001), childhood cancer in the offspring (Fraga *et al.*, 1996; Ji *et al.*, 1997; Aitken and Krausz, 2001) and imprinting diseases (Cox *et al.*, 2002; DeBaun *et al.*, 2003). So far, however, follow-up studies of children born after ICSI compared with children born after conventional IVF have not been conclusive regarding the risks of congenital malformations and health problems in general (Bonduelle *et al.*, 1996, 1998, 2003; Kurinczuk and Bower, 1997; Sutcliffe *et al.*, 1999; Wennerholm and Berg, 2000; Hansen *et al.*, 2002). It has been reported previously that the highest level of sperm DNA damage usually is found in men with the poorest semen quality, i.e. ICSI candidates (Aitken and Krausz, 2001). In the present study, the men in the ICSI group had significantly higher DFI levels compared with the men in the IVF group (median 18.0 versus 15.0%, respectively). It gives reason for concern that the most efficient ART techniques are used to treat males with the highest level of sperm DNA damage. SCSA parameters, however, refer to the DNA status of the whole cell population, not excluding the presence of a subpopulation with no significant DNA damage. Host *et al.* (2000), suggested that the technician who performs ICSI, attempts to select spermatozoa with normal morphology and thereby reduces the risk of introducing spermatozoa with strand breaks. This statement may, however, be questioned, since the traditional sperm parameters such as sperm count, motility and morphology have been proven to be poorly correlated to the SCSA parameters (Evenson *et al.*, 1991, 1999; Spano *et al.*, 1998, 2000). A future challenge will be to develop methods to identify and select spermatozoa with intact DNA during the ICSI procedure.

We selected a DFI level of 27% as a cut-off for calculation of ORs in relation to the reproductive outcome. This level was defined previously in reports where the SCSA measurement was performed in other laboratories. One could ask whether each laboratory performing the SCSA analysis should have its own reference levels or not. Apart from being subject to a very limited intra-laboratory variation (Giwerzman *et al.*, 1999), however, the SCSA analysis was shown to be very robust to variation between laboratories. In an external quality control based on >180 samples, a high ($\rho = 0.8$) correlation was found between the values obtained by our laboratory and those from a control laboratory. Furthermore, not only was there a high level of correlation between the results reported by two independent laboratories that strictly followed the SCSA protocol, but the absolute DFI values obtained at two different places, using different equipment, did not on average differ by >1% (Giwerzman *et al.*, 2003). Furthermore, in the present study, viewing the distribution of DFI values between those who became pregnant and those who did not, 27% was confirmed to be an appropriate cut-off level.

The results from this study confirm previous reports indicating a low chance of obtaining *in vivo* pregnancy when the DFI exceeds the level of 27%. Contrary to previous reports, the present study has shown that a DFI level >27% is not

incompatible with pregnancy and delivery following IVF and ICSI. ICSI seems to be the most efficient method in cases with high levels of sperm DNA breaks. Additional large-scale trials are needed to confirm the results of the present study and to investigate further the value of SCSA in relation to other available pregnancy predictors *in vivo* or *in vitro*. Methods for selection of spermatozoa with intact DNA should be developed in order to avoid the potential risk of using gametes with chromatin breaks for ICSI treatment.

Acknowledgements

The assistance of the staff of the clinic, in particular Mrs Anette Lybye Poulsen, Mrs Betina Boel Povlsen, Mrs Kirsten Madsen, Mrs Pia Elliman Jensen and Mrs Henni Blaaberg, is gratefully acknowledged. This study was supported by grants from the Swedish Research Council (grant no.521-2002-3907) University Hospital MAS Foundation in Malmö, Crafoordska Fund, Ove Tulefjords Fund, Swedish governmental funding for clinical research and the Foundation for Urological Research.

References

- Agarwal A and Said T (2003) Role of sperm chromatin abnormalities and DNA damage in male infertility. *Hum Reprod Update* 9,331–345.
- Aitken RJ and Krausz C (2001) Oxidative stress, DNA damage and the Y chromosome. *Reproduction* 122,497–506.
- Benchab M, Braun V, Lornage J, Hadj S, Salle B, Lejeune H and Guerin JF (2003) Sperm DNA fragmentation decreases the pregnancy rate in an assisted reproductive technique. *Hum Reprod* 18,1023–1028.
- Bonduelle M, Legein J, Buysse A, Van Assche E, Wisanto A, Devroey P, Van Steirteghem AC and Liebaers I (1996) Prospective follow-up study of 423 children born after intracytoplasmic sperm injection. *Hum Reprod* 11,1558–1564.
- Bonduelle M, Joris H, Hofmans K, Liabears I and Van Steirteghem AC (1998) Mental development of 201 ICSI children at 2 years of age. *Lancet* 351,1553.
- Bonduelle M, Ponjaert I, Steirteghem AV, Derde MP, Devroey P and Liebaers I (2003) Developmental outcome at 2 years of age for children born after ICSI compared with children born after IVF. *Hum Reprod* 18,342–350.
- Braun ER (2001) Packaging paternal chromosomes with protamine. *Nature Genet* 28,10–12.
- Bungum M, Bungum L, Humaidan P and Yding-Andersen C (2003) Day 3 versus day 5 embryo transfer; a prospective randomized study. *Reprod Biomed Online* 7,98–104.
- Carrell DT and Liu L (2001) Altered protamine 2 expression is uncommon in donors of known fertility, but common among men with poor fertilizing capacity, and may reflect other abnormalities of spermiogenesis. *J Androl* 22,604–610.
- Cox GF, Burger J, Lip V, Mau UA, Sperling K, Wu BL and Horsthemke B (2002) Intracytoplasmic sperm injection may increase the risk of imprinting defects. *Am J Hum Genet* 71,162–164.
- DeBaun MR, Niemitz EL and Feinber AP (2003) Association of *in vitro* fertilization with Beckwith–Wiedemann syndrome and epigenetic alterations of LIT1 and H19. *Am J Hum Genet* 72,156–160.
- Duran EH, Morshedi M, Taylor S and Oehninger S (2002) Sperm DNA quality predicts intrauterine insemination outcome—a prospective cohort study. *Hum Reprod* 17, 3122–3128.
- Edwards RG (2003) New outlooks on IVF spermatozoa. *Reprod Biomed Online* 5,89–91.
- Erenpreisa J, Erenpreiss J, Freivalds T, Slaidina M, Krampe R, Butikova J, Ivanov A and Pjanova D (2003) Toluidine blue test for sperm DNA integrity and elaboration of image cytometry algorithm. *Cytometry* 52A,19–27.
- Evenson DP and Jost LK (2000) Sperm chromatin structure assay is useful for fertility assessment. *Methods Cell Sci* 22,169–189.
- Evenson DP, Darzynkiewicz Z and Melamed MR (1980) Relation of mammalian sperm chromatin heterogeneity to fertility. *Science* 210,1131.
- Evenson DP, Jost LK, Baer RK, Turner TW and Schrader SM (1991) Individuality of DNA denaturation patterns in human sperm measured by the sperm chromatin structure assay. *Reprod Toxicol* 5,115–125.

- Evenson DP, Jost LK, Marshall D, Zinaman MJ, Clegg E, Purvis K, de Angelis P and Clausen OP (1999) Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum Reprod* 14,1039–1049.
- Evenson DP, Larson KL and Jost LK (2002) Sperm chromatin structure assay: its clinical use for detecting sperm DNA fragmentation in male infertility and comparisons with other techniques. *J Androl* 23,25–43.
- Fraga CG, Motchnik PA, Wyrobek AJ, Rempel DM and Ames BN (1996) Smoking and low antioxidant levels increase oxidative damage to sperm DNA. *Mutat Res* 351,199–203.
- Gandini L, Lombardo F, Paoli D, Caponecchia L, Familiari G, Verlengia C, Dondero F and Lenzi A (2000) Study of apoptotic DNA fragmentation in human spermatozoa. *Hum Reprod* 15,830–839.
- Gardner DK and Schoolcraft W (1999) *In vitro* culture of human blastocysts. In Jansen R and Mortimer D (eds), *Towards Reproductive Certainty: Infertility and Genetics Beyond 1999*. Parthenon Press, Carnforth, pp. 378–388.
- Giwerzman A, Spanò M, Lahdetie J and Bonde JP (1999) Quality assurance of semen analysis in multicenter studies. *Asclepios. Scand J Work Environ Health* 25, Suppl 1,23–25.
- Giwerzman A, Richthoff J, Hjöllund H, Bonde JP, Jepson K, Frohm B and Spanò M. (2003) Correlation between sperm motility and sperm chromatin structure assay parameters. *Fertil Steril* 80,1404–1412.
- Hammadeh ME, Zeginiadov T, Rosenbaum P, Georg T, Schmidt W and Strehler E (2001) Predictive value of sperm chromatin condensation (aniline blue staining) in the assessment of male fertility. *Arch Androl* 46,99–104.
- Hansen M, Kurinczuk JJ, Bower, C and Webb S (2002) The risk of major birth defects after intracytoplasmic sperm injection and *in vitro* fertilization. *N Engl J Med* 346,725–730.
- Host E, Lindenberg S, Kah JA and Christensen F (1999) DNA strand breaks in human sperm cells: a comparison between men with normal and oligozoospermic sperm samples. *Acta Obstet Gynecol Scand* 78,336–339.
- Host E, Lindenberg S and Smidt-Jensen S (2000) The role of DNA strand breaks in human spermatozoa used for IVF and ICSI. *Acta Obstet Gynecol Scand* 79,599–563.
- Irvine S, Twigg JP, Gordon EL, Fulton N, Milne PA and Aitken RJ (2000) DNA integrity in human spermatozoa: relationships with semen quality. *J Androl* 31,33–44.
- Ji, BT, Shu XO, Linet MS, Zheng W, Wacholder S, Ying DM and Jin F (1997) Paternal cigarette smoking and the risk of childhood cancer among offspring of non-smoking mothers. *J Natl Cancer Inst* 89,238–244.
- Kramer JA and Krawetz SA (1997) RNA in spermatozoa: implications for the alternative haploid genome. *Mol Hum Reprod* 3,473–478.
- Kurinczuk JJ and Bower C (1997) Birth defects in infants conceived by intracytoplasmic sperm injection: an alternative interpretation. *Br Med J* 315,1260–1265.
- Larson KL, De Jonge CJ, Barnes AM, Jost LK and Evenson DP (2000) Sperm chromatin structure assay parameters as predictors of failed pregnancy following assisted reproductive techniques. *Hum Reprod* 15,1717–1722.
- Larson-Cook KL, Brannian JD, Hansen KA, Kaspersen KM, Aarnold ET and Evenson DP (2003) Relationship between the outcomes of assisted reproductive techniques and sperm DNA fragmentation as measured by the sperm chromatin structure assay. *Fertil Steril* 80,895–902.
- Lopes S, Sun JG, Juriscova A, Meriano J and Casper RF (1998) Sperm deoxyribonucleic acid fragmentation is increased in poor-quality semen samples and correlates with failed fertilization in intracytoplasmic sperm injection. *Fertil Steril* 69,528–532.
- Morris ID, Iltot S, Dixon L and Brison DR (2002) The spectrum of DNA damage in human sperm assessed by single cell gel electrophoresis (Comet assay) and its relationship to fertilization and embryo development. *Hum Reprod* 17,990–998.
- Muratori M, Maggi M, Spinelli S, Filimberti E, Forti G and Baldi E (2003) Spontaneous DNA fragmentation in swim-up selected human spermatozoa during long term incubation. *J Androl* 24,253–262.
- Perreault SD, Aitken RJ, Baker HW, Evenson DP, Huszar G, Irvine DS, Morris ID, Morris RA, Robbins, WA, Sakkas D *et al.* (2003) Integrating new tests of sperm genetic integrity into semen analysis: breakout group discussion. *Adv Exp Med Biol* 518,253–268.
- Sakkas D, Manethoz E, Manicardi G, Bizzarro D, Bianchi PG and Bianchi U (1999) Origin of DNA damage in ejaculated human spermatozoa. *Rev Reprod* 4,31–37.
- Sakkas D, Moffatt O, Manicardi GC, Mariethoz E, Tarozzi N and Bizzarro D (2002) Nature of DNA damage in ejaculated human spermatozoa and the possible involvement of apoptosis. *Biol Reprod* 66,1061–1067.
- Sakkas D, Manicardi GC and Bizzarro D (2003) Sperm nuclear DNA damage in the human. *Adv Exp Med Biol* 518,73–84.
- Saleh RA, Agarwal A, Nelson DR, Nada EA, El-Tonsy MH, Alvarez JG, Thomas AJ and Sharma RK (2002) Increased sperm nuclear DNA damage in normozoospermic infertile men: a prospective study. *Fertil Steril* 78,313–318.
- Saleh RA, Agarwal A, Nada EA, El-Tonsy MH, Sharma RK, Meyer A, Nelson DR and Thomas AJ (2003) Negative effects of increased sperm DNA damage in relation to seminal oxidative stress in men with idiopathic and male factor infertility. *Fertil Steril* 79, Suppl 3,1597–605.
- Schultz RM and Williams CJ (2002) The science of ART. *Science* 296,2188–2190.
- Spano M, Bonde JP, Hjöllund HI, Kolstad HA, Cordelli E and Leter G (2000) Sperm chromatin damage impairs human fertility. The Danish First Pregnancy Planner Study Team. *Fertil Steril* 73,43–50.
- Spano M, Kolstad AH, Larsen SB, Cordelli E, Leter G, Giwerzman A, Bonde JP (1998) The applicability of the flow cytometric sperm chromatin structure assay in epidemiological studies. *Asclepios. Hum Reprod* 13,2495–2505.
- Tomsu M, Sharma V and Miller D (2002) Embryo quality and IVF treatment outcomes may correlate with different sperm comet assay parameters. *Hum Reprod* 17,1856–1862.
- Sutcliffe AG, Saunders K, Thornton S, Grudzinskas JG and Lieberman, BA (1999) Children born after intracytoplasmic sperm injection: population control study. *Br Med J* 318,704–705.
- Twigg J, Fulton N, Gomez E, Irvine DS and Aitken RJ (1998a) Analysis of the impact of intracellular reactive oxygen species generation on the structural and functional integrity of human spermatozoa: lipid peroxidation, DNA fragmentation and effectiveness of antioxidants. *Hum Reprod* 13,1429–1436.
- Twigg JP, Irvine DS and Aitken RJ (1998b) Oxidative damage to DNA in human spermatozoa does not preclude pronucleus formation in intracytoplasmic sperm injection. *Hum Reprod* 13,1864–1871.
- Ward DC (1994) The structure of the sleeping genome: implications of sperm DNA organization for somatic cells. *J Cell Biochem* 55,77–82.
- Wennerholm UB and Bergh C (2000) Obstetric outcome and follow-up of children born after *in vitro* fertilization (IVF). *Hum Fertil* 3,52–64.
- World Health Organization. (1999) *WHO Laboratory Manual for the Examination of Human Sperm and Sperm–Cervical Mucus Interaction*. Cambridge University Press.
- Zini A, Bielecki R, Phang D and Zenzes MT (2001) Correlations between two markers of sperm DNA denaturation and DNA fragmentation, in fertile and infertile men. *Fertil Steril* 75,674–677.
- Zini A, Fischer MA, Sharir S, Shayegan B, Phang D and Jarvi K (2002) Prevalence of abnormal sperm DNA denaturation in fertile and infertile men. *Urology* 60,1069–1072.

Submitted on January 9, 2004; accepted on March 31, 2004