

The distribution of chondrichthyans along the northern coast of Norway

Tom Williams, Kristin Helle, and Michaela Aschan

Williams, T., Helle, K., and Aschan, M. 2008. The distribution of chondrichthyans along the northern coast of Norway. – ICES Journal of Marine Science, 65: 1161–1174.

The relationship between temperature, latitude, and depth and the distribution and relative abundance of chondrichthyans along the northern coast of Norway was examined based on catches made in scientific trawls north of 62°N from 1992 to 2005. It appears that *Chimaera monstrosa*, *Etmopterus spinax*, *Squalus acanthias*, and *Galeus melastomus* were more abundant in the south, and *Amblyraja radiata* more common in the north. Between 1992 and 2005, the distribution and relative abundance did not appear to change significantly, although average water temperatures rose during the period. Current fishing levels do not appear to be impacting the populations of the more common species, but the status of species rarely found in the survey catches is unclear.

Keywords: abundance, bottom trawl survey, Chondrichthyes, distribution, elasmobranchs, Norway, shark, skate.

Received 4 February 2008; accepted 16 May 2008; advance access publication 27 June 2008.

T. Williams and M. Aschan: Norwegian College of Fishery Science, University of Tromsø, Breivika, N-9037 Tromsø, Norway. K. Helle: Institute of Marine Research, N-5817 Bergen, Norway. Correspondence to M. Aschan: tel: +47 776 46953; fax: +47 776 46020; e-mail: michaela.aschan@nfh.uit.no.

Introduction

There is increased awareness that stocks of chondrichthyans are particularly susceptible to overexploitation as either target or bycatch species. As large-bodied species with few natural predators when fully mature, they have not evolved strategies to withstand rapid changes in mortality (Stevens *et al.*, 2000). To varying degrees, all these species are slow-growing, have a long lifespan and low fecundity, and attain sexual maturity at a late age (Holden, 1977; Walker and Heessen, 1996). Population growth is influenced by juvenile survivorship and age at maturity (Sosebee, 2005). Although theoretical mechanisms of density-dependence have been looked at, empirical evidence of chondrichthyans (elasmobranchs) is limited and often confounded by methodological issues (Ellis *et al.*, 2008). Their limited ability to compensate for being depleted has often been exemplified by the poor record of sustainability by fisheries that have targeted them (Stevens *et al.*, 2000) and by elasmobranch stocks that have declined either unnoticed or unchecked. The reasons for these stock declines have been described by a number of authors (Bonfil, 1994; Dulvy *et al.*, 2000; Stevens *et al.*, 2000; ICES, 2006).

Fisheries research has traditionally focused on the more commercially important teleost and shellfish species, and little research has been undertaken on chondrichthyans. Catch and landings data from commercial fisheries are often poor because of a general lack of species-specific recording (Johnston *et al.*, 2005), and bycatch data only became more available recently. As a result, even the most basic data are unavailable for quantitative studies of the stock status of most chondrichthyans, including those in the Northeast Atlantic (ICES, 2006). The uncertainties in historical total landings and bycatch data attributable to the common practice of recording catches generically, e.g. as “dogfish and hounds”,

rather than by species, make fishery-independent data from surveys an important source for studying the distribution and relative abundance of chondrichthyans. To varying degrees, research has been undertaken throughout much of the ICES Area (Walker and Hislop, 1998; Daan *et al.*, 2005; Ellis *et al.*, 2005a), and studies have been conducted along the coast of southwestern Norway (Skjæraasen and Bergstad, 2000, 2001), and northwards into the Barents Sea (Dolgov, 1997, 2006; Dolgov *et al.*, 2004, 2005a). However, the chondrichthyan species along the northern coast of Norway have received little attention and are poorly understood.

Norwegian fisheries targeted Greenland shark (*Somniosus microcephalus*) until 1960 and basking shark (*Cetorhinus maximus*) until 2006 (ICES, 2006). Other chondrichthyans have not been and are not currently targeted by commercial fisheries operating along the northern coast of Norway, but are taken as bycatch in the coastal fishery. The coastal fleet consists of vessels using a variety of gears, including gillnet, longline, trawl, Danish seine, handline, and pots. Gillnet and longline fisheries targeting demersal fish (e.g. cod, *Gadus morhua*, and haddock, *Melanogrammus aeglefinus*) generate the bulk of the chondrichthyan bycatch (Table 1), and management strategies are in place to minimize the bycatch of undersized commercial species, though not relating directly to chondrichthyans (Nakken, 2003). The introduction of sorting grids (Nordmøre grid) in the shrimp fishery in 1992 reduced the bycatch significantly, and only juvenile chondrichthyans (generally <25 cm) have been caught since (Reithe and Aschan, 2004).

Climate may also play a part in determining the biogeographical distribution of the group. Therefore, studies on the distribution and movement of elasmobranchs should examine environmental

Table 1. Average bycatch (kg) and standard deviation (s.d.) by gear type, and the number of reported landings (*n*) of chondrichthyan species along the northern coast of Norway during the period 2000–2005, based on official landings data provided by the Norwegian Directorate of Fisheries.

Taxon	Gillnet		Longline		Other		Total		<i>n</i>
	Average	s.d.	Average	s.d.	Average	s.d.	Average	s.d.	
<i>Chimaera monstrosa</i>	7	6	3 501	7 305	0	–	3 239	7 080	40
<i>Lamna nasus</i>	190	454	79	99	68	64	177	431	339
<i>Squalus acanthias</i>	6 020	15 626	1 383	3 751	1 845	452	2 735	9 998	1 188
Dogfish and hounds	20	12	952	2 001	13	9	485	1 444	14
<i>Dipturus batis</i>	212	337	73	97	19	22	143	273	121
Rajidae	656	946	468	1 181	272	461	465	946	1 441
Total	2 357	9 548	926	2 937	225	449	1 315	6 338	3 143

Some of the landings are reported in aggregated categories: “Dogfish and hounds” and “Rajidae”. Such records may contain landings of the main species and other species not identified when caught or delivered.

parameters associated with the distribution of the various species (Pawson and Ellis, 2005). Dolgov *et al.* (2005a) suggested that the distribution of various skate species in the Barents Sea appeared to be related to sea temperature. Since the 1990s, there has been a marked increase in sea temperature, particularly in the southern part of the coast of northern Norway (Pawson and Ellis, 2005). Many chondrichthyans found along this coast are close to their geographical limits, so changes in environmental conditions may be expected to affect their local abundance.

The aim of this study is (i) to identify species present along the northern coast of Norway between 1992 and 2005, (ii) to describe their distribution and abundance, (iii) to identify significant spatial or temporal differences between species, and (iv) to evaluate whether changes observed in distribution and abundance over time are affected by sea temperature. Such information will, we believe, help to resolve the issue of identifying appropriate stock units for management.

Material and methods

The study area consists of fjords and offshore banks along the northern coast of Norway from south of Ålesund (62°00'N 4°50'E) to the Russian border near Kirkenes (69°50'N 30°50'E; Figure 1). Fjords north of 68°N are mainly shallower than 300 m, though those to the south are generally deeper. The coastal banks outside the fjords range in depth from 50 to

400 m (Berg and Albert, 2003). Despite the high latitude, sea temperatures are generally warmer than in other northern coastal areas because of the influence of the Norwegian Current, a branch of the Gulf Stream that flows northeast along the coast. As the current passes through higher latitudes, there is an overall reduction in sea temperature. Temperatures are not constant and fluctuate in short- and long-term intervals (Gyory *et al.*, 2005), and average sea temperatures have increased over the past century (Berstad *et al.*, 2003).

Temperature data

During the period 1935–1947, several permanent hydrographical sampling stations were established along the Norwegian coast by the Institute of Marine Research (IMR, 2007). Temperature data from four stations, Bud (62°56'N 6°47'E), Eggum (68°22'N 13°38'E), Ingøy (71°08'N 24°01'E), and Vardø (70°45'N 31°03'E), were used in this study (Figure 1). Water temperature and salinity were measured using CTD sensors deployed from research vessels. Approximate bottom temperatures were measured as close to the seabed as the equipment would allow (generally within 10 m) and registered to an accuracy of 0.01°C. Annual temperatures at each station were calculated as the mean of quarter-year values.

Survey data

The survey data were from the annual combined trawl and acoustic surveys conducted each autumn by the Norwegian Institute of Fisheries and Aquaculture Research (Fiskeriforskning) from 1992 to 2001 and by IMR during the period 2002–2005 (Table 2). The main aim of these coastal surveys was to investigate commercial species such as coastal cod, haddock, saithe (*Pollachius virens*), and juvenile herring (*Clupea harengus*).

At the start of the survey in 1992, the intended survey area was divided into three sections to determine the feasibility of the survey and to facilitate the development of a practical design. One section was intensely sampled each year, the northern section in 1992, the central section in 1993, and the southern section in 1994 (Table 2). Since 1995, the entire area has been sampled annually. The surveys then began at the northeastern limit of the area, covered the Norwegian coast to 62°N, and lasted ~30 d (Figure 1). Sampling was evenly distributed along the coast, and included fjords and offshore areas near the coast (Figure 1). Sampling stations were not selected randomly,





Figure 1. Survey area and strata along the northern coast of Norway. Positions of permanent hydrographic stations are marked with circles.

Table 2. Survey period, vessel, statistical areas (Directorate of Fisheries), trawl gear specifications, and number of stations for each coastal survey conducted along the Norwegian coast from 62 to 70°N during the period 1992–2005.

Survey period	Vessel	Statistical areas	Inner net mesh size (mm)	Strapping	Number of stations
25 August to 3 October 1992	RV "Johan Ruud"	03, 04, and northern sector of 05	20	Yes	127
1 August to 8 September 1993	RV "Mikael Sars"	05 and eastern limit of 37	20	Yes	163
22 September to 19 October 1994	RV "Mikael Sars"	06, 07, and eastern limit of 37	35	No	106
13 September to 11 October 1995	RV "Mikael Sars"	00, 03, 04, 05, 06, and 07	35	No	129
11 September to 6 October 1996	RV "Mikael Sars"	As above	35	No	130
20 August to 23 September 1997	RV "Mikael Sars"	As above	20	For most	159
26 October to 19 November 1998	RV "Jan Mayen"	As above	20	Yes	132
22 October to 19 November 1999	RV "Jan Mayen"	As above	20	Yes	154
23 October to 16 November 2000	RV "Jan Mayen"	As above	20	Yes	154
23 October to 17 November 2001	RV "Jan Mayen"	As above	20	Yes	125
29 October to 26 November 2002	RV "Jan Mayen"	As above	20	Yes	168
11 October to 14 November 2003	RV "Jan Mayen"	As above	20	For most	133
	RV "Johan Hjort"				
13 October to 9 November 2004	RV "Jan Mayen"	As above	20	For most	128
	RV "Johan Hjort"				
11 October to 8 November 2005	RV "Jan Mayen"	As above	20	For most	124
	RV "Johan Hjort"				

because the seabed in fjords and over the shelf is often too rough to permit trawling (Berg and Albert, 2003). The same stations were approximately fixed for each survey from 1995 to 2005, although poor weather conditions or technical difficulties resulted in some stations occasionally being omitted. Catches were considered reasonably representative of substrata suitable for trawling at 30–700 m. The deepest average depth trawled was around 65°N (Figure 2).

The sampling trawl was a Campelen 1800 shrimp trawl with a 30-m headline, 19-m groundrope, 80–42-mm knot-to-knot stretched mesh in the body, and 20-mm standard mesh size in the inner net (Table 2; Aschan and Sunnanå, 1997). The gear had 40-m bridles and rock-hopper groundgear, with eight steel spacers between 14 rubber discs. Sensors monitored trawl geometry, and strapping constrained the distance between doors to ~47 m while trawling (Aschan and Sunnanå, 1997). At this door spread, the silt plume was directed towards the trawl wings and considered to maximize the herding effect between doors and net.

**Figure 2.** Average bottom depth with standard deviation of trawl stations taken in each area along the northern coast of Norway during the coastal surveys of 1992–2005.

Species identification

All species were identified, counted, and weighed. In 14 of the trawl samples taken between 1997 and 2004, all the skate species (Rajidae) were grouped and recorded generically, so these observations were excluded from this study. Originally, the names were in Norwegian, according to the species list in the IMR quality control system (Mjanger *et al.*, 2004), but the species list of 2004 included Latin names. These names have been monitored and, where appropriate, updated to valid scientific names according to the Integrated Taxonomic Information System (ITIS, 2008).

The reliability of the results from this study depends on the species identifications being accurate and consistent. Unfortunately, no voucher specimens were collected because the main target of the survey was the commercially important teleosts. Therefore, uncertainties in species identification were analysed during the 2006 survey. Participating scientists were observed sorting and identifying fish, then interviewed to understand better the limitations of the identification procedures, especially for skates, which are a problematic taxonomic group (e.g. *Raja montagui* is sometimes confused with *Raja brachyura*, according to J. R. Ellis, pers. comm., and *Raja clavata* and *Amblyraja radiata* are often confused—Daan, 2001). Because of the uncertainties involved in modifying the raw data, changes in identification (described by Williams, 2007) were limited to clear inconsistencies that were demonstrated between personnel shifts during some surveys. Individual *A. radiata* may have been misidentified as *R. clavata* during the 1990s, but *R. clavata* has rarely been taken, then mainly south of 68°N (W. Richardsen, pers. comm.). However, we cannot confirm these recordings, so have excluded *R. clavata* from our analyses.

Abundance and distribution mapping

The estimated spatial distribution for a species was based on data from the entire survey period (1992–2005). Abundance was expressed as number of individuals km^{-2} at each station. Abundance was estimated based on the area trawled at each

station using the method of Jakobsen *et al.* (1997):

$$\rho_s = \frac{f_s}{a_s}, \quad (1)$$

where ρ_s denotes the abundance in number of fish km^{-2} at sample station s , f_s the number captured (frequency) at sample station s , and a_s the area swept (km^2) at sample station s (nautical miles trawled $\times 1.852 \times 0.047 \text{ km}$).

The door spread of 47 m was assumed to be the upper limit of the mean effective catching width of the gear. A more precise estimate of catch efficiency was beyond the scope of this study.

The annual mean catch (number km^{-2}) was used to compare species abundance. Species with an annual mean catch >5 animals km^{-2} were grouped as common species, and included for further statistical analysis. Species with lower catch rates were either grouped as infrequent, if previously recorded in the survey area, or rare if not recorded previously.

Pethon (2005) and FishBase (Froese and Pauly, 2007) give previously estimated distributions. The two sources were generally consistent, although they differed for *Dipturus batis*, *Leucoraja fullonica*, *Dipturus lineatus*, *Bathyraja spinicauda*, and *Galeus melastomus*. All five of these species are described by Pethon (2005) as having a more northerly distribution than given in FishBase (Froese and Pauly, 2007). In our opinion, the species distributions of Pethon (2005), which are updated based on recent data, were more accurate than those in FishBase, so were taken as the standard distributions for the present study.

Statistical analysis

For seven common species, the relative abundance (number km^{-2}) in each trawl sample was used to assess the statistical significance of temporal and spatial variation in abundance. Year, depth, and area were the independent variables. The survey area was divided into seven subareas by degree latitude from 62 to 69°N, and a northern region (from 69 to 71°N) was divided into two subareas east and west of 25°E (Figure 1). Samples were identified by 50-m depth interval, encompassing the depth range covered by the survey. Average abundance for each species was calculated, and differences in abundance were tested against the three variables, depth, latitude, and year, using one-way single factor ANOVA. The significance level was set at $p = 0.05$.

A constrained (canonical) correspondence analysis (CCA) was run in R 2.5.0 (R Development Core Team, 2007) using the vegan package (Oksanen, 2007). As the species data contained many zeros, they were analysed using unimodal methods (ter Braak and Verdonschot, 1995). The ten most common species were included in the analysis, and their abundance estimates were $\log(x + 1)$ transformed. Potential explanatory variables were longitude, latitude, depth, and year of sampling. Shifts in abundance of each species were shown as percentages of deviation from the average abundance within the total survey area and presented with respect to the strongest explanatory variable.

Correlations were tested for all species with distributions significantly correlated with latitude. The locations of the four temperature stations determined the areas 63°N, 68°N, West (69–71°N), and East (69–71°N; Figure 1) used for assessing correlations between abundance and temperature. Spearman's rank-order correlation (ρ) was used to test whether latitudinal or annual differences in temperature significantly influenced the distribution of nine common species.

Results

Abundance and distribution

During the survey period 1992–2005, 18 species of Chondrichthyes were recorded at 1932 stations (Table 3). The most diverse orders observed were skates (Rajiformes, 13 species) and dogfish sharks (Squaliformes, 3 species). In contrast, only one species of catshark (Scyliorhinidae) and rabbitfish (Holocephali) was identified in the samples. Six species were observed across the entire latitudinal range (62–71°N), and many species seemed to have a boundary in the north (Figures 3 and 4). Mean annual catch rates and frequency of occurrence for each species reveal that *Chimaera monstrosa*,

Table 3. Chondrichthyan species identified along the northern Norwegian coast during the coastal surveys of 1992–2005.

Scientific name	Family	Common name	Norwegian Red List*
<i>Chimaera monstrosa</i> Linnaeus, 1758	Chimaeridae	Rabbitfish	No
<i>Etmopterus spinax</i> (Linnaeus, 1758)	Dalatiidae	Velvet belly	No
<i>Somniosus microcephalus</i> (Bloch and Schneider, 1801)	Dalatiidae	Greenland shark	Near-threatened
<i>Galeus melastomus</i> Rafinesque, 1810	Scyliorhinidae	Blackmouth catshark	No
<i>Squalus acanthias</i> Linnaeus, 1758	Squalidae	Spurdog	Critically endangered
<i>Amblyraja hyperborea</i> (Collett, 1879)	Rajidae	Arctic skate	Data-deficient
<i>Amblyraja radiata</i> (Donovan, 1808)	Rajidae	Thorny skate	No
<i>Bathyraja spinicauda</i> (Jensen, 1914)	Rajidae	Spinetail ray	Data-deficient
<i>Dipturus batis</i> (Linnaeus, 1758)	Rajidae	Blue skate	Data-deficient
<i>Dipturus lineatus</i> (Fries, 1838)	Rajidae	Sailray	Data-deficient
<i>Dipturus nidarosiensis</i> (Storm, 1881)	Rajidae	Norwegian skate	Data-deficient
<i>Dipturus oxyrinchus</i> (Linnaeus, 1758)	Rajidae	Longnosed skate	Data-deficient
<i>Leucoraja circularis</i> (Couch, 1838)	Rajidae	Sandy ray	No
<i>Leucoraja fullonica</i> (Linnaeus, 1758)	Rajidae	Shagreen ray	Data-deficient
<i>Raja brachyura</i> Lafont, 1873	Rajidae	Blonde ray	No
<i>Raja clavata</i> Linnaeus, 1758	Rajidae	Thornback ray	No
<i>Raja montagui</i> Fowler, 1910	Rajidae	Spotted ray	Data-deficient
<i>Rajella fyllae</i> (Lütken, 1887)	Rajidae	Round skate	No

*Species entry in the 2006 Norwegian Red List—Artsdatabanken Norwegian Biodiversity Information Centre.




Figure 3. Distribution and abundance of (a) *Chimaera monstrosa*, (b) *Etmopterus spinax*, (c) *Galeus melastomus*, (d) *Amblyraja radiata*, (e) *Squalus acanthias*, (f) *Dipturus oxyrinchus*, (g) *Rajella fyllae*, and (h) *Dipturus batis* along the northern coast of Norway from the coastal surveys of 1992–2005. Note that the abundance scales differ between panels. The shaded area is the distribution according to Pethon (2005).

Etmopterus spinax, *G. melastomus*, *A. radiata*, *Squalus acanthias*, and *Dipturus oxyrinchus* were the main species (Table 4). For many species, abundance sometimes varied greatly between years, but there were no obvious negative or positive trends, indicating no recent changes in abundance of the more common species (Figure 5). One exception was *Rajella fyllae*, which exhibited an increasing trend in the data. Annual changes in abundance were

only significant ($p < 0.05$) for *A. radiata* (Table 5), which increased significantly between 2002 and 2003, decreased by the same extent between 2004 and 2005, but had no apparent long-term trend (Figure 5). Latitude was a significant factor (ANOVA, $p < 0.05$) affecting the abundance of all common species (Table 6).

Chimaera monstrosa and *E. spinax* (Figures 3a and b) were observed in all areas except in the far northeast (East 69–70°N).




Figure 4. Distribution and abundance of (a) *Leucoraja fullonica*, (b) *Leucoraja circularis* and *Bathyraja spinicauda*, (c) *Amblyraja hyperborea*, (d) *Dipturus nidarosiensis*, (e) *Raja montagui* and *Raja brachyura*, and (f) *Dipturus linteus* and *Somniosus microcephalus* along the northern coast of Norway from the coastal surveys of 1992–2005. Note that the abundance scales differ between panels. The shaded area is the distribution according to Pethon (2005).

Abundance was greatest in the south, where catch rates were occasionally >2000 fish per haul, suggesting an aggregating behaviour by the two species. Distributions of *G. melastomus*, *S. acanthias*, and *D. oxyrinchus* (Figures 3c, e, and f) appeared to be constrained to areas south of 68°N, with most catches south of 65°N. Catches of *G. melastomus* of >200 animals in each of 10 hauls highlights the aggregating nature of this species. *Amblyraja radiata* was the dominant skate species and was caught throughout the survey area, but in greater abundance in the north. The distribution of *R. fyllae* was similar (Figures 3d and g).

Abundance appeared to be greatest for *C. monstrosa* at 450–550 m, for *G. melastomus* deeper than 150 m, for *A. radiata* at 600–650 m, and for *R. fyllae* and *D. batis* shallower than 500 m (Table 4; Figure 6). However, differences in depth-dependent abundance were only statistically significant for *G. melastomus* (Table 7), with greatest abundance in the 500-m interval. The three other common species were more evenly distributed with respect to depth (Figure 6).

The constrained correspondence biplot shows the species scores that may be taken as the optimal location for nine species in the environment field spanned by the site scores (Figure 7). The constrained axis CCA1 (eigenvalue 0.457, 88%) has a much larger explanatory value than CCA2 (eigenvalue 0.049, 9%). Latitude seemed to structure the chondrichthyan assemblage into three groups (Figure 7), a northern component consisting of *A. radiata* and *R. fyllae*, a central component with *C. monstrosa*, *D. batis*, *E. spinax*, and *L. fullonica*, and a southern component consisting of *D. oxyrinchus*, *G. melastomus*, and *S. acanthias*. Relative abundance along the latitude shows the same pattern for species of each group (Figure 8). Depth and year were of little importance in defining species distribution (Figure 7).

Of the four species ranked as infrequent, *D. batis* was observed along the entire coast at depths of 85–425 m, and *L. fullonica* from 77–512 m in all areas except the eastern sector of 69–70°N (Table 4; Figure 4). *Dipturus nidarosiensis* was found as far north as 68°N at depths of 140–590 m, whereas its known

Table 4. Average annual catch rates and standard deviations, proportion of trawl stations where the species was observed in the coastal surveys of 1992–2005, qualitative abundance category, and latitude and depth ranges.

Species	Mean catch (number km ⁻²)	s.d.	Proportion of stations at which observed (%)	Abundance	Latitude (°N)	Depth (m)
<i>Chimaera monstrosa</i>	2 586.2	3 250.7	29.5	Common	62–71	42–665
<i>Etmopterus spinax</i>	2 078.4	2 234.0	17.7	Common	62–71	47–730
<i>Galeus melastomus</i>	605.7	598.0	13.2	Common	62–68	51–665
<i>Amblyraja radiata</i>	55.2	20.7	19.6	Common	62–72	30–635
<i>Squalus acanthias</i>	49.3	49.3	4.5	Common	62–69	42–515
<i>Dipturus oxyrinchus</i>	7.7	15.6	2.0	Common	62–68	45–665
<i>Rajella fyllae</i>	5.5	8.3	2.2	Common	62–72	83–415
<i>Dipturus batis</i>	2.2	2.2	1.0	Infrequent	62–71	85–425
<i>Leucoraja fullonica</i>	1.8	2.7	1.0	Infrequent	62–71	77–512
<i>Leucoraja circularis</i>	1.7	3.2	0.4	Rare	62	88–244
<i>Bathyraja spinicauda</i>	1.1	1.4	0.5	Rare	67–72	48–410
<i>Amblyraja hyperborea</i>	0.6	1.9	0.3	Rare	67–71	125–620
<i>Dipturus nidarosiensis</i>	0.5	0.9	0.3	Infrequent	62–69	132–588
<i>Raja montagui</i>	0.4	0.6	0.3	Rare	62	63–120
<i>Raja brachyura</i>	0.3	1.1	0.02	Rare	62–63	99
<i>Dipturus linteus</i>	0.1	0.3	0.1	Rare	68	588
<i>Somniosus microcephalus</i>	0.1	0.3	0.02	Infrequent	69	480

distribution was primarily south of 65°N (Pethon, 2005). A single *S. microcephalus* was caught 480 m deep at 69°10'N 16°19'E in 1993.

From 1992 to 2005, six rare species were reported, extending the distributions given by Pethon (2005). *Raja brachyura*, *R. montagui*, and *Leucoraja circularis* were all caught south of 64°N, but at higher latitudes than previously documented (Pethon, 2005). Of these three, *L. circularis* was caught most frequently and in greatest number, 23 animals in six trawls. The depths of capture for *L. circularis* were 88–244 m, and for *R. montagui* 63–120 m, and four *R. brachyura* were caught at 99 m. Totals of 11 *B. spinicauda* and nine *Amblyraja hyperborea* were caught in trawls north of 67°N at depth ranges of 48–410 m and 125–620 m, respectively, and one *D. linteus* was caught at 588 m off Lofoten at 68°N.

Abundance and temperature

Except the eastern sector at 69–71°N, all areas appeared to show an overall increase in sea temperature between 1992 and 2005 (Figure 9). Except *R. fyllae*, the abundance of chondrichthyans showed no obvious trend during the study period (Figure 5). Therefore, the Spearman rank analysis showed no significant correlations between annual sea temperature and abundance for each species in the areas 63°N, 68°N, 69–71°N (West), and 69–71°N (East). Bottom temperature decreased in a northward direction along the coast. Latitudinal shifts in abundance for *C. monstrosa*, *E. spinax*, *G. melastomus*, and *S. acanthias* showed positive correlations ($\rho > 0.9$) with bottom temperature, whereas *A. radiata* abundance showed a negative correlation ($\rho > -0.9$) with temperature.

Discussion

Survey data uncertainty

The surveys were designed primarily to assess the commercially important teleosts, so chondrichthyans were not taken into account in the design. Consequently, it is uncertain how accurately the catch rates reflect the relative abundance of the various species

(Kotwicki and Weinberg, 2005). As trawl catchability varies with bottom type and species, there is little information available for estimating absolute stock size (Bonfil, 1994; Abella and Serena, 2005; Daan *et al.*, 2005; Dolgov *et al.*, 2005a). Moreover, species that favour hard rocky substrata (e.g. *D. batis*) are likely to be underrepresented because most trawling was conducted on seabed that could be trawled, i.e. was fairly smooth.

Species identification issues are important, although this was corrected where possible (Williams, 2007). Misidentification of skates is common (Daan, 2001), and except the more visually distinct species such as *D. nidarosiensis*, there is still concern regarding the validity of the skate identifications. The main uncertainty in our results is the soundness of the estimated distribution of the infrequent and rare species such as *R. clavata* (which was excluded from the analyses). For the common species, the data are considered to be valid for describing their relative abundance and distribution.

An update on chondrichthyan distributions

The porbeagle (*Lamna nasus*), which is recorded as a bycatch in the area (Table 1), was not caught at all during the coastal surveys, because the species is rarely taken by trawl (Daan *et al.*, 2005). Basking sharks (*C. maximus*) are caught as a bycatch in gillnet fisheries, but were not in our survey data.

From 1992 to 2005, the distribution and abundance of the common species appeared to remain stable (Figure 5). Latitudinal trends in distribution correlated well with previous distributions given by Pethon (2005). Shark species and *C. monstrosa* were clearly more abundant south of 65°N. *Chimaera monstrosa* and *E. spinax* appeared to be the most abundant species, including north of 70°N (Figure 3). *Chimaera monstrosa* has recently also been observed in the southern Barents Sea (Dolgov, 2006; Byrkjedal and Høines, 2007). *Amblyraja radiata* is uniformly and widely distributed, and was the dominant skate, followed by *R. fyllae*. The dominance of these two species agrees with studies undertaken in the neighbouring




Figure 5. Mean abundance of all chondrichthyan species along the northern coast of Norway from the coastal surveys of 1992–2005. Note that the abundance scales differ between panels.

Barents Sea, northeastern North Sea, and Norwegian Sea (Skjæraasen and Bergstad, 2001; Dolgov *et al.*, 2005a; Dolgov, 2006; Byrkjedal and Høines, 2007). In the south, *D. oxyrinchus* appeared to be more abundant than *R. fyllae*. This may be a fairly localized population, because *D. linteus* replaces *D. oxyrinchus* in the skate assemblage that dominates the neighbouring northeastern North Sea and Norwegian Sea (Skjæraasen and Bergstad, 2001). *Raja clavata* has been recorded as far north as the Barents Sea (Hognestad and Vader, 1979; Fossheim *et al.*, 2006; Byrkjedal and Høines, 2007), but was not observed during Russian surveys from 1996 to 2007 (A. V. Dolgov, pers. comm.). We believe that this species may be a sporadic visitor to the whole northern coast of Norway and may also be taken occasionally in the southwestern Barents Sea. However, the real distribution of *R. clavata* needs further clarification because

identification of this species in our data seems to have been biased; voucher specimens are required to confirm its occurrence in northern Norwegian waters.

Because of a lack of knowledge and infrequency of recordings of the rare and infrequent species, it is impossible to be certain of any distribution shifts. Our observations show that *R. brachyura*, *L. circularis*, and *R. montagui*, commonly associated with the North Sea and Atlantic areas south of 62°N (Dulvy *et al.*, 2000; Pethon, 2005; Froese and Pauly, 2007), may all be present as far north as 64°N. This is probably not attributable to a change in distribution, but rather because of poor data historically. *Amblyraja hyperborea* and *B. spinicauda* are associated with offshore areas (Mahon *et al.*, 1998; Pethon, 2005; Fossheim *et al.*, 2006; Byrkjedal and Høines, 2007), but were found closer to the coast in our study.

Table 5. One-way single factor ANOVA schemes for determining the significance of temporal differences in abundance of the main species caught during the coastal surveys of 1992–2005.

Species	Sum of squares	d.f.	Mean square	F	p	F _{crit}
<i>Chimaera monstrosa</i>	3 717 741.6	11	337 976.5	0.98	0.473	1.88
<i>Etmopterus spinax</i>	893 976.4	11	81 270.6	0.58	0.840	1.88
<i>Galeus melastomus</i>	231 510.3	11	21 046.4	0.68	0.751	1.88
<i>Squalus acanthias</i>	5 282.6	11	480.2	0.76	0.675	1.88
<i>Amblyraja radiata</i>	184.2	11	16.7	1.92	0.044	1.88
<i>Dipturus oxyrinchus</i>	52.4	11	4.8	1.00	0.449	1.88
<i>Rajella fyllae</i>	14.0	11	1.3	1.61	0.107	1.88

Significant differences ($p = 0.05$) emboldened.

Table 6. One-way single factor ANOVA schemes for determining the significance of latitudinal differences in abundance of the main species caught during the coastal surveys of 1992–2005.

Species	Sum of squares	d.f.	Mean square	F	p	F _{crit}
<i>Chimaera monstrosa</i>	7 537 614.7	8	942 201.8	2.8	0.007	2.03
<i>Etmopterus spinax</i>	5 141 623.0	8	642 702.9	6.0	<0.001	2.03
<i>Galeus melastomus</i>	1 078 162.9	8	134 770.4	5.5	<0.001	2.03
<i>Squalus acanthias</i>	13 594.7	8	1 699.3	2.8	0.007	2.03
<i>Amblyraja radiata</i>	294.3	8	36.8	5.3	<0.001	2.03
<i>Dipturus oxyrinchus</i>	162.5	8	20.3	5.0	<0.001	2.03
<i>Rajella fyllae</i>	10.3	8	1.3	2.8	0.008	2.03

Significant differences ($p = 0.05$) emboldened.

Spatial distribution

Amblyraja radiata has a wide and uniform distribution throughout the study area, with biomass increasing to the north, and is found also in the Barents Sea (Dolgov *et al.*, 2005a; Byrkjedal and Høines, 2007). Annual distribution maps show that the distributions of *C. monstrosa*, *E. spinax*, *G. melastomus*, and *S. acanthias* are patchy (Williams, 2007). The large catches (>500 animals in a single haul) underscore the aggregating behaviour of these species. *Squalus acanthias* occurs often in shoals of the same sex and/or size (Ellis *et al.*, 2005b; Stenberg, 2005), similar to *G. melastomus*, for which there are also bathymetric patterns (Massuti and Moranta, 2003; Calis *et al.*, 2005; Coelho *et al.*, 2005). These uneven distributions can be linked to the availability of suitable bottom substrata or food availability, as has been suggested to explain similar distribution patterns in the neighbouring North Sea and Skagerrak (Skjæraasen and Bergstad, 2000). Tagging studies indicate little mixing of *S. acanthias* between northern and southern areas of the North Sea (Aasen,

1962; Holden, 1967), and Holden (1968) claimed that the Norwegian–Scottish and Channel populations are separate stocks. Despite assertions of transatlantic migration (Holden, 1967; Templeman, 1984), recent analysis of tag returns indicates that Northeast and Northwest Atlantic stocks should be managed independently. No decline over time was observed for *S. acanthias* in our study area, but declines have been documented for the North Sea, Celtic Sea, and off Northwest Scotland (Daan *et al.*, 2005; Dobby *et al.*, 2005; Ellis *et al.*, 2005a; ICES, 2007). Those studies, however, were based on surveys that started in the 1970s and 1980s. The few large catches (7–19 fish) indicate that, although *D. oxyrinchus* is relatively scarce, local aggregations may occur.

The abundance of some species appeared to be depth-dependent. *Dipturus batis* and *R. fyllae* were confined to water shallower than 500 m (Table 4). However, *R. fyllae* probably exists over a wider depth range (Dolgov *et al.*, 2005a; Jørgensen *et al.*, 2005; Pethon, 2005) and probably prefers deeper waters in the south (Skjæraasen and Bergstad, 2001). As observed in other areas, *G. melastomus* preferred depths deeper than 150 m (Magnussen, 2002; Massuti and Moranta, 2003; Rey *et al.*, 2004; Costa *et al.*, 2005; Serena *et al.*, 2006). *Chimaera monstrosa* was found mainly in deeper water (>400 m), but may migrate to shallower water in spring and summer to deposit egg capsules (Bristow, 1992; Pethon, 2005). *Etmopterus spinax* was found in both deep and shallow water in the north and the south of the survey area, and the depth range was not clearly dependent on latitude in the survey area, as suggested by Pethon (2005). *Amblyraja radiata* and *D. oxyrinchus* were present at all depths.

Species assemblages

The northern, central, and southern species assemblages shown by the constrained correspondence biplot revealed that factors linked to latitude play an important role in determining species distribution and abundance. The sharp decline in abundance north of 65°N for species in the southern and central group was particularly noteworthy because it suggests a latitudinal change in conditions, resulting in a shift in biodiversity. The 65°N region is, on average, deeper than the other areas surveyed, and the deep trench may function as a biogeographical barrier (Figure 2). However, depth alone does not appear to explain this trend, and the Lofoten Peninsula may function as a physical barrier.

Temporal and spatial shifts with regard to sea temperature

Because of the small number of stations with temperature measurements, sea temperature had limited ability to explain shifts in abundance in our data. Skate distribution in the Barents Sea has been linked to changes in bottom temperature (Dolgov *et al.*, 2005a). In some areas of the Northeast Atlantic, the increase in sea temperature in recent years has coincided with a gradual northward shift in the distribution of some species (Perry *et al.*, 2005; Dolgov, 2006). So far, though, there appear to have been no such shifts along the northern coast of Norway. Demersal fish species such as cod seem to be able to adapt to moderate changes in their ambient temperature and do not necessarily respond to it with a change in distribution (Ottersen *et al.*, 1998), at least to the same extent as do pelagic fish, such as blue whiting (*Micromesistius poutassou*), capelin (*Mallotus villosus*), and herring (Bergstad *et al.*, 1999; Toresen and Østvedt, 2000; Fosshem *et al.*, 2006). Depth and temperature are unlikely to be the only factors involved in determining the




Figure 6. Mean abundance and standard error for the common species in each depth zone along the northern coast of Norway from the coastal surveys of 1992–2005. Note that the abundance scales differ between panels.

apparent trends in distribution and general stability in abundance of each chondrichthyan listed here. As the distributions of the southern and central species assemblages have not expanded north in response to ocean warming, other factors may be playing a role in determining abundance and distribution.

Little is known about how fisheries have impacted chondrichthyan populations along the northern coast of Norway. However, we know that they provide a significant and probably underestimated bycatch in gillnet and longline fisheries (Table 1). It is therefore reasonable to assume that the demersal fisheries in particular have a negative impact on chondrichthyan stocks (Bonfil, 1994; Stevens *et al.*, 2000; Dolgov *et al.*, 2005b; Drevetnyak *et al.*, 2005). Any major changes in population structure in response to fishing probably took place in the survey area before 1992, when the most significant development in the fisheries there took place. As mentioned, the introduction of sorting grids in the shrimp fishery reduced bycatch to include only juveniles. This has not

resulted in an obvious increase in chondrichthyan abundance, but may be one reason the stocks are not decreasing. The populations of *A. radiata* and *R. fyllae* in the neighbouring Barents Sea appear to be stable at current fishing levels (Drevetnyak *et al.*, 2005), so both species are probably similarly tolerant to current fishing pressure within our study area.

The processes influencing the population dynamics of the chondrichthyans described here are undoubtedly complex. Geographical barriers, particularly the deep trenches in the region of 67°N (Vestfjord) and the Lofoten Peninsula, could restrict passage and inhibit an expansion in the distribution of a species. Also, chondrichthyans are generally long-lived, and their distribution may to some degree be the consequence of territorial behaviour and a slow rate of migration. For example, some skates do not migrate great distances (Hunter *et al.*, 2005) and may show clear gaps between areas of high concentration, perhaps indicating that they may form local stock units (Daan *et al.*, 2005). However,

Table 7. One-way single factor ANOVA schemes for determining the significance of depth-dependent differences in abundance of the main species caught during the coastal surveys of 1992–2005.

Species	Sum of squares	d.f.	Mean square	F	p	F _{crit}
<i>Chimaera monstrosa</i>	40 047 077	9	4 449 675.2	1.46	0.170	1.95
<i>Etmopterus spinax</i>	1 386 782	9	154 086.9	1.69	0.098	1.95
<i>Galeus melastomus</i>	352 063	9	39 118.1	3.23	0.001	1.95
<i>Squalus acanthias</i>	4 807	9	534.1	1.65	0.107	1.95
<i>Amblyraja radiata</i>	155	9	17.2	1.21	0.297	1.95
<i>Dipturus oxyrinchus</i>	135	9	15.0	0.97	0.469	1.95
<i>Rajella fyllae</i>	5	9	0.6	1.81	0.072	1.95

Significant differences ($p = 0.05$) emboldened.




Figure 7. Ordination biplot of constrained correspondence analysis (CCA) relating the abundance of the ten most abundant chondrichthyan species along the northern Norwegian coast to the potential explanatory variables, longitude, latitude, depth, and year of sampling. Am_rad, *Amblyraja radiata*; Ch_mon, *Chimaera monstrosa*; Di_bat, *Dipturus batis*; Di_ox, *Dipturus oxyrinchus*; Et_spi, *Etmopterus spinax*; Ga_mel, *Galeus melastomus*; Le_ful, *Leucoraja fullonica*; Ra_fyl, *Rajella fyllae*; and Sq_aca, *Squalus acanthias*.

A. radiata seems to be an exception, because the continental shelf edge apparently does not present a barrier to its migration, and there are no significant population gaps in the North Atlantic generally (Chevolot *et al.*, 2007). This is one explanation for the uniform distribution of *A. radiata* in the study area and in the Barents Sea.

Although chondrichthyans are potentially vulnerable to fisheries (Stevens *et al.*, 2000; Priede *et al.*, 2006), the current populations in the study area of the more common species appear not to be adversely affected by bycatch at the current levels of fishing activity (although historical estimates of abundance are




Figure 8. Shifts in abundance, given as percentages of deviation from mean abundance (number km⁻²) within the total survey area, with respect to latitude area for species grouped as (a) northern, (b) central, and (c) southern assemblages.




Figure 9. Mean annual temperature at the hydrological stations along the northern coast of Norway at Vardø, Ingøy, Bud, and Eggum.

not available). Stocks of *C. monstrosa*, *E. spinax*, *S. microcephalus*, *G. melastomus*, *S. acanthias*, *D. batis*, and recently also *A. radiata* have declined in the North Sea (Daan *et al.*, 2005; Ellis *et al.*, 2005a; Jones *et al.*, 2005; ICES, 2007), but there has been no such decline in our area of interest. This may be due to the lower effort in the demersal fishery in general and in the elasmobranch fishery in particular along the northern coast of Norway over the period 1992–2005. However, because of the limitations of our data, we cannot be certain that this is the case for the less frequently caught species, especially given the comparatively short period covered by the study. For example, species such as

S. microcephalus are reported to have had longer term declines in Norwegian waters (Ruud, 1968).

As species identification is currently difficult and potentially inaccurate, effort should be put into quality controlling available taxonomic keys for northern areas, especially for skates, which are known to have high morphological interspecific variability (Serena *et al.*, 2005). To address the distribution of skate species in Norwegian waters, future surveys should place more emphasis on species identification, including the collection of voucher specimens, and the development of appropriate field identification guides should be given high priority. Knowledge of chondrichthyan species along the coast of Norway remains limited compared with that for the neighbouring North Sea (ICES, 2007) and further work is required.

Acknowledgements

An anonymous referee and Jim Ellis provided valuable advice and recommendations for improving the manuscript, and Frøydis Strand kindly helped us finalize the figures. We thank Erik Berg and Ivan Ahlquist (IMR) for making the fish abundance data available, and Øyvind Strand (IMR) and all others who were involved in helping to find suitable temperature data. Andrey Dolgov supported us through interesting discussions. Finally, we thank the crew of RV “Jan Mayen” for their support.

References

- Aasen, O. 1962. Norwegian dogfish tagging. *Annals of Applied Biology*, 17: 106–107.
- Abella, A. J., and Serena, F. 2005. Comparison of elasmobranch catches from research trawl surveys and commercial landings at port of Viareggio, Italy, in the last decade. *Journal of Northwest Atlantic Fishery Science*, 35: 345–356.
- Aschan, M., and Sunnanå, K. 1997. Evaluation of the Norwegian shrimp surveys conducted in the Barents Sea and Svalbard area 1980–1997. ICES Document CM 1997/Y: 07. 24 pp.
- Berg, E., and Albert, O. T. 2003. Cod in fjords and coastal waters of North Norway: distribution and variation in length and maturity at age. *ICES Journal of Marine Science*, 60: 787–797.
- Bergstad, O. A., Bjelland, O., and Gordon, J. D. M. 1999. Fish communities on the slope of the eastern Norwegian Sea. *Sarsia*, 84: 67–78.
- Berstad, I. M., Sejrup, H. P., Klitgaard-Kristensen, D., and Haflidason, H. 2003. Variability in temperature and geometry of the Norwegian Current over the past 600 years; stable isotope and grain size evidence from the Norwegian margin. *Journal of Quaternary Science*, 18: 591–602.
- Bonfil, R. 1994. Overview of world elasmobranch fisheries. FAO Fisheries Technical Paper, 341. 119 pp.
- Bristow, G. A. 1992. On the distribution, ecology and evolution of *Gyrocotyle urna*, *G. confusa* and *G. nybelini* (Cerceromorpha, Gyrocotylidae) and their host *Chimera monostrosa* (Holocephalidae) in Norwegian waters, with a review of the species question. *Sarsia*, 77: 119–124.
- Byrkjedal, I., and Høines, A. 2007. Distribution of demersal fish in the south-western Barents Sea. *Polar Research*, 26: 135–151.
- Calis, E., Jackson, E. H., Nolan, C. P., and Jeal, F. 2005. Preliminary age and growth estimates of the rabbitfish, *Chimaera monstrosa*, with implications for future resource management. *Journal of Northwest Atlantic Fishery Science*, 35: 15–26.
- Chevolot, M., Wolfs, P. H. J., Palsson, J., Rijnsdorp, A. D., Stam, W. T., and Olsen, J. L. 2007. Population structure and historical demography of the thorny skate (*Amblyraja radiata*, Rajidae) in the North Atlantic. *Marine Biology*, 151: 1275–1286.
- Coelho, R., Figueiredo, I., Machado, P. B., and Erzini, K. 2005. Depth distribution of the velvet belly lantern shark, *Etmopterus spinax*, in southern Portugal. ICES Document CM 2005/N: 30.
- Costa, M. E., Erzini, K., and Borges, T. C. 2005. Reproductive biology of the blackmouth catshark, *Galeus melastomus* (Chondrichthyes: Scyliorhinidae) off the south coast of Portugal. *Journal of the Marine Biological Association of the UK*, 85: 1173–1183.
- Daan, N. 2001. The IBTS database: a plea for quality control. ICES Document CM 2001/T: 03. 5 pp.
- Daan, N., Heessen, H., and ter Hofstede, R. 2005. North Sea elasmobranchs: distribution, abundance and biodiversity. ICES Document CM 2005/N: 06. 15 pp.
- Dobby, H., Beare, D., Jones, E., and MacKenzie, K. 2005. Comparison of trends in long term survey data from *Squalus acanthias* with a preliminary stock assessment for this species. ICES Document CM 2005/N: 01. 19 pp.
- Dolgov, A. V. 1997. Distribution, abundance, biomass and feeding of thorny skate, *Raja radiata*, in the Barents Sea. ICES Document 1997/G: 04. 21 pp.
- Dolgov, A. V. 2006. New data on the distribution of rare and new fish species in Russian waters of the Barents Sea. *Journal of Ichthyology*, 46: 139–147.
- Dolgov, A. V., Drevetnyak, K. V., and Gusev, E. V. 2005a. The status of skate stocks in the Barents Sea. *Journal of Northwest Atlantic Fishery Science*, 35: 1–13.
- Dolgov, A. V., Drevetnyak, K. V., Sokolov, K. M., and Gusev, E. V. 2004. Skates. In *Investigations of PINRO in the Area off Spitsbergen Archipelago*, pp. 265–274. PINRO Press, Murmansk. 550 pp. (in Russian).
- Dolgov, A. V., Grekov, A. A., Shestopal, I. P., and Sokolov, K. M. 2005b. By-catch of skates in trawl and long-line fisheries in the Barents Sea. *Journal of Northwest Atlantic Fishery Science*, 35: 357–366.
- Drevetnyak, K. V., Dolgov, A. V., Sokolov, K. M., Gusev, E. V., and Grekov, A. A. 2005. Skates in the Barents Sea: stock status and catch by fishing fleet. ICES Document CM 2005/N: 11. 7 pp.
- Dulvy, N. K., Metcalfe, J. D., Glanville, J., Pawson, M. G., and Reynolds, J. D. 2000. Fishery stability, local extinctions, and shifts in community structure in skates. *Conservation Biology*, 14: 283–293.
- Ellis, J., Dulvy, N. K., Jennings, S., Parker-Humphreys, M., and Rogers, S. I. 2005a. Assessing the status of demersal elasmobranchs in UK waters: a review. *Journal of the Marine Biological Association of the UK*, 85: 1025–1047.
- Ellis, J., Clarke, M., Cortés, E., Heessen, H. J. L., Apostolaki, P., Carlson, J. K., and Kulka, D. W. 2008. Management of elasmobranch fisheries in the North Atlantic. In *Advances in Fisheries Science*. 50 years on from Beverton and Holt, pp. 184–228. Ed. by A. Payne, J. Cotter, and T. Potter. Blackwell Publishing, Oxford. 547 pp.
- Ellis, J. R., Cruz-Martinez, A., Rackham, B. D., and Rogers, S. I. 2005b. The distribution of chondrichthyan fishes around the British Isles and implications for conservation. *Journal of Northwest Atlantic Fishery Science*, 35: 195–213.
- Fossheim, M., Nilssen, E. M., and Aschan, M. 2006. Fish assemblages in the Barents Sea. *Marine Biology Research*, 2: 260–269.
- Froese, R., and Pauly, D. (Eds.) 2007. FishBase. World Wide Web Electronic Publication. www.fishbase.org
- Gyory, J., Mariano, A. J., and Ryan, E. H. 2005. The Norwegian and North Cape Currents. The Cooperative Institute for Marine and Atmospheric Studies.
- Hognestad, P. T., and Vader, W. 1979. Saltvannsfiskene i Nord-Norge (The marine fish in North Norway). *Tromsø*, 6: 1–74.
- Holden, M. J. 1967. Transatlantic movement of a tagged spurdogfish. *Nature*, 214: 1140–1141.

- Holden, M. J. 1968. The rational exploitation of Scottish–Norwegian stocks of spurdogs (*Squalus acanthias* L.). Fishery Investigations, Series II, 25. 27 pp.
- Holden, M. J. 1977. Elasmobranchs. In *Fish Population Dynamics*, pp. 187–214. Ed. by J. A. Gulland. John Wiley, New York. 384 pp.
- Hunter, E., Blickley, A. A., Stewart, C., and Metcalf, J. D. 2005. Repeated seasonal migration by a thornback ray in the southern North Sea. *Journal of the Marine Biological Association of the UK*, 85: 1199–1200.
- ICES. 2006. Report of the Working Group on Elasmobranch Fishes (WGEF). ICES Document CM 2006/ACFM: 31. 291 pp.
- ICES. 2007. Report of the Working Group on Elasmobranch Fishes (WGEF). ICES Document CM 2006/ACFM: 27. 318 pp.
- IMR. 2007. Faste hydrografiske stasjoner langs norskekysten (Permanent Hydrographical Stations along the Norwegian Coast). Institute for Marine Research, Bergen. <http://atlas.nodc.no/stasjoner/>.
- ITIS. 2008. Accessed 12/04/2008, from the Integrated Taxonomic Information System On-line Database. <http://www.itis.gov>
- Jakobsen, T., Korsbrekke, K., Mehl, S., and Nakken, O. 1997. Norwegian combined acoustic and bottom trawl surveys for demersal fish in the Barents Sea during winter. ICES Document CM 1997/Y: 17. 26 pp.
- Johnston, G., Figueiredo, I., Machado, P., Clarke, M., Bladsdale, T., Ellis, J., and Hareide, R. 2005. Separation of species data from national landing figures. ICES Document CM 2005/N: 22. 16 pp.
- Jones, E., Bare, D., Dobby, H., Trinkler, N., Burns, F., Peach, K., and Bladsdale, T. 2005. The potential impact of commercial fishing activity on the ecology of deepwater chondrichthyans from the west of Scotland. ICES Document CM 2005/N: 16. 19 pp.
- Jørgensen, O. A., Hvingel, C., Møller, P. R., and Treble, M. A. 2005. Identification and mapping of bottom fish assemblages in Davis Strait and southern Baffin Bay. *Canadian Journal of Fisheries and Aquatic Sciences*, 62: 1833–1852.
- Kotwicki, S., and Weinberg, K. L. 2005. Estimating capture probability of a survey bottom trawl for Bering Sea skates (*Bathyraja* spp.) and other fish. *Alaska Fishery Research Bulletin*, 11: 135–145.
- Magnussen, E. 2002. Demersal fish assemblages of Faroe Bank: species composition, distribution, biomass spectrum and diversity. *Marine Ecology Progress Series*, 238: 211–225.
- Mahon, R., Brown, S. K., Zwanenburg, K. C. T., Atkinson, D. B., Buja, K. R., Clafin, L., Howell, G. D., et al. 1998. Assemblages and biogeography of demersal fishes of the east coast of North America. *Canadian Journal of Fisheries and Aquatic Sciences*, 55: 1704–1738.
- Massuti, E., and Moranta, J. 2003. Demersal assemblages and depth distribution of elasmobranchs from the continental shelf and slope off the Balearic Islands (western Mediterranean). *ICES Journal of Marine Science*, 60: 753–766.
- Mjanger, H., Alsvåg, J., Hetsenes, K., Svendsen, B. V., and Wenneck, T. L. 2004. Håndbok for prøvetaking av fisk og krepsdyr (Manual for Sampling Fish and Crustaceans), version 3.15. 169 pp.
- Nakken, O. 2003. Hva betyr utkast for bestandene? (What influence does discards have on the fish stocks). In *Report from a Workshop on Discarding in Nordic Fisheries*, pp. 39–43. Ed. by J. W. Valdemarsen. Nordic Council of Ministers, Copenhagen, Tema Nord 2003. 537 pp.
- Oksanen, J. 2007. *Multivariate Analysis of Ecological Communities in R: vegan tutorial*, 39. University of Oulu. 580 pp. Available at <http://cc.oulu.fi/~jarioksa/opetus/metodi/vegantutor.pdf>
- Ottersen, G., Michalsen, K., and Nakken, O. 1998. Ambient temperature and distribution of North-east Arctic cod. *ICES Journal of Marine Science*, 55: 67–85.
- Pawson, M. G., and Ellis, J. 2005. Stock identity of elasmobranchs in the Northeast Atlantic in relation to assessment and management. *Journal of Northwest Atlantic Fishery Science*, 35: 173–193.
- Perry, A. L., Low, P. J., Ellis, J. R., and Reynolds, J. D. 2005. Climate change and distribution shifts in marine fishes. *Science*, 308: 1912–1915.
- Pethon, P. 2005. *Aschehaugs Store Fiskebok*. H. Aschehaug & Co., Oslo. 468 pp.
- Priede, I. G., Froese, R., Bailey, D. M., Bergstad, O. A., Collins, M. A., Dyb, J. E., Henriques, C., et al. 2006. The absence of sharks from abyssal regions of the world's oceans. *Proceedings of the Royal Society Series B*, 273: 1435–1441.
- R Development Core Team. 2007. R: a Language and Environment for Statistical Computing, version R 2.5.0. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>
- Reithe, S., and Aschan, M. 2004. Bioeconomic analysis of by-catch of juvenile fish in the shrimp fisheries—an evaluation of management procedures in the Barents Sea. *Environmental and Resource Economics*, 28: 55–72.
- Rey, J., de Sola, L. G., and Massuti, E. 2004. Distribution and biology of the blackmouth catshark *Galeus melastomus* in the Alboran Sea (Southwestern Mediterranean). *Journal of Northwest Atlantic Fishery Science*, 35: 215–223.
- Ruud, J. T. 1968. Changes since the turn of the century in the fish fauna and the fisheries of the Oslofjord. *Helgoländer Wissenschaftliche Meeresuntersuchungen*, 17: 510–517.
- Serena, F., Baino, R., Rey, J., Papacostantinou, C., and Relini, G. 2006. Catch composition and abundance of deep-sea elasmobranchs based on the MEDITS1 trawl surveys. In *Deep Sea 2003: Conference on the Governance and Management of Deep-sea Fisheries. Part 2: Conference Poster Papers and Workshop Papers*, Queenstown, New Zealand, 1–5 December 2003; Dunedin, New Zealand, 27–29 November 2003, pp. 395–408. Ed. by R. Shotton. *FAO Fisheries Proceedings*, 3/2. 500 pp.
- Serena, F., Mancousi, C., Barone, M., and Abella, A. J. 2005. Abundance and distribution of rays in the South Ligurian and North Tyrrhenian Sea. ICES Document CM 2005/N: 20. 16 pp.
- Skjæraasen, J. E., and Bergstad, O. A. 2000. Distribution and feeding ecology of *Raja radiata* in the northeastern North Sea and Skagerrak (Norwegian Deep). *ICES Journal of Marine Science*, 57: 1249–1260.
- Skjæraasen, J. E., and Bergstad, O. A. 2001. Notes on the distribution and length composition of *Raja lintea*, *R. fyllae*, *R. hyperborea* and *Bathyraja spinicauda* (Pisces: Rajidae) in the deep northeastern North Sea and on the slope of the eastern Norwegian Sea. *ICES Journal of Marine Science*, 58: 21–28.
- Sosebee, K. A. 2005. Are density-dependent effects on elasmobranch maturity possible? *Journal of Northwest Atlantic Fishery Science*, 35: 115–124.
- Stenberg, C. 2005. Life history of the piked dogfish (*Squalus acanthias* L.) in Swedish waters. *Journal of Northwest Atlantic Fishery Science*, 35: 155–164.
- Stevens, J. D., Bonfil, R., Dulvy, N. K., and Walker, P. A. 2000. The effects of fishing on sharks, rays, and chimaeras (chondrichthyans), and the implications for marine ecosystems. *ICES Journal of Marine Science*, 57: 476–494.
- Templeman, W. 1984. Migrations of spiny dogfish, *Squalus acanthias*, and recapture success from tagging in the Newfoundland area, 1963–65. *Journal of Northwest Atlantic Fisheries Science*, 5: 47–53.
- ter Braak, C. J. F., and Verdonschot, P. F. M. 1995. Canonical correspondence-analysis and related multivariate methods in aquatic ecology. *Aquatic Sciences*, 57: 255–289.
- Toresen, R., and Østvedt, O. J. 2000. Variations in the abundance of Norwegian spring-spawning herring (*Clupea harengus* L., Clupeidae) throughout the 20th century and the influence of climatic variations. *Fish and Fisheries*, 1: 231–256.
- Walker, P. A., and Heessen, H. J. L. 1996. Long-term changes in ray populations in the North Sea. *ICES Journal of Marine Science*, 53: 1085–1093.

- Walker, P. A., and Hislop, J. R. G. 1998. Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central and north-western North Sea between 1930 and the present day. *ICES Journal of Marine Science*, 55: 392–402.
- Williams, T. 2007. Cartilaginous fishes along the North Norwegian coast. Distributions and densities with regard to fishing and sea temperature. Masters thesis in International Fisheries Management, Norwegian College of Fishery Science, University of Tromsø. 62 pp.

doi:10.1093/icesjms/fsn103