

gaps exist. The goal was to study the impact of a multi-component, experiential, brief curriculum on attitudes of dementia care. Methods: 108 medical students participated in a curriculum including didactics, clinical, and team-based learning followed by pre-post assessments. Results: Student's perception of their ability to assess multiple facets of dementia such as behaviors, caregiver burden, and cognition improved significantly ( $p < 0.001$ ). Students' perception of the role of social worker improved significantly ( $p = 0.002$ ). Conclusion: An interdisciplinary curriculum, improved medical students' perception of their ability to assess for dementia in a cohort of predominantly rural Veterans.

## SESSION 7220 (SYMPOSIUM)

### SENSORY LOSS AND THE HEALTHCARE SYSTEM: OUTCOMES AND NAVIGATION

Chair: Nicholas Reed

Discussant: Charlotte Yeh

Communication is fundamental to patient-centered care. However, sensory impairment limits communication among older adults. Specifically, hearing impairment strains communication via degraded auditory encoding while vision impairment distresses ability to read and interpret visual cues. The presence of dual sensory impairment, defined as concurrent hearing and vision impairment, may exacerbate these effects. The potential consequences of age-related sensory loss on health care interactions and outcomes are beginning to surface in epidemiologic studies demonstrating poorer patient-provider communication, higher medical expenditures, increased risk of 30-day readmission, and longer length of stay when compared to individuals without sensory loss. Importantly, these associations may be amenable to intervention via sensory aids; however, uptake to sensory care is low. Notably, less than 20% of persons with hearing impairment have hearing aids and over 55% of Medicare Beneficiaries with reported vision problems have not had an eye examination in the prior year. Affordability and access may contribute to lack of sensory care uptake as Medicare explicitly excludes coverage of vision and hearing services. In this symposium, we will review current and new evidence for whether sensory loss affects health care outcomes, including satisfaction with care, incident delirium during hospitalization, navigation of Medicare, and present data on how persons with sensory loss are more likely to delay their care independent of cost and insurance factors suggesting fundamental changes in health care system interaction. We will place these results within the context of current national quality care and policy initiatives and review methods to address sensory loss.

### HEARING LOSS AND HELP-SEEKING BEHAVIOR

Nicholas Reed, *Johns Hopkins University, Baltimore, Maryland, United States*

Hearing Loss (HL) is common among older adults and is associated with poor health care quality outcomes include 30-day readmissions, length of stay, poorer satisfaction, and increased medical expenditures. These associations

may manifest in changes in help-seeking behaviour. In the 2015 Current Medicare Beneficiary Study (MCBS) ( $n = 10848$ ; weighted sample = 46.3 million), participants reported whether they knowingly had avoided seeking care in the past year and self-reported HL was measured as degree of trouble (none, a little, or a lot) hearing when using a hearing aid if applicable. In a model adjusted for demographic, socioeconomic, and health factors, those with a little trouble (OR = 1.612; 95% CI = 1.334-1.947;  $P < 0.001$ ) and a lot of trouble hearing (OR = 2.011; 95% CI = 1.443-2.801;  $P < 0.001$ ) had 61.2% and 101.1% higher odds of avoiding health care over the past year relative to participants with no trouble hearing. Future work should examine whether hearing care modifies this association.

### SENSORY LOSS AND DELIRIUM AMONG MEDICARE BENEFICIARIES

Emmanuel Garcia Morales,<sup>1</sup> and Nicholas Reed,<sup>2</sup> *1. Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, United States, 2. Johns Hopkins University, Baltimore, Maryland, United States*

Sensory impairment is prevalent among older adults and may increase risk for delirium via mechanisms including sensory deprivation and poor communication which may result in confusion and agitation. In the Medicare Current Beneficiary Study (MCBS), delirium was measured using a validated algorithm of claims data. Sensory impairment was defined as any self-reported trouble hearing or seeing, with the use of aids, and was categorized as no impairment, hearing impairment only (HI), vision impairment only (VI), and dual sensory impairment (DSI). Among, 3,240 hospitalized participants in 2016-2017, 346 (10.7%) experienced delirium. In a model adjusted for socio-demographic and health characteristics, those with HI only, VI only, and DSI had 0.84 (95% CI: 0.6-1.3), 1.1 (95% CI 0.7-1.7), and 1.5 (95% CI 1.0-2.1) times the odds of experiencing delirium compared to those without sensory impairment. Future research should focus on mechanisms underlying association and determine the impact of treatment of sensory loss.

### UNDERSTANDING MEDICARE WITH HEARING LOSS

Amber Willink, *The University of Sydney, Sydney, New South Wales, Australia*

Medicare has become an increasingly complex program to navigate with numerous choices available to beneficiaries with important implications on their financial exposure and access to care. While research has identified poor health literacy as a barrier to understanding Medicare, little information is available on the experience of individuals with hearing loss. Using the Medicare Current Beneficiary Survey (2016), a nationally-representative sample of 10,841 beneficiaries, we examined if difficulty understanding Medicare was associated with reported trouble hearing, while controlling for socio-demographic and health literacy factors. Compared to no trouble, Medicare beneficiaries with a little or a lot of trouble hearing had 44% (95% CI OR: 1.34-1.55) and 63% (95% CI OR: 1.44-1.83) increased odds of reporting greater