

Systematic Review

Barriers to evidence-based practice implementation in physiotherapy: a systematic review and meta-analysis

MATTEO PACI¹, GIANNI FAEDDA², ALESSANDRO UGOLINI³, and LEONARDO PELLICCIARI⁴

¹Unit of Functional Rehabilitation, Azienda USL Toscana Centro, Viale Michelangiolo, 41, Florence 50125, Italy,

²Graduate Course in Rehabilitation Sciences, University of Florence, Largo Brambilla, 3, Florence 50134, Italy,

³Program in Physical Therapy, University of Florence, Largo Brambilla, 3, Florence 50134, Italy, and ⁴IRCCS Fondazione Don Carlo Gnocchi, Via di Scandicci, 269 I, Florence 50143, Italy

Address reprint requests to: Matteo Paci, Unit of Functional Rehabilitation, Azienda USL Toscana Centro, Viale Michelangiolo, 41, Florence 50125, Italy. Tel: +390556937265; E-mail: matteo.paci@applicazione.it

Received 25 October 2020; Editorial Decision 25 May 2021; Revised 9 April 2021; Accepted 10 June 2021

Abstract

Background: To review and meta-analyse the evidence about the prevalence of barriers to evidence-based practice (EBP) reported in physiotherapy.

Methods: Two independent investigators conducted an extensive electronic search in EMBASE, PubMed, Scopus, Web of Science and CINAHL databases from their inception to July 2020 and included the retrieved articles if they investigated barriers to EBP among physiotherapy professionals. Subsequently, they extracted data and assessed the methodological quality using a scale described in a similar previous study. The outcome for meta-analysis was frequency of each reported barrier. Sub-analyses were performed grouping studies based on countries where surveys were performed, classified as either developed or developing countries.

Results: Twenty-nine articles were included in the systematic reviews and meta-analysis. Risk of bias assessment of included studies showed a median score: 4 points (interquartile range: 3–4). The findings of meta-analysis revealed that lack of time was the most frequently reported barrier (53.0% [95% confidence interval, 95%CI, 44.0–62.0]), followed by language (36.0% [95%CI 16.0–62.0]), lack of access (34.0% [95%CI 23.0–27]) and lack of statistical skills (31.0% [95%CI 20.0–44.0]). Lack of skills and lack of generalizability were declared as barriers by 27.0% [95%CI 18.0–38.0] and 23.0% [95%CI 15.0–33.0] of responders, respectively. Lack of support and lack of interest are less frequent, with 16.0% [95%CI 11.0–24.0] and 9.0% [95%CI 6.0–15.0] of responses, respectively. Barriers reported in investigations performed in developed countries were less frequent when compared to those performed in developing countries.

Conclusion: Organizational issues and methodological skills seem key issues to allow the implementation of EBP, suggesting the need to adopt or enhance organizational and training strategies to facilitate the implementation of the EBP. Quantitative synthesis showed high heterogeneity for all analyses, and therefore, pooled data should be interpreted with caution.

Key words: evidence-based practice, physiotherapy, barriers, systematic review, rehabilitation

Introduction

Evidence-based practice (EBP) was defined as integrating the best research evidence with clinical expertise and patient values [1], considering the environmental (e.g. local health laws) and organizational (e.g. available resources) context [2]. Its implementation requires specific skills (e.g. literature search and critical appraisal for validity, clinical relevance and applicability) and assigned resources (e.g. accessibility to database and full text of articles) [2]. EBP is widely considered a standard of care throughout all of health care because it can be able to promote individualization of care and support the quality of health care [3]. At the same time, there are concerns about its application and use in clinical practice [4, 5].

Barriers to EBP can be considered as modifiable factors against the EBP implementation, and their knowledge could help to improve environmental and organizational context, as well as to enhance education. At the same time, improving education, environmental and organizational context, can be able to change attitudes and beliefs about EBP.

A number of surveys [6–9] and reviews [10, 11] were conducted in order to investigate barriers to EBP. The most common barriers to EBP are found to be lack of resources, lack of time, inadequate skills, inadequate access, lack of knowledge and financial barriers [10]. In a more recent review, patient overload and lack of personal time, knowledge and skills rather than a lack of facilities and resources are reported as main barriers [12].

In physiotherapy, two reviews summarized literature about the attitudes and use of EBP [9, 13], reporting inability to understand statistics, generalization of results and lack of time, support, resources and interest as main barriers to implement EBP. However, these reviews included articles published until 2012 [13] and 2014 [9] and provided results in a descriptive way, without a quantitative synthesis (e.g. meta-analysis); moreover, one of these [13] was limited to investigation performed in Western regions. Specific challenges for health research in developing countries have been stressed, in terms of lack of research culture, limited research education and limited access to research information [14], and, in the past years, few surveys were conducted in low- and middle-income countries [10].

In recent years, the dissemination of resources to access to literature may be increased (e.g. widespread use of the Internet and open access to literature) and several surveys addressed to physiotherapists were published, especially in developing countries. In addition, despite data obtained from primary research can be meta-analyzed in reviews of observational studies statistics of prevalence, no previous quantitative literature synthesis on barriers to EBP in physiotherapy has been performed.

Therefore, the aim was to perform a systematic review with meta-analysis on the prevalence of reported barriers to EBP in physiotherapy, exploring differences between developed and developing countries.

Methods

The reporting of this systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses checklist [15] (Supplementary Appendix 1).

Identification and selection of studies

An electronic search was conducted to retrieve complete original studies, reporting on barriers that restrain the EBP implementation in physiotherapy.

Studies were searched in PubMed, Scopus, Web of Science, EMBASE and CINAHL from their inception to July 2020. Full research strings for each database are reported in Supplementary Appendix 2.

To ensure retrieval of all potentially relevant publications, reference lists of related and included articles were checked by two independent reviewers (G.F. and M.P.).

Firstly, two independent reviewers (G.F. and M.P.) excluded irrelevant articles by reading the title and the abstract; then, the full texts of eligible articles were retrieved and assessed by the same reviewers for their inclusion. Disagreement was resolved by the involvement of a third reviewer (L.P.).

Studies were considered for inclusion if they met the following criteria: surveys on barriers in the use of EBP by physiotherapy professionals, published in English and Italian, involving different categories of health-care professionals only when it was possible to extract distinct data referred to physiotherapists. Studies on physiotherapy students and on the effectiveness of interventions to increase EBP skills, abstracts from conference proceedings and qualitative studies were excluded.

Methodological quality assessment

Risk of bias was assessed using the scale described by da Silva *et al.* [9], using criteria developed by Ferreira *et al.* [16] and Leboeuf-Yde and Lauritsen [17]. It consists of six criteria describing representativeness of the sample (three criteria) and quality of data (three criteria). The total score, summing up the number of criteria met, ranges from 0 (higher risk of bias) to 6 (lower risk of bias) (Supplementary Appendix 3). The risk of bias was rated by two independent reviewers (G.F. and L.P.), and disagreements were resolved by a third reviewer (M.P.).

Statistical analysis

Data from the eligible studies were combined to compute a pooled prevalence for the outcomes. Barriers were grouped in homogeneous areas to combine extracted frequencies: ‘generalizability,’ ‘interest,’ ‘general skills’ (e.g. searching for evidence or critically appraise an article), ‘statistical skills,’ ‘support’ (i.e. inadequate support from colleagues, managers and other health professionals), ‘access’ (i.e. access to electronic databases and articles’ full texts), ‘time’ and ‘language.’

The generalized linear mixed model (GLMM), a random intercept logistic regression model for the meta-analysis of proportions, was used for the meta-analysis [18]. GLMM uses the maximum likelihood estimation for the Tau-squared calculation and the logit transformation, without the reported weight of each study. We reported results through forest plots including both fixed and random effects.

Statistical heterogeneity was tested with statistical Q and its P-value, while it was quantified with Tau² and I² [19]. To reduce the heterogeneity, sub-group analyses were performed for each barrier, grouping studies based on countries where surveys were performed, classified as either developed or developing countries according to the International Statistical Institute (<https://www.isi-web.org/capacity-building/developing-countries>). Additional sub-analyses were conducted by grouping studies considering the continents where surveys were conducted, methodological quality (score <4 versus the others) and year of publication (before or after 2014, the year of the last published review in the physiotherapy field).

The publication bias was assessed using the funnel-plot graph method and the statistic test proposed by Thompson and Sharp [20],




Figure 1 Flowchart of study selection.

based on a weighted linear regression of the effect on its standard error using the method of moments estimator for the additive between-study variance component. A P -value < 0.05 indicated significant publication bias.

Meta-analyses were performed using the software R and the ‘meta’ package [21].

If the frequencies of the demographic data of the included studies were not reported for all the samples, the missing data were reported in the table and the frequencies were recomputed. Finally, the scores of the risk of bias assessment of studies conducted in developed and developing countries were compared using the Mann–Whitney test. Descriptive and non-parametric statistics were calculated using SPSS software (Version 23 for Windows; SPSS Inc., Chicago, IL, USA).

Results

Study inclusion and characteristics

Of 3602 titles, 29 articles fulfilled the selection criteria [22–50] (participants = 9337, mean number of participants per article \pm standard deviation = 338.0 ± 341.4) (Figure 1). The main characteristics of the study samples are reported in Table 1.

Lack of time and lack of skills were the most reported barriers, reported in 27 [22–32, 34–50] and 24 [22–31, 33–38, 40–42, 44–46, 49, 50] of the included studies, followed by lack of access ($n = 23$) [22, 24–31, 33–35, 37, 38, 40, 41, 43–47, 49, 50], lack of support ($n = 23$) [22, 24–30, 33–35, 37–41, 44–50], lack of interest ($n = 21$) [23, 24, 26–29, 31, 34–36, 38–42, 44–46, 48–50], lack

of generalizability ($n = 19$) [22, 24–27, 29, 31–36, 40, 42, 44, 45, 48–50] and lack of statistical skills ($n = 17$) [22, 24–26, 29, 31–36, 42, 44, 45, 48–50]. Language was reported as a barrier in few studies ($n = 8$) [30, 31, 34–36, 42, 43, 49]. Studies were performed in 15 countries from 6 different continents: 9 in Asia [29, 30, 32, 33, 38, 40, 43, 45, 46], 6 in South America [34–36, 42, 47, 48], 5 in North America [22, 24, 26, 27, 41] and Europe [23, 28, 31, 39, 49], 3 in Africa [37, 44, 50] and 1 in Oceania [25]. Response rate ranged from 17.5% [31] to 95.0% [37] (Table 2).

Fourteen studies were included in the sub-group developing countries [29, 32, 34–37, 40, 42–44, 46–48, 50]. Response rate ranged from 28.8% [40] to 95.0% [37]. Fifteen studies were conducted in developed countries [22–28, 30, 31, 33, 38, 39, 41, 45, 49], with a response rate ranging from 17.5% [31] to 92.0% [45] (Table 2).

Methodological quality

The score ranged from 2 to 6 points (median = 4; interquartile range = 3–4) for all included studies. For developed countries, the total score varied between 2 and 6 points (median = 4; interquartile range = 3–4.5), while for developing countries the total score ranged between 3 and 5 points (median = 3.5; interquartile range = 3–4); the result of Mann–Whitney test revealed no significant difference between the scores of studies conducted in developed and developing countries. ‘Data collection’ item was satisfied by all articles, while ‘non-responders’ and ‘power calculation’ were the less frequently

Table 1 Main characteristics of the sample in the included studies

Reference and country	Age		Gender		Place of employment		Education		Professional experience		Working hours per week		Patients per day	
	Years	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	Years	n (%)	Hours	n (%)	n (%)	n (%)	
Metcalfe et al., 2001 [22] ^a USA	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Kamwendo et al., 2002 [23] Sweden	Age for male: 36.9 ± 7.8; age for female: 41.3 ± 10.2	63 (18.4) 279 (81.3) 1 (0.3)	Male Female Missing	NR	Private outpatient clinic Facility-based outpatient clinic Acute care hospital Home care School system Skilled nursing facility Acute rehabilitation Subacute rehabilitation University Other	157 (32.2) 118 (24.2) 62 (12.7) 36 (7.4) 26 (5.3) 23 (4.7) 20 (4.1) 14 (2.9) 7 (1.4) 8 (1.6) 17 (3.5)	Baccalaureate Professional master's Advanced master's Professional doctorate Advanced doctorate Other Missing	188 (38.6) 187 (38.4) 80 (16.4) 10 (2.0) 10 (2.0) 3 (0.6) 10 (2.0)	Experience for male: 10.3 ± 6.8; experience for female: 14.1 ± 10.1	NR	NR	NR	NR	NR
Jette et al., 2003 [24] USA	20–29 30–39 40–49 50+ Missing	99 (20.3) 160 (32.9) 149 (30.5) 71 (14.5) 9 (1.8)	Male Female Missing	NR	Private outpatient clinic Facility-based outpatient clinic Acute care hospital Home care School system Skilled nursing facility Acute rehabilitation Subacute rehabilitation University Other	141 (28.9) 338 (69.3) 9 (1.8)	Baccalaureate Professional master's Advanced master's Professional doctorate Advanced doctorate Other Missing	188 (38.6) 187 (38.4) 80 (16.4) 10 (2.0) 10 (2.0) 3 (0.6) 10 (2.0)	Experience for male: 10.3 ± 6.8; experience for female: 14.1 ± 10.1	NR	NR	NR	NR	NR
Grimmer-Somers et al., 2007 [25] Australia	NR	NR	NR	NR	Private practice Public or private hospital Aged care or disability services Government department or university Missing	96 (57.8) 44 (26.5) 14 (8.4) 11 (6.6) 1 (0.7)	Bachelor's or equivalent Master's Postgraduate diploma Honours degree Missing	100 (60.2) 30 (18.1) 17 (10.2) 13 (7.8) 6 (3.7)	Mean time ± SD 16.8 ± 10.6 (range 0.5–47)	NR	NR	NR	NR	NR
Salbach et al., 2007 [26] Canada	20–29 30–39 40–49 50+ Missing	40 (14.8) 93 (34.4) 75 (27.8) 60 (22.2) 2 (0.8)	Male Female Missing	30 (11.1) 239 (88.5) 1 (0.4)	Acute care hospital Rehabilitation hospital Private practice Home visiting agency Community care access center Long-term care facility Complex continuing care Community health center University Other	106 (39.3) 43 (15.9) 28 (10.4) 17 (6.3) 14 (5.2) 13 (4.8) 10 (3.7) 3 (1.1) 1 (0.4) 33 (12.2) 2 (0.7)	Bachelor's Certificate/diploma Applied or research master's Professional master's Missing	203 (75.2) 30 (11.1) 23 (8.5) 8 (3.0) 6 (2.2)	<5 5–10 11–15 >15 Missing	40 (14.9) 59 (21.9) 48 (17.8) 122 (45.2) 1 (0.4)	<20 20–30 31–40 >40 Missing	28 (10.4) 51 (18.9) 154 (57.0) 35 (13.0) 2 (0.7)	1–10 11–15 >15 Missing 3 (1.1)	
Fruth et al., 2010 [27] ^b USA	20–29 30–39 40–49 50+ Missing	7 (29.2) 10 (41.7) 4 (16.7) 3 (12.5) 3 (12.5)	Male Female	9 (25.0) 18 (75.0)	Missing Hospital-based inpatient Hospital-based general outpatient Private practice outpatient orthopedic Long-term neurological rehabilitation	11 (45.8) 7 (29.2) 3 (12.5) 3 (12.5)	Master's Bachelor's Doctor of physical therapy Transitional doctor of physical therapy	11 (45.8) 7 (29.2) 3 (12.5) 3 (12.5)	<6 6–10 11–15 15+	7 (29.2) 6 (25.0) 5 (20.8) 6 (25.0)	NR	NR	NR	NR

Continued.

Table 1 (Continued)

Reference and country	Age		Gender		Place of employment		Education		Professional experience		Working hours per week		Patients per day	
	Years	n (%)	Male	n (%)	Primary care Hospital Company health service University Health maintenance and preventive care services Other	n (%)	<3-year ground education 3-year education Master's Specialist Master's 2 years Missing	Years	n (%)	Hours	n (%)	<5 5-10 11-15 Missing	NR	n (%)
Nilsågård and Lohse, 2010 [28] Sweden	NR	133 (16.0) 700 (84.0)	Male Female	133 (16.0) 700 (84.0)	Primary care Hospital Company health service University Health maintenance and preventive care services Other	255 (30.6) 183 (22.0) 46 (5.5) 26 (3.1) 11 (1.3) 77 (9.3)	<3-year ground education 3-year education Master's Specialist Master's 2 years Missing	<14 14-29 >30	120 (14.4) 393 (47.2) 320 (38.4)	NR	NR	<5 5-10 11-15 >15	NR	NR
Gorgon <i>et al.</i> , 2013 [29] Philippines	20-29 30-39	120 (79.0) 32 (21.0)	Male	51 (33.6%) 101 (66.4%)	Government tertiary hospital Private tertiary hospital	53 (34.9) 99 (65.1)	Bachelor's Master's	<5 5-10 11-15 Missing	92 (60.6) 49 (32.2) 9 (5.9) 2 (1.3)	<20 21-30 31-40 >40	13 (8.6) 5 (3.3) 56 (36.8) 78 (51.3)	<5 5-10 11-15 >15	NR	25 (16.4) 93 (61.2) 20 (13.2) 14 (9.2)
Weng <i>et al.</i> , 2013* [30] Taiwan	NR	NR	NR	NR	Private tertiary hospital	NR	NR	NR	NR	NR	NR	NR	NR	NR
Diermayr <i>et al.</i> , 2015 [31] Austria	20-29 30-39 40-49 50+	83 (14.1) 189 (32.1) 168 (28.6) 99 (16.8)	Male Female Missing	97 (16.5) 438 (74.5) 53 (9.0)	Private practice Shared private practice Acute care hospital Rehabilitation center Home care Outpatient clinic University of applied sciences Home for the elderly Missing	199 (33.8) 95 (16.2) 83 (14.1) 41 (7.0) 37 (6.3) 31 (5.3) 16 (2.7) 12 (2.0) 74 (12.6)	Diploma physical therapy Master/Master's student Bachelor's Dr/PhD/doctoral student Missing	<5 5-10 11-15 >15	69 (11.7) 103 (17.5) 115 (19.6) 252 (42.9) 49 (8.3)	<20 20-30 31-40 >40 Missing	64 (10.9) 175 (29.8) 158 (26.9) 141 (24.0) 50 (8.4)	<5 5-10 11-15 >15 Missing	66 (11.2) 294 (50.0) 120 (20.4) 51 (8.7) 57 (9.7)	
Panhale and Bellare [32] India	Mean age: 27	13 (22.0) 47 (78.0)	Male Female	13 (22.0) 47 (78.0)	NR	NR	NR	NR	48 (80.0) 12 (20.0)	NR	NR	NR	NR	NR
Park <i>et al.</i> , 2015 [33] Korea	<24 25-29 30-34 35+	19 (12.0) 61 (38.6) 52 (32.9) 26 (16.5)	Male Female	103 (65.2) 53 (34.8)	General hospital University hospital Rehabilitation hospital Clinic	81 (51.3) 51 (32.3) 21 (13.3) 5 (3.2)	University Graduate school or more College	1-5 Missing	44 (73.0) 12 (20.0) 4 (7.0)	NR	NR	NR	NR	NR
Ramirez-Vélez <i>et al.</i> , 2015 [34] Ramirez-Vélez <i>et al.</i> , 2015 [35] Colombia	20-29 30-39 40-49 50+	845 (79.4) 155 (14.6) 58 (5.5) 6 (0.6)	Male Female	243 (22.8) 821 (77.2)	Private outpatient clinic Acute care hospital Facility-based outpatient clinic Skilled nursing facility University Acute rehabilitation Home care Sub-acute rehabilitation School system Other	412 (38.7) 175 (16.4) 124 (11.7) 91 (8.6) 82 (7.7) 55 (5.2) 36 (3.4) 34 (3.2) 7 (0.7) 48 (4.5)	Undergraduate Specialized Master's Doctorate	<5 5-10 11-15 >15	342 (32.1) 450 (42.3) 128 (12) 144 (13.5)	<20 20-30 31-40 >40	114 (10.7) 256 (24.1) 349 (32.8) 345 (32.4)	<5 5-10 11-15 >15	159 (14.9) 327 (30.7) 284 (26.7) 294 (27.6)	
Silva <i>et al.</i> , 2015 [36] Brazil	NR	49 (19.1) 207 (80.9)	Male Female	49 (19.1) 207 (80.9)	Self-employed Private Public More than one	129 (50.4) 84 (32.8) 32 (12.5) 11 (4.3)	Bachelor's Master's by coursework Master's by research Post-doctoral	<5 5-9 10-14 15-19 20-24 >24	99 (38.6) 81 (31.6) 45 (17.6) 13 (5.1) 5 (2.0)	NR	NR	NR	NR	NR
Wanjiru <i>et al.</i> , 2016 [37] Kenya	Mean age: 30.5 (range 27-63)	17 (42.5) 23 (57.5)	Male Female	17 (42.5) 23 (57.5)	NR	NR	Diploma Higher diploma Degree Master's	<5 5-10 11-20 21-30 31-40 >40	26 (65.0) 6 (15.0) 8 (20.0) 9 (22.5) 4 (10.0) 2 (5.0)	NR	NR	NR	NR	NR

Continued.

Table 1 (Continued)

Reference and country	Age		Gender		Place of employment		Education		Professional experience		Working hours per week		Patients per day	
	Years	n (%)	Male	n (%)	Ministry of health	n (%)	Bachelor's	n (%)	Years	n (%)	Hours	n (%)	n (%)	n (%)
Alshetri et al., 2017 [38] Saudi Arabia	20–25	118 (31.4)	Male	227 (60.4)	Ministry of health	153 (40.7)	Bachelor's	249 (66.2)	<3	67 (29.3)	<20	NR	NR	NR
	26–30	119 (31.6)	Female	149 (39.6)	Private health sectors	62 (16.5)	Master's	75 (19.9)	3–5	30 (13.1)	20–30	NR	NR	NR
	31–35	73 (19.4)			University	61 (16.2)	Diploma	18 (4.8)	6–10	50 (21.8)	31–40	NR	NR	NR
	36–40	34 (9.0)			Other health sectors	100 (26.6)	Doctorate	31 (8.2)	11–15	23 (10.0)	>40	NR	NR	NR
	>40	32 (8.5)				Doctor of physical therapy	3 (0.8)	16–20	23 (10.0)					
Krutulyte et al., 2017 [39] Lithuania	20–29	106 (46.3)		NR	NR		NR	NR	<3	67 (29.3)	<20	NR	NR	NR
	30–39	58 (25.3)							3–5	30 (13.1)	20–30	NR	NR	NR
	40–49	29 (12.7)							6–10	50 (21.8)	31–40	NR	NR	NR
	50+	36 (15.7)							11–15	23 (10.0)	>40	NR	NR	NR
Yahui and Swaminathan, 2017 [40] Malaysia	25–34	87 (85.3)	Male	27 (26.5)	Private clinic	33 (32.4)	Bachelor's	54 (52.9)	2–5	59 (57.8)	<20	10 (9.8)	NR	NR
	35–44	11 (10.8)	Female	75 (73.5)	Private hospital	30 (29.4)	Diploma certificate	45 (44.1)	>5	24 (23.5)	20–30	10 (9.8)	NR	NR
	45–54	3 (2.9)			Government hospital	22 (21.6)	Master's	1 (1.0)	>10	7 (6.9)	31–40	34 (33.3)	NR	NR
	55+	1 (1.0)			Own practice	9 (8.8)	Doctorate	2 (2.0)	>15	7 (6.9)	>40	48 (47.1)	NR	NR
				University	4 (3.9)			>20	5 (4.9)					
				Home care	3 (2.9)									
				Government health clinics	1 (1.0)									
Gardin and Hudson, 2018 [41] ^a USA	NR	NR		NR	NR		NR	NR	NR	NR	NR	NR	NR	NR
	NR	NR	Male	12 (11.9)	Self-employed	46 (45.5)	Master's of coursework	60 (59.4)	<5	32 (31.7)	NR	NR	NR	NR
	NR	NR	Female	89 (88.1)	Private	34 (33.7)	Master's of research	23 (22.8)	5–9	25 (24.8)	NR	NR	NR	NR
	NR	NR			Public	11 (10.9)	Bachelor's	11 (10.8)	10–14	28 (27.7)	NR	NR	NR	NR
Claudio et al., 2018 [42] Brazil	NR	NR			Both	10 (9.9)	Doctorate	4 (4.0)	15–19	9 (8.8)	NR	NR	NR	NR
	NR	NR							20–24	4 (4.0)	NR	NR	NR	NR
	NR	NR	Male	146 (38.3)	NR				>24	3 (3.0)	NR	NR	NR	NR
	NR	NR	Female	235 (61.7)					<6	116 (30.4)	NR	NR	NR	NR
Dao and Pichaiyongwongdee, 2018 [43] Vietnam	20–29	138 (36.2)	Male	146 (38.3)	NR				6–9	88 (23.1)	NR	NR	NR	NR
	30–39	159 (41.7)	Female	235 (61.7)					10–14	79 (20.7)	NR	NR	NR	NR
	40–49	61 (16.0)							15+	98 (25.8)	NR	NR	NR	NR
	50+	23 (6.1)									NR	NR	NR	NR
Quarrey and Kwakye, 2018 [44] Ghana	20–25	16 (13.2)	Male	74 (61.2)	General hospital	103 (85.1)	doctorate	111 (91.7)	<5	45 (37.1)	NR	NR	NR	NR
	26–30	63 (52.1)	Female	47 (38.8)	University	11 (9.2)	Bachelor's	6 (5.0)	5–8	68 (56.2)	NR	NR	NR	NR
	31–35	29 (24.0)			Private practice	5 (4.1)	Applied or research master's	4 (3.3)	>8	8 (6.6)	NR	NR	NR	NR
	36–40	13 (10.7)			Rehabilitation hospital	1 (0.8)	Entry level master's				NR	NR	NR	NR
Alrowayeh and Baabbas, 2019 [45] Kuwait	20–29	42 (22.9)	Male	72 (40.2)	General hospital	70 (38.0)	Bachelor's	131 (71.2)	0–5	33 (17.9)	<10	22 (12.0)	1–7	92 (50.0)
	30–39	79 (42.9)	Female	108 (58.7)	Rehabilitation hospital	69 (37.5)	Master's	48 (26.1)	6–10	32 (17.4)	10–19	5 (2.6)	8–12	69 (37.5)
	40–49	50 (27.2)	Missing	2 (1.1)	Specialized hospital	43 (23.4)	Missing	5 (2.7)	11–15	38 (20.7)	20–29	38 (20.7)	>12	11 (6.0)
	>50	12 (6.5)			Missing	2 (1.1)			16–20	38 (20.7)	30–39	62 (33.7)	Missing	12 (6.2)
	Missing	1 (0.5)						>20	33 (17.9)	>40	45 (24.5)			
Bajracharya et al., 2019 [46] Nepal	20–30	129 (78.7)	Male	78 (47.6)	Acute care hospital	89 (54.3)	NR	NR	Missing	10 (5.4)	Missing	12 (6.5)	<5	14 (8.5)
	31–40	27 (16.5)	Female	86 (52.4)	Missing	75 (45.7)			<2	65 (39.6)	<20	6 (3.7)	<5	67 (40.9)
	41–50	3 (1.8)							2–5	55 (33.5)	20–30	11 (6.7)	5–10	44 (26.8)
	>50	5 (3.0)							6–10	30 (18.3)	31–40	36 (22)	11–15	39 (23.8)
								>10	14 (8.5)	>40	111 (67)	>15	39 (23.8)	

Continued.

Table 1 (Continued)

Reference and country	Age		Gender		Place of employment		Education		Professional experience		Working hours per week		Patients per day	
	Years	n (%)	Male	n (%)		n (%)		n (%)	Years	n (%)	Hours	n (%)		n (%)
Cobo-Sevilla <i>et al.</i> , 2019 [47] Ecuador	20–29	24 (35.8)	Male	25 (37.3)	Acute care	55 (82.0)	Bachelor's	0–5	24 (35.8)	<20	7 (10.5)	<5	1 (1.5)	
	30–39	31 (46.2)	Female	42 (62.7)	Outpatient	6 (9.0)	Graduate studies	6–10	18 (26.9)	20–30	2 (3.0)	5–10	4 (6.0)	
	40–49	6 (9.0)			Private office	2 (3.0)	Technologist	11–15	16 (23.9)	31–40	24 (35.8)	11–15	5 (7.5)	
	>50	6 (9.0)			Private outpatient	2 (3.0)	Missing	>15	9 (13.4)	>40	34 (50.7)	>15	57 (85.0)	
Nascimento <i>et al.</i> , 2019 [48] Brazil	20–29	63 (38.4)	Male	40 (24.4)	Private long-term facility	74 (45.1)	Specialization	<5	65 (39.6)	NR	NR	<5	12 (7.3)	
	30–39	79 (48.1)	Female	122 (74.4)	Hospital	24 (14.6)	Bachelor's	5–10	61 (37.2)			5–10	45 (27.4)	
	40–49	16 (9.8)	Missing	2 (1.2)	University	23 (14.0)	Master's	11–15	21 (12.8)			11–15	18 (11.0)	
	50+	6 (3.7)			Public long-term facility	19 (11.6)	Doctorate	>15	15 (9.1)			>15	17 (10.4)	
Castellini <i>et al.</i> , 2020 [49] ^a Italy	<29	361 (28.0)	Male	619 (48.0)	Rehabilitation hospital	12 (7.3)	Missing	Missing	2 (1.2)			Missing	72 (43.9)	
	29–38	400 (31.0)	Female	670 (52.0)	Community care access center	6 (3.7)								
	39–49	232 (18.0)			Missing	6 (3.7)								
	>49	296 (23.0)			Private office	657 (59.0)	NR	NR	NR				NR	
Ibikunle <i>et al.</i> , 2020 [50] Nigeria	20–29	98 (57.6)	Male	106 (62.4)	Hospital	223 (20.0)	Bachelor's	0–4	97 (57.1)	<20	34 (20.0)	<5	23 (13.5)	
	30–39	58 (34.1)	Female	64 (37.6)	General hospital	65 (38.2)	Entry-level master's	5–9	39 (22.9)	20–30	32 (18.8)	5–10	110 (64.7)	
	40–49	11 (6.5)			Private practice	20 (11.8)	Applied or research master's	10–14	26 (15.3)	31–40	52 (30.6)	11–15	24 (14.1)	
	50+	3 (1.8)			Rehabilitation hospital	8 (4.7)	Doctoral	15–19	5 (2.9)	>40	52 (30.6)	>15	13 (7.6)	
				Consulting firm	7 (4.1)									
				Specialist hospital	5 (2.9)									
				Federal medical center	3 (1.8)									
				Tertiary hospital	1 (0.6)									

Abbreviations: n, number; %, percentages; NR, not reported; SD, standard deviation.

^a Characteristics were not available for physical therapists but only for the total sample (including other health professionals).

^b Demographic data are reported for 24 subjects; barriers data are reported for 43 subjects including test and retest.

^c For place of employment, percentage exceed 100% because this question allowed more than one answer in the original study.

Table 2 Total sample size and response rate of the included studies

Study	Total sample (responded—response rate)
Metcalfe <i>et al.</i> , 2001 [22]	348 (287—82.5%)
Kamwendo <i>et al.</i> , 2002 [23]	556 (343—61.7%)
Jette <i>et al.</i> , 2003 [24]	1000 (488—48.8%)
Grimmer-Somers <i>et al.</i> , 2007 [25]	335 (166—49.5%)
Salbach <i>et al.</i> , 2007 [26]	334 (270—80.8%)
Fruth <i>et al.</i> , 2010 [27]	NR (43—NR)
Nilsagård and Lohse, 2010 [28]	2160 (833—38.6%)
Gorgon <i>et al.</i> , 2013 [29]	188 (150—79.8%)
Weng <i>et al.</i> , 2013 [30]	NR (149—NR)
Diermayr <i>et al.</i> , 2015 [31]	3360 (588—17.5%)
Panhale and Bellare [32]	100 (60—60.0%)
Park <i>et al.</i> , 2015 [33]	200 (158—79.0%)
Ramírez-Vélez <i>et al.</i> , 2015 [34]	1250 (1064—85.1%)
Ramírez-Vélez <i>et al.</i> , 2015 [35]	1250 (1064—85.1%)
Silva <i>et al.</i> , 2015 [36]	490 (256—52.2%)
Wanjiru <i>et al.</i> , 2016 [37]	42 (40—95%)
Alshehri <i>et al.</i> , 2017 [38]	604 (376—62.2%)
Krutulytė <i>et al.</i> , 2017 [39]	300 (229—76.3%)
Yahui and Swaminathan, 2017 [40]	354 (102—28.8%)
Cardin and Hudson, 2018 [41]	NR (30—NR)
Claudino <i>et al.</i> , 2018 [42]	250 (101—40.4%)
Dao and Pichaiyongwongdee, 2018 [43]	453 (381—84.1%)
Quartey and Kwakye, 2018 [44]	130 (121—93.0%)
Alrowayeh and Buabbas, 2019 [45]	200 (184—92.0%)
Bajracharya <i>et al.</i> , 2019 [46]	NR (164—NR)
Cobo-Sevilla <i>et al.</i> , 2019 [47]	NR (67—NR)
Nascimento <i>et al.</i> , 2019 [48]	260 (164—63.1%)
Castellini <i>et al.</i> , 2020 [49]	2000 (1289—64.5%)
Ibikunle <i>et al.</i> , 2020 [50]	220 (170—85%)

Abbreviation: NR, not reported.

satisfied items (only by 4 [13.8%] and 7 [24.1%] articles, respectively) (Table 3).

Meta-analyses

Pooled prevalence, with 95% confidence interval (95% CI), of each barrier for both the whole sample and studies conducted in developed and developing countries is reported in Table 4. The forest plots for each barrier are reported in Supplementary Appendix 4.

Lack of time was the most frequently reported barrier (53.0% [95%CI=44.0–62.0]), followed by language (36.0% [95%CI=16.0–62.0]), lack of access (34.0% [95%CI=23.0–47.0]) and lack of statistical skills (31.0% [95%CI=20.0–44.0]). Lack of skills and lack of generalizability were declared as barriers by 27.0% [95%CI=18.0–38.0] and 23.0% [95%CI=15.0–33.0] of responders, respectively. Lack of support and lack of interest are less frequent, with 16.0% [95%CI=11.0–24.0] and 9.0% [95%CI=6.0–15.0] of responses, respectively.

With the exception of the group ‘generalizability,’ barriers reported in investigations performed in developed countries were less frequent when compared to those performed in developing countries, especially for ‘time,’ ‘general skills,’ ‘statistical skills,’ ‘access’ and ‘language’ (Table 4, Supplementary Appendix 4).

Heterogeneity was found to be high for all analyses, ranging from 98% to 99% (Table 4). All additional sub-group analyses, according to continents (i.e. Europe, Africa, Asia, Oceania, North and South America), methodological quality (i.e. best [≥ 4 points] or worst [< 4 points] quality) and year of publication (i.e. articles published before and after 2014) were not able to explain heterogeneity.

The *P*-value for publication bias was significant for meta-analyses regarding barriers on time ($P=0.007$), skills ($P=0.004$), interest ($P=0.003$) and statistical skills ($P=0.027$) but not for the other barriers (Supplementary Appendix 5).

Discussion

Statement of principal findings

Barriers to the EBP implementation in the clinical practice are related to the organizational context (lack of time, access and support), education (language, lack of research and statistical skills), personal behavior (lack of interest) and limits of EBP (lack of generalizability).

These results are similar to those reported in previous reviews [9, 13]. This systematic review reports quantitative pooled data on each barrier also grouped based on countries where surveys were performed (developed/developing countries).

Quantitative synthesis shows high heterogeneity for all analyses, and therefore, pooled data should be interpreted with caution.

Strengths and limitations

To our knowledge, this systematic review reports, for the first time, quantitative pooled data on barriers to EBP implementation in the field of physiotherapy and compares surveys conducted in developed and developing countries. Moreover, a large number of surveys were included ($n=29$), balanced between developed ($n=15$) and developing ($n=14$) countries.

Twenty studies (70.4%) were published after 2014, and all surveys conducted in developing countries, with the exception of Gorgon *et al.* [29], as well as some studies published before 2014 [22, 23, 27, 30] were not included in previous systematic reviews [9, 13]. In addition, quantitative synthesis allows to highlight the high heterogeneity among studies.

However, this study also presented limitations that deserve to be discussed. A potential limitation is that grouping barriers in eight areas was an arbitrary choice. For example, ‘lack of support’ includes support from colleagues, managers and other health professionals. However, an external and standardized criterion for grouping the results is not available in the literature, and we tried to put together different items in a coherent way. Secondly, detection of frequencies of barriers is related to the structure of used questionnaires. Surveys include a finite set of pre-determined items that assess the individual importance of each barrier and might not include all potential factors [51]. Moreover, interaction among different barriers was not investigated, despite it might have synergistic effects on EBP implementation [51]. In addition, the scale used to assess the risk of bias, proposed by da Silva *et al.* [9], could lack content validity (i.e. it does not include specific aspects related to the validity of the measurements and analyses); however, no consensus is available in literature on the best to use to evaluate the methodological quality of surveys; therefore, future researches in this topic should be implemented. Finally, the search was restricted to studies in English and Italian.

Interpretation within the context of the wider literature

Barriers referred to organizational context built a complex area with a potential interaction among single barriers, without relevant differences across countries. Lack of time is reported as a barrier by >50% of responders. It should be recognized that ‘lack of time’ may include different barriers, as lack of willingness to change clinical

Table 3 Quality assessment of the studies included in the systematic review

Study	Description of the sample	Non-responders	Response rate	Data collection	Questionnaire validated	Power calculation	Total score*
<i>Developed countries</i>							
Metcalfe <i>et al.</i> , 2001 [22]	+	-	+	+	+	-	4
Kamwendo <i>et al.</i> , 2002 [23]	+	-	+	+	+	-	4
Jette <i>et al.</i> , 2003 [24]	+	-	+	+	+	-	4
Grimmer-Somers <i>et al.</i> , 2007 [25]	+	-	+	+	+	+	5
Salbach <i>et al.</i> , 2007 [26]	+	+	+	+	-	-	4
Fruth <i>et al.</i> , 2010 [27]	+	-	-	+	-	-	2
Nilsagård and Lohse, 2010 [28]	+	-	+	+	-	-	3
Weng <i>et al.</i> , 2013 [30]	+	-	-	+	+	-	3
Diermayr <i>et al.</i> , 2015 [31]	+	+	+	+	+	-	5
Park <i>et al.</i> , 2015 [33]	+	-	+	+	-	-	3
Alshehri <i>et al.</i> , 2017 [38]	+	+	+	+	+	+	6
Krutulytė <i>et al.</i> , 2017 [39]	+	-	+	+	+	-	4
Cardin and Hudson, 2018 [41]	+	-	-	+	+	-	3
Alrowayeh and Buabbas, 2019 [45]	+	-	+	+	+	-	4
Castellini <i>et al.</i> , 2020 [49]	+	-	+	+	+	+	5
<i>Developing countries</i>							
Gorgon <i>et al.</i> , 2013 [29]	+	-	+	+	-	-	3
Ramírez-Vélez <i>et al.</i> , 2015 [34]	+	-	-	+	+	-	3
Ramírez-Vélez <i>et al.</i> , 2015 [35]	+	-	-	+	+	-	3
Silva <i>et al.</i> , 2015 [36]	+	-	+	+	+	+	5
Panhale and Bellare, 2015 [32]	-	-	+	+	-	-	2
Wanjiru <i>et al.</i> , 2016 [37]	+	-	+	+	-	-	3
Yahui and Swaminathan, 2017 [40]	+	-	+	+	+	-	4
Claudino <i>et al.</i> , 2018 [42]	+	-	+	+	-	-	3
Dao and Pichaiyongwongdee, 2018 [43]	+	-	+	+	+	-	4
Quartey and Kwakye, 2018 [44]	+	-	+	+	-	-	3
Bajracharya <i>et al.</i> , 2019 [46]	+	-	-	+	+	+	4
Cobo-Sevilla <i>et al.</i> , 2019 [47]	+	-	+	+	+	+	5
Nascimento <i>et al.</i> , 2019 [48]	+	+	+	+	+	-	5
Ibikunle <i>et al.</i> , 2020 [50]	+	-	+	+	-	+	4

Notes: Total score ranges from 0 (worst quality) from 6 (best quality).

Table 4 Results of the meta-analyses for study conducted in developed and developing countries and for all included studies

Study	Time	Skills	Access	Support	Interest	Generalizability	Statistical skills	Language
<i>Developed countries</i>								
Metcalfe et al., 2001 [22]	0.62 [0.56; 0.67]	0.67 [0.61; 0.72]	0.45 [0.39; 0.51]	0.37 [0.31; 0.43]	0.05 [0.03; 0.07]	0.50 [0.44; 0.56]	0.80 [0.75; 0.85]	
Kamwendo et al., 2002 [23]	0.52 [0.47; 0.58]	0.02 [0.01; 0.04]	0.02 [0.01; 0.04]	0.03 [0.02; 0.05]	0.01 [0.00; 0.02]	0.09 [0.07; 0.12]	0.04 [0.02; 0.06]	
Jette et al., 2003 [24]	0.46 [0.41; 0.51]	0.03 [0.02; 0.05]	0.02 [0.01; 0.04]	0.25 [0.18; 0.32]	0.04 [0.02; 0.07]	0.20 [0.14; 0.27]	0.54 [0.46; 0.62]	
Grimmer-Somers et al., 2007 [25]	0.31 [0.24; 0.39]	0.43 [0.35; 0.51]	0.20 [0.15; 0.27]	0.19 [0.08; 0.33]	0.03 [0.02; 0.06]	0.34 [0.28; 0.40]	0.30 [0.25; 0.36]	
Salbach et al., 2007 [26]	0.74 [0.69; 0.80]	0.31 [0.25; 0.37]	0.16 [0.12; 0.21]	0.04 [0.02; 0.07]	0.09 [0.03; 0.22]	0.23 [0.12; 0.39]		
Fruth et al., 2010 [27]	0.84 [0.69; 0.93]	0.26 [0.14; 0.41]	0.40 [0.25; 0.56]	0.19 [0.08; 0.33]	0.46 [0.43; 0.49]			
Nilsagård and Lohse, 2010 [28]	0.86 [0.83; 0.88]	0.55 [0.52; 0.58]	0.80 [0.77; 0.83]					0.61 [0.53; 0.69]
Weng et al., 2013 [30]	0.60 [0.52; 0.68]	0.36 [0.29; 0.45]	0.62 [0.54; 0.70]	0.12 [0.07; 0.18]	0.05 [0.03; 0.07]	0.08 [0.06; 0.10]	0.15 [0.12; 0.18]	0.07 [0.05; 0.09]
Diermayr et al., 2015 [31]	0.22 [0.18; 0.25]	0.06 [0.04; 0.08]	0.09 [0.07; 0.11]	0.09 [0.07; 0.12]		0.64 [0.56; 0.71]		
Park et al., 2015 [33]	0.30 [0.23; 0.38]	0.39 [0.31; 0.47]	0.58 [0.50; 0.66]	0.39 [0.31; 0.47]				
Alshehri et al., 2017 [38]	0.23 [0.19; 0.27]	0.36 [0.32; 0.42]	0.36 [0.31; 0.41]	0.31 [0.26; 0.36]				
Krutulytė et al., 2017 [39]	0.30 [0.24; 0.36]			0.07 [0.04; 0.11]				
Cardin and Hudson, 2018 [41]	0.37 [0.20; 0.56]	0.10 [0.02; 0.27]	0.20 [0.08; 0.39]	0.17 [0.06; 0.35]				
Alrowayeh and Buabbas, 2019 [45]	0.59 [0.52; 0.66]	0.07 [0.04; 0.12]	0.49 [0.41; 0.56]	0.02 [0.01; 0.05]		0.41 [0.34; 0.48]	0.03 [0.01; 0.07]	
Castellini et al., 2020 [49]	0.23 [0.21; 0.25]	0.09 [0.08; 0.11]	0.08 [0.07; 0.10]	0.12 [0.10; 0.14]		0.11 [0.09; 0.13]	0.10 [0.08; 0.11]	0.05 [0.04; 0.06]
Total (fixed effects)	0.45 [0.43; 0.46]	0.24 [0.23; 0.25]	0.30 [0.29; 0.32]	0.14 [0.13; 0.15]		0.20 [0.19; 0.21]	0.21 [0.19; 0.22]	0.10 [0.08; 0.11]
Total (random effects)	0.48 [0.36; 0.60]	0.19 [0.10; 0.33]	0.28 [0.16; 0.46]	0.12 [0.07; 0.20]	0.07 [0.03; 0.13]	0.25 [0.14; 0.39]	0.22 [0.09; 0.46]	0.15 [0.03; 0.51]
I ² (heterogeneity, %)	98	99	99	97	98	98	99	99
<i>Developing countries</i>								
Gorgon et al., 2013 [29]	0.85 [0.78; 0.90]	0.40 [0.32; 0.48]	0.44 [0.36; 0.52]	0.27 [0.20; 0.34]	0.16 [0.11; 0.23]	0.26 [0.19; 0.34]	0.44 [0.36; 0.52]	
Panhale and Bellare, 2015 [32]	0.32 [0.20; 0.45]					0.18 [0.10; 0.30]	0.18 [0.10; 0.30]	
Ramírez-Vélez et al., 2015 [34]	0.35 [0.32; 0.38]	0.41 [0.38; 0.44]	0.09 [0.07; 0.11]	0.03 [0.02; 0.04]	0.01 [0.01; 0.02]	0.02 [0.01; 0.03]	0.39 [0.36; 0.42]	0.22 [0.19; 0.25]
Ramírez-Vélez et al., 2015 [35]	0.44 [0.41; 0.47]	0.56 [0.53; 0.59]	0.20 [0.18; 0.23]	0.15 [0.13; 0.17]	0.11 [0.09; 0.13]	10.1 (8.3–12.0)	0.53 [0.50; 0.56]	0.33 [0.30; 0.36]
Silva et al., 2015 [36]	0.61 [0.55; 0.67]	0.41 [0.35; 0.47]			0.28 [0.23; 0.34]	0.55 [0.49; 0.62]	0.54 [0.48; 0.61]	0.70 [0.64; 0.76]
Wanjiru et al., 2016 [37]	0.80 [0.64; 0.91]	0.72 [0.56; 0.85]	0.85 [0.70; 0.94]	0.75 [0.59; 0.87]				
Yahui and Swaminathan, 2017 [40]	0.28 [0.20; 0.38]	0.17 [0.10; 0.25]	0.17 [0.10; 0.25]	0.16 [0.09; 0.24]	0.14 [0.08; 0.22]	0.21 [0.13; 0.30]	0.54 [0.44; 0.64]	0.47 [0.37; 0.57]
Claudio et al., 2018 [42]	0.45 [0.35; 0.55]	0.42 [0.32; 0.52]			0.28 [0.19; 0.38]	0.61 [0.51; 0.71]		0.84 [0.80; 0.88]
Dao and Pichaiyongwongdee, 2018 [43]	0.72 [0.67; 0.76]		0.73 [0.68; 0.77]					
Quarrey and Kwakye, 2018 [44]	0.46 [0.37; 0.56]	0.18 [0.12; 0.26]	0.43 [0.34; 0.52]	0.09 [0.05; 0.16]	0.33 [0.25; 0.42]	0.15 [0.09; 0.22]	0.36 [0.27; 0.45]	
Bajracharya et al., 2019 [46]	0.67 [0.59; 0.74]	0.63 [0.55; 0.70]	0.53 [0.45; 0.61]	0.34 [0.27; 0.42]	0.24 [0.17; 0.31]			
Cobo-Sevilla et al., 2019 [47]	0.96 [0.87; 0.99]		0.54 [0.41; 0.66]	0.51 [0.38; 0.63]				
Nascimento et al., 2019 [48]	0.62 [0.54; 0.70]			0.32 [0.25; 0.39]	0.05 [0.02; 0.09]	0.33 [0.26; 0.41]	0.32 [0.25; 0.39]	
Ibikunle et al., 2020 [50]	0.48 [0.40; 0.55]	0.31 [0.24; 0.38]	0.43 [0.35; 0.51]	0.14 [0.09; 0.20]	0.13 [0.08; 0.19]	0.26 [0.19; 0.33]	0.28 [0.21; 0.35]	0.39 [0.38; 0.41]
Total (fixed effects)	0.49 [0.48; 0.51]	0.45 [0.43; 0.47]	0.29 [0.27; 0.30]	0.15 [0.13; 0.16]	0.11 [0.10; 0.12]	0.16 [0.15; 0.17]	0.44 [0.42; 0.46]	0.52 [0.30; 0.73]
Total (random effects)	0.59 [0.46; 0.71]	0.41 [0.30; 0.52]	0.42 [0.26; 0.59]	0.22 [0.12; 0.38]	0.13 [0.07; 0.23]	0.22 [0.12; 0.36]	0.40 [0.32; 0.48]	0.52 [0.30; 0.73]
I ² (heterogeneity, %)	98	97	98	98	97	98	94	99
Total (fixed effects)	0.47 [0.46; 0.48]	0.32 [0.31; 0.33]	0.30 [0.29; 0.31]	0.14 [0.14; 0.15]	0.13 [0.13; 0.14]	0.18 [0.17; 0.19]	0.32 [0.31; 0.33]	0.27 [0.26; 0.28]
Total (random effects)	0.53 [0.44; 0.62]	0.27 [0.18; 0.38]	0.34 [0.23; 0.47]	0.16 [0.11; 0.24]	0.09 [0.06; 0.15]	0.23 [0.15; 0.33]	0.31 [0.20; 0.44]	0.36 [0.16; 0.62]
I ² (inconsistency, %)	98	99	99	98	98	98	99	99

practice or low priority of learning EBP principles over practical skills [52]. Indeed, ideally, lack of time should be perceived as a barrier only when skills and access are satisfied, but it seems to be not true. Lack of access to research resources was reported as a barrier by more than one-third of responders, with a higher frequency in developing countries. Limited access to information has been found to be a relevant barrier also for nurses working in low- and middle-income countries [53]. However, since access restrictions can be also due to the high number of subscription only databases and journals and access to these resources are frequently provided by academic institutions, physiotherapists freelance or working in non-academic institutions can have problems to EBP access across all countries [54]. For example, apart from the Hinari database sponsored by the World Health Organization, many Nigerian public health establishments do not subscribe to other research databases [50]. The theme 'lack of support' includes different types of support (i.e. inadequate support from colleagues, managers and other health professionals) at different levels (i.e. individual, workplace and extra-organizational levels) [55]. This is not a frequently reported barrier, mainly in developed countries, although it could be ascribed, at least partially, to other reported barriers as lack of time or lack of access, because they also depend on organization restrictions.

Barriers referred to education are more frequently reported in surveys performed in developing countries. Language is frequently reported as a barrier, although it was investigated only in some studies conducted in non-native English countries, reaching to be perceived as a barrier for >50% of responders in developing countries. This issue was not analyzed in previous reviews, probably as seven of the eight studies are published after 2014. This issue is relevant since English is currently the predominant language in the field of biomedical publications [56], and language seems to be a relevant barrier in developing countries. Lack of research skills is reported by nearly one-third of responders. This group of skills includes finding the best evidence, formulating a relevant and clear question, evaluating evidences and applying them and combining them with clinical experience and patient values [57]. Furthermore, inability to understand statistical data was reported by more than one-third of responders. Both these types of barriers are more frequently reported in developing countries, suggesting specific needs in these areas [10, 14]. Since the level of academic education seems to be related to EBP acknowledgement [58], the lower ability in research in developing countries can be explained by a poor availability of advanced education (e.g. doctorate) or specific EBP-related training. For example, it has been reported that there were few or no postgraduate degree and no doctorate programs in Kuwait [59] and Saudi Arabia [38].

The lack of interest is confirmed as a rarely declared barrier, and this result links to the finding that physiotherapists generally have a positive opinion about EBP [9]. Anyway, the so-called 'eminence-based therapy' (i.e. when clinical experience entails established reputations of credibility despite the high risk of bias) [60] seems to be still highly valued in physiotherapy [61], and a possible mismatch between reported attitudes and daily practice should be considered. Education and interest in EBP are strictly related. Undergraduate education is a key occasion to implement and develop methodological skills, attitudes and interest [62].

Lack of generalizability of research results is reported as a frequent barrier in all countries. This issue can effectively be relevant [63] and can explain the underuse in the routine practice of research literature [64, 65]. However, it is the sole issue investigated on limits of EBP, despite other objections to EBP have been reported [66].

High heterogeneity was found for all analyses, despite sub-group analyses, according to the type of country, continent, methodological quality and publication year, not clarifying on the heterogeneity. This result is likely to be expected since studies included in this study differ in cultural, economic and social contexts, as in main characteristics of the included subjects, especially the place of employment, education and professional experience (Table 1). Furthermore, this result could also be due to different physiotherapy education programs and health-care systems across countries. Moreover, the diversity of questionnaires used in the surveys, or aspects of the quality of the studies not captured by the checklist used, could further justify the heterogeneity. Two reviews [67, 68] found 15 and 34 different survey instruments testing EBP knowledge, skills, attitudes and behaviors, and barriers. Additional sources of variability may be related to different operational definitions or diverse subjective perceptions of the respondents. For example, Castellini *et al.* [49], reported that 52% of responders stated they understood the meaning of 'meta-analysis' term, but only 17% and 38% were able to explain 'forest plot' and 'confidence intervals' terms, respectively.

Moreover, we found a publication bias in pooled meta-analysis for four out of eight barriers (i.e. time, skills, interest and statistical skills) that could concur with the high heterogeneity. However, in our analyses, we considered the random effect model that accounts for unexplained heterogeneity [69].

The results of our meta-analysis revealed a high heterogeneity across studies that should be considered for future investigations; hence, guidelines and checklists about this type of study should be produced to minimize the heterogeneity for the studies; for example, a consensus on how the questionnaire should be developed should fill the agenda for future researches, as the checklist development for assessing the methodological quality of these types of studies. Finally, our results should be interpreted with caution, providing initial quantitative results on this topic.

Included investigations were assessed to have a low-to-moderate risk of bias. However, response rate is <50% in six investigations [24, 25, 28, 31, 40, 42], and it is not reported in five studies [27, 30, 41, 46, 47]. Literature on non-response bias through mailed surveys shows that non-response bias is a serious concern in survey studies [70] that could affect the interval validity of surveys.

Implications for policy, practice and research

Since barriers to the EBP implementation in the clinical practice are mainly related to organizational and educational factors, university trainings, continuing education providers, health policymakers and health-care companies should further support research educations and activities. Particularly, universities should implement programs to increase the lack of statistical and research skills and courses in scientific English; health-care organizations should contemplate time for EBP (i.e. reading articles and journal clubs) and provide access and support; finally, research should provide evidence with such generalizability that they can be used in clinical practice. Furthermore, research can stimulate a scientific method of approach to problems for the health care, management and organization of clinical services and practices. Indeed, it is well known that a research culture is associated with greater service efficiency and efficacy from clinical and organizational perspectives [71].

Finally, a consensus on structure and core set of areas of investigation should be developed to compare results coming from studies using different survey instruments.

Conclusion

Organizational strategies and methodological skills seem key issues to allow the EBP implementation, although results should be interpreted with caution, considering the low response rate of primary studies and the high heterogeneity. Therefore, our data suggest the need to adopt or enhance organizational and training strategies aimed at providing opportunities and tools to facilitate the EBP implementation. Barriers referred to education seem to be an important EBP obstacle in developing countries. It is necessary to explore effective methods to support the EBP adoption by students and physiotherapists [72], especially in developing countries.

Supplementary material

Supplementary material is available at *International Journal for Quality in Health Care* online.

Funding

This study was not funded.

Contributorship

M.P. and G.F. were primarily responsible for study conception and design and for data analysis and interpretation. All authors (M.P., G.F., A.U. and L.P.) participated in the phases of the systematic review. M.P. and A.U. analyzed the data. All authors (M.P., G.F., A.U. and L.P.) interpreted the results. M.P. led the writing of the first draft of the manuscript. All authors (M.P., G.F., A.U. and L.P.) contributed to drafting and revising the manuscript. M.P. is the guarantor. All authors (M.P., G.F., A.U. and L.P.) read and approved the final manuscript.

Ethics and other permissions

There were no permissions needed to undertake this work.

Data availability statement

No new data were generated or analyzed in support of this review.

References

- Sackett DL, Rosenberg WM, Gray JA *et al.* Evidence based medicine: what it is and what it isn't. *BMJ* 1996;312:71–2.
- Dawes M, Summerskill W, Glasziou P *et al.* Sicily statement on evidence-based practice. *BMC Med Educ* 2005;5:1.
- Evidence-Based Medicine Working Group. Evidence-based medicine. A new approach to teaching the practice of medicine. *JAMA* 1992;268:2420–5.
- Naylor CD. Grey zones of clinical practice: some limits to evidence-based medicine. *Lancet* 1995;345:840–2.
- Straus SE, McAlister FA. Evidence-based medicine: a commentary on common criticisms. *CMAJ* 2000;163:837–41.
- González-Torrente S, Pericas-Beltrán J, Bennasar-Veny M *et al.* Perception of evidence-based practice and the professional environment of primary health care nurses in the Spanish context: a cross-sectional study. *BMC Health Serv Res* 2012;12:227.
- Durr ES. How to apply evidence-based principles in clinical dentistry. *J Multidiscip Healthc* 2019;12:131–6.
- Weiss CH. Why do we fail to deliver evidence-based practice in critical care medicine? *Curr Opin Crit Care* 2017;23:400–5.
- da Silva TM, Costa Lda C, Garcia AN *et al.* What do physical therapists think about evidence-based practice? A systematic review. *Man Ther* 2015;20:388–401.
- Sadeghi-Bazargani H, Tabrizi JS, Azam-Aghdash S. Barriers to evidence-based medicine: a systematic review. *J Eval Clin Pract* 2014;20:793–802.
- Zwolsman S, Te Pas E, Hooft L *et al.* Barriers to GPs' use of evidence-based medicine: a systematic review. *Br J Gen Pract* 2012;62:e511–21.
- Barzkar F, Baradaran HR, Koohpayehzadeh J. Knowledge, attitudes and practice of physicians toward evidence-based medicine: a systematic review. *J Evid Based Med* 2018;11:246–51.
- Scurlock-Evans L, Upton P, Upton D. Evidence-based practice in physiotherapy: a systematic review of barriers, enablers and interventions. *Physiotherapy* 2014;100:208–19.
- Rahman MM, Ghoshal UC, Raguath K *et al.* Biomedical research in developing countries: opportunities, methods, and challenges. *Indian J Gastroenterol* 2020;39:292–302.
- Moher D, Liberati A, Tetzlaff J *et al.* Preferred Reporting Items for Systematic Reviews and Meta-Analyses: the PRISMA statement. *Ann Int Med* 2009;151:264–9, w64.
- Ferreira ML, Machado G, Latimer J *et al.* Factors defining care-seeking in low back pain—a meta-analysis of population based surveys. *Eur J Pain* 2010;14:747.e1–7.
- Leboeuf-Yde C, Lauritsen JM. The prevalence of low back pain in the literature. A structured review of 26 Nordic studies from 1954 to 1993. *Spine* 1995;20:2112–8.
- Stijnen T, Hamza TH, Ozdemir P. Random effects meta-analysis of event outcome in the framework of the generalized linear mixed model with applications in sparse data. *Stat Med* 2010;29:3046–67.
- Higgins JP, Thomas J, Chandler J *et al.* *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester (UK): John Wiley & Sons, 2019.
- Thompson SG, Sharp SJ. Explaining heterogeneity in meta-analysis: a comparison of methods. *Stat Med* 1999;18:2693–708.
- Schwarzer G, Carpenter JR, Rücker G. *Meta-analysis with R*. Cham, Switzerland: Springer, 2015.
- Metcalfe C, Lewin R, Wisher S *et al.* Barriers to implementing the evidence base in four NHS therapies: dietitians, occupational therapists, physiotherapists, speech and language therapists. *Physiotherapy* 2001; 87:433–41.
- Kamwendo K. What do Swedish physiotherapists feel about research? A survey of perceptions, attitudes, intentions and engagement. *Physiother Res Int* 2002;7:23–34.
- Jette DU, Bacon K, Batty C *et al.* Evidence-based practice: beliefs, attitudes, knowledge, and behaviors of physical therapists. *Phys Ther* 2003;83:786–805.
- Grimmer-Somers K, Lekkas P, Nyland L *et al.* Perspectives on research evidence and clinical practice: a survey of Australian physiotherapists. *Physiother Res Int* 2007;12:147–61.
- Salbach NM, Jaglal SB, Korner-Bitensky N *et al.* Practitioner and organizational barriers to evidence-based practice of physical therapists for people with stroke. *Phys Ther* 2007;87: 1284–303.
- Fruth SJ, Van Veld RD, Despos CA *et al.* The influence of a topic-specific, research-based presentation on physical therapists' beliefs and practices regarding evidence-based practice. *Physiother Theory Pract* 2010;26:537–57.
- Nilsagård Y, Lohse G. Evidence-based physiotherapy: a survey of knowledge, behaviour, attitudes and prerequisites. *Adv Physiother* 2010;12:179–86.
- Gorgon EJ, Barrozo HG, Mariano LG *et al.* Research evidence uptake in a developing country: a survey of attitudes, education and self-efficacy, engagement, and barriers among physical therapists in the Philippines. *J Eval Clin Pract* 2013;19:782–90.
- Weng YH, Kuo KN, Yang CY *et al.* Implementation of evidence-based practice across medical, nursing, pharmacological and allied healthcare professionals: a questionnaire survey in nationwide hospital settings. *Implement Sci* 2013;8:112.

31. Diermayr G, Schachner H, Eidenberger M *et al.* Evidence-based practice in physical therapy in Austria: current state and factors associated with EBP engagement. *J Eval Clin Pract* 2015;21:1219–34.
32. Panhale VP, Bellare B. Evidence-based practice among physiotherapy practitioners in Mumbai, India. *Educ Health* 2015;28:154–5.
33. Park K, Choi B, Han D. Clinical physical therapists research activity reality and barriers to their utilizing research findings. *J Phys Ther Sci* 2015;27:2091–5.
34. Ramírez-Vélez R, Correa-Bautista JE, Muñoz-Rodríguez DI *et al.* Evidence-based practice: beliefs, attitudes, knowledge, and skills among Colombian physical therapists. *Colomb Med* 2015;46:33–40.
35. Ramírez-Vélez R, Bagur-Calafat MC, Correa-Bautista JE *et al.* Barriers against incorporating evidence-based practice in physical therapy in Colombia: current state and factors associated. *BMC Med Educ* 2015;15:220.
36. Silva TM, Costa LC, Costa LO. Evidence-based practice: a survey regarding behavior, knowledge, skills, resources, opinions and perceived barriers of Brazilian physical therapists from São Paulo state. *Braz J Phys Ther* 2015;19:294–303.
37. Wanjiru N, Kabara S, Milimo J. Factors affecting implementation of evidence based practice among physiotherapists in Moi teaching referral hospital, Kenya. *Int J Physiother* 2016;3:267–72.
38. Alshehri MA, Alalawi A, Alhasan H *et al.* Physiotherapists' behaviour, attitudes, awareness, knowledge and barriers in relation to evidence-based practice implementation in Saudi Arabia: a cross-sectional study. *Int J Evid Based Healthc* 2017;15:127–41.
39. Krutulytė G, Lendraitienė E, Petruševičienė D *et al.* Evidence-based physiotherapy practice in Lithuania: attitudes, application, and barriers. *Fizjoterapia Polska* 2017;17:42–49.
40. Yahui HC, Swaminathan N. Knowledge, attitudes, and barriers towards evidence-based practice among physiotherapists in Malaysia. *Hong Kong Physiother J* 2017;37:10–8.
41. Cardin AD, Hudson MB. Evidence-based practice in the hospital setting: views of interdisciplinary therapy practitioners. *J Allied Health* 2018;47:81–9.
42. Claudino R, De Pietro Simoes N, Da Silva T. Evidence-based practice: a survey of Brazilian physical therapists from the dermatology subdiscipline. *Braz J Phys Ther* 2019;23:395–401.
43. Dao HT, Pichaiyongwongdee S. Are physical therapists in Viet Nam ready to implement evidence-based practice? A survey. *BMC Med Educ* 2018;18:317.
44. Quartey J, Kwakye S. Barriers to evidence-based physiotherapy practice for stroke survivors in Ghana. *S Afr J Physiother* 2018;74:423.
45. Alwayeh HN, Buabbas AJ. Evidence-based physical therapy practice in the state of Kuwait: a survey of attitudes, beliefs, knowledge, skills, and barriers. *JMIR Med Educ* 2019;5:e12795.
46. Bajracharya S, Tharu NS, Bokali D. Knowledge, attitude and barrier to evidence-based practice among physiotherapists in selected districts of Nepal. *J Nepal Health Res Counc* 2019;17:215–21.
47. Cobo-Sevilla V, De Oliveira-Ferreira I, Moposita-Baño L *et al.* Evidence-based physiotherapy clinical practice in the public health-care service in Ecuador. *Physiother Res Int* 2019;24:e1745.
48. Nascimento LR, Fernandes MOP, Teixeira-Salmela LF *et al.* Personal and organizational characteristics associated with evidence-based practice reported by Brazilian physical therapists providing service to people with stroke: a cross-sectional mail survey. *Braz J Phys Ther* 2020;24:349–57.
49. Castellini G, Corbetta D. Twenty-five years after the introduction of evidence-based medicine: knowledge, use, attitudes and barriers among physiotherapists in Italy - a cross-sectional study. *BMJ Open* 2020;10:e037133.
50. Ibikunle PO, Onwuakagba IU, Maduka EU *et al.* Perceived barriers to evidence-based practice in stroke management among physiotherapists in a developing country. *J Eval Clin Pract* 2021;27:291–306.
51. Nilsen P, Bernhardsson S. Towards evidence-based physiotherapy - research challenges and needs. *J Physiother* 2013;59:143–4.
52. Stander J, Grimmer K, Brink Y. Time as a barrier to evidence uptake—a qualitative exploration of the concept of time for clinical practice guideline uptake by physiotherapists. *J Eval Clin Pract* 2021;27:280–90.
53. Shayan SJ, Kiwanuka F. Barriers associated with evidence-based practice among nurses in low- and middle-income countries: a systematic review. *Worldviews Evid Based Nurs* 2019;16:12–20.
54. Herbert R, Jamtvedt G, Hagen KB *et al.* *Practical Evidence-Based Physiotherapy-E-Book*. London: Elsevier Health Sciences, 2011.
55. Dannapfel P, Peolsson A, Nilsen P. What supports physiotherapists' use of research in clinical practice? A qualitative study in Sweden. *Implement Sci* 2013;8:31.
56. Valkimadi P, Karageorgopoulos D, Vliagoftis H *et al.* Increasing dominance of English in publications archived by PubMed. *Scientometrics* 2009;81:219–23.
57. Sackett DL, Straus SE, Richardson WS *et al.* *Evidence-Based Medicine: Teach EBM*. 2nd ed. Edinburgh: Churchill Livingstone, 2000.
58. Mikhail C, Korner-Bitensky N, Rossignol M *et al.* Physical therapists' use of interventions with high evidence of effectiveness in the management of a hypothetical typical patient with acute low back pain. *Phys Ther* 2005;85:1151–67.
59. Al-Enezi L, May S. Why do physiotherapists do what they do? A study of Kuwaiti physiotherapists. *Physiother Res Int* 2017;22:e1640.
60. Isaacs D, Fitzgerald D. Seven alternatives to evidence based medicine. *BMJ* 1999;319:1618.
61. Bø K, Herbert RD. When and how should new therapies become routine clinical practice? *Physiotherapy* 2009;95:51–7.
62. Olsen NR, Bradley P, Lomborg K *et al.* Evidence based practice in clinical physiotherapy education: a qualitative interpretive description. *BMC Med Educ* 2013;13:52.
63. Paci M, Prestera C, Ferrarello F. Generalizability of results from randomized controlled trials in post-stroke physiotherapy. *Physiother Can* 2020.
64. Rothwell PM. External validity of randomised controlled trials: “to whom do the results of this trial apply?” *Lancet* 2005;365:82–93.
65. Uhlig K, Leff B, Kent D *et al.* A framework for crafting clinical practice guidelines that are relevant to the care and management of people with multimorbidity. *J Gen Intern Med* 2014;29:670–9.
66. Herbert RD, Sherrington C, Maher C *et al.* Evidence-based practice—imperfect but necessary. *Physiother Theory Pract* 2001;17:201–11.
67. Glegg SM, Holsti L. Measures of knowledge and skills for evidence-based practice: a systematic review. *Can J Occup Ther* 2010;77:219–32.
68. Buchanan H, Siegfried N, Jelsma J. Survey instruments for knowledge, skills, attitudes and behaviour related to evidence-based practice in occupational therapy: a systematic review. *Occup Ther Int* 2016;23:59–90.
69. Riley RD, Higgins JP, Deeks JJ. Interpretation of random effects meta-analyses. *BMJ* 2011;342:d549.
70. Groves RM, Cialdini RB, Couper MP. Understanding the decision to participate in a survey. *Public Opin Q* 1992;56:475–95.
71. Harding K, Lynch L, Porter J *et al.* Organisational benefits of a strong research culture in a health service: a systematic review. *Aust Health Rev* 2017;41:45–53.
72. Stander J, Grimmer K, Brink Y. Training programmes to improve evidence uptake and utilisation by physiotherapists: a systematic scoping review. *BMC Med Educ* 2018;18:14.