

Spectropolarimetry of the Type Ia SN 2019ein rules out significant global asphericity of the ejecta

Kishore C. Patra¹,¹★† Yi Yang (杨轶),^{1,2}★‡ Thomas G. Brink,¹ Peter Höflich,³ Lifan Wang,⁴ Alexei V. Filippenko,^{1,5}§ Daniel Kasen,^{1,6} Dietrich Baade,⁷ Ryan J. Foley,⁸ Justyn R. Maund⁹, WeiKang Zheng,¹ Tiara Hung,⁸ Aleksandar Cikota¹⁰, J. Craig Wheeler¹¹ and Mattia Bulla¹²

¹Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

²Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot 76100, Israel

³Department of Physics, Florida State University, 77 Chieftan Way, Tallahassee, FL 32306, USA

⁴George P. and Cynthia Woods Mitchell Institute for Fundamental Physics & Astronomy, Texas A&M University, 4242 TAMU, College Station, TX 77843, USA

⁵Miller Institute for Basic Research in Science, University of California, Berkeley, CA 94720, USA

⁶Departments of Physics and Astronomy and Lawrence Berkeley National Laboratory, University of California, Berkeley, CA 94720, USA

⁷European Organisation for Astronomical Research in the Southern Hemisphere (ESO), Karl-Schwarzschild-Str 2, D-85748 Garching b. München, Germany

⁸Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA

⁹Department of Physics and Astronomy, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, UK

¹⁰European Southern Observatory, Alonso de Córdova 3107, Vitacura, Casilla 19001, Santiago de Chile, Chile

¹¹Department of Astronomy, University of Texas, Austin, TX 78712, USA

¹²The Oskar Klein Centre, Department of Astronomy, Stockholm University, AlbaNova, SE-10691 Stockholm, Sweden

Accepted 2021 October 25. Received 2021 October 25; in original form 2021 August 11

ABSTRACT

Detailed spectropolarimetric studies may hold the key to probing the explosion mechanisms and the progenitor scenarios of Type Ia supernovae (SNe Ia). We present multi-epoch spectropolarimetry and imaging polarimetry of SN 2019ein, an SNe Ia showing high expansion velocities at early phases. The spectropolarimetry sequence spans from ~ -11 to $+10$ d relative to peak brightness in the B band. We find that the level of the continuum polarization of SN 2019ein, after subtracting estimated interstellar polarization, is in the range 0.0–0.3 per cent, typical for SNe Ia. The polarization position angle remains roughly constant before and after the SN light-curve peak, implying that the inner regions share the same axisymmetry as the outer layers. We observe high polarization (~ 1 per cent) across both the Si II $\lambda 6355$ and Ca II near-infrared triplet features. These two lines also display complex polarization modulations. The spectropolarimetric properties of SN 2019ein rule out a significant departure from spherical symmetry of the ejecta for up to a month after the explosion. These observations disfavour merger-induced and double-detonation models for SN 2019ein. The imaging polarimetry shows weak evidence for a modest increase in polarization after ~ 20 d since the B -band maximum. If this rise is real and is observed in other SNe Ia at similar phases, we may have seen, for the first time, an aspherical interior similar to what has been previously observed for SNe IIP. Future polarization observations of SNe Ia extending to post-peak epochs will help to examine the inner structure of the explosion.

Key words: polarization – techniques: polarimetric – supernovae: individual: SN 2019ein – white dwarfs.

1 INTRODUCTION

During the last half century, Type Ia supernovae (SNe Ia; for reviews of SNe, see Filippenko 1997; Gal-Yam 2017) have answered (and posed) a myriad of interesting questions in astrophysics. These range from nucleosynthesis, chemical enrichment (Renzini 1999), and heating of the interstellar medium (ISM; Ciotti et al. 1991) to the discovery of the accelerating expansion of the Universe (Riess et al. 1998; Perlmutter et al. 1999), and more recently the so-called

Hubble tension (as summarized by Riess 2020). Yet, the nature of the progenitor systems of SNe Ia is still unclear. It has been generally established that the rise of SNe Ia is powered by the thermonuclear runaway of white dwarfs (WDs; see Hoyle & Fowler 1960; Howell 2011; Hillebrandt et al. 2013; Maoz, Mannucci & Nelemans 2014; Höflich 2017; Soker 2019 for recent reviews). However, the exact mechanism by which a WD's explosion is triggered and propagates through the progenitor still remains poorly understood (Arnett 1969; Nomoto, Sugimoto & Neo 1976; Khokhlov 1991; Niemeyer, Hillebrandt & Woosley 1996; Reinecke, Hillebrandt & Niemeyer 2002; Plewa, Calder & Lamb 2004; Röpke 2007; Pakmor et al. 2011; Seitenzahl et al. 2013).

Multiple channels of progenitors have been theorized, including the double-degenerate scenario in which two WDs merge (Iben & Tutukov 1984; Webbink 1984), the single-degenerate scenario in

* E-mail: kcpatra@berkeley.edu (KCP); yi.yang@berkeley.edu (YY)

† Nagaraj-Noll Graduate Fellow.

‡ Bengier-Winslow-Robertson Fellow.

§ Miller Senior Fellow.

which a WD accretes matter from a non-degenerate companion until the Chandrasekhar mass ($M_{\text{Ch}} \approx 1.4 M_{\odot}$) is approached (Whelan & Iben 1973), and tidal disruption of a WD by a compact companion and subsequent detonation of the WD (Luminet & Pichon 1989; Rosswog, Ramirez-Ruiz & Hix 2009).

Among these progenitor scenarios, a range of explosion mechanisms might be possible: delayed detonation, in which an initial deflagration front transitions to a detonation in a WD (Khokhlov 1991); double detonation, where a thin He layer on the WD detonates first, starting a detonation front in the WD (see e.g. Taam 1980; Fink et al. 2010; Shen et al. 2010); and compressional heating of WDs triggered by the dynamic merger of two C–O WDs (Hayden et al. 2010; Pakmor et al. 2010) or head-on collisions of WDs (Kushnir et al. 2013).

The shape of the ejecta and their circumstellar configuration are spatially unresolvable for extragalactic SN explosions, even with the best of ground-based interferometers.¹ Conventional photometry and spectroscopy provide a way to probe the kinematics and chemical structures of SN ejecta and their interaction with any pre-explosion circumstellar matter (CSM; see e.g. Nugent et al. 2011; Gal-Yam et al. 2014). However, these observations only offer crude clues on the structures of the ejecta and the interaction between the ejecta and any existing CSM. Such information is projected and smeared into the single dimension of radial velocity. Fortunately, spectropolarimetry, which measures polarization as a function of wavelength, provides a unique approach to the study of the SN explosion geometry. Any asphericity of the SN ejecta and the distribution of various elements formed in the ejecta are traced by the level of the continuum and the profiles of associated spectral lines in the polarization spectra, respectively. Additionally, the footprint of the interaction between the SN ejecta and any companion and CSM is encoded in the polarization spectra since such processes may create non-spherically symmetric emission and/or scattering photon sources.

In SN atmospheres, photons are scattered by free electrons (Thomson scattering). The polarization state of the emitted photons is determined when they escape the last-scattering surface, known as the photosphere. A photon that undergoes Thomson scattering will be polarized perpendicularly to the plane of scattering, which is defined as the plane containing the incident and scattered rays. For a spatially unresolved source, the observed polarization is an integration of the photons' electric vectors (E-vectors) projected in the plane of the sky. If the projected photosphere is circularly symmetric, a complete cancellation of the E-vectors results in zero net polarization. However, if the projected photosphere deviates from circular symmetry, incomplete cancellation of the E-vectors would lead to non-zero polarization across the spectrum. Additionally, any clumps of high-opacity absorbing material present above the photosphere along the observer's line of sight may block parts of the underlying photosphere. Therefore, an incomplete cancellation of the E-vectors will occur across the corresponding spectral lines, further producing non-zero polarization at the extinguished wavelengths.

SN 2019ein [α (J2000) = 13:53:29.11, δ (J2000) = +40:16:31.33] was discovered on 2019 May 1.47 (UT dates are used throughout this paper; Tonry et al. 2019) on the outskirts of the nearby galaxy NGC 5353. The host of SN 2019ein is a lenticular galaxy (Hubble type S0). A redshift of $z = 0.00775$ taken from the NASA/IPAC (Infrared Pro-

cessing and Analysis Center) Extragalactic Database² was adopted in this study. The spectrum obtained by the Las Cumbres Observatory Global SN project on 2021 May 2.3 (about 2 weeks before the *B*-band light-curve peak) shows a very high expansion velocity of $\sim 24\,000 \text{ km s}^{-1}$ as inferred from the absorption minimum of the Si II $\lambda 6355$ line. The Ca II near-infrared triplet (hereafter Ca II NIR3) and the O I lines are blended, creating a broad absorption profile. Curiously, the entire spectrum was slightly blueshifted with respect to the host-galaxy redshift, with the emission peaks of Si II, Ca II, and S II exhibiting velocities of $\sim 10\,000 \text{ km s}^{-1}$ towards the observer. Pellegrino et al. (2020) suggested that the apparent blueshift may be caused by an asymmetric explosion resulting in enhanced abundance of material at high velocities or due to optical-depth effects in the photosphere, in which most of the flux comes from material moving along the SN–Earth line of sight. The rise time of SN 2019ein was short ($15.37 \pm 0.55 \text{ d}$), after which the SN faded rapidly with a 15 d post-peak *B*-band magnitude decline (Phillips 1993) $\Delta m_{15(B)} = 1.36 \pm 0.02 \text{ mag}$ (Kawabata et al. 2020).

These features put SN 2019ein in the rare company of high-velocity SNe Ia like SNe 2004dt and 2006X, for which spectropolarimetric data have been obtained (Wang et al. 2006; Patat et al. 2009). In this work, we present five epochs of spectropolarimetry of SN 2019ein. We describe our observations in Section 2 and present our results along with the analysis in Section 3. We discuss the interpretations of the data in Section 4 and provide a concluding summary in Section 5.

2 OBSERVATIONS AND DATA REDUCTION

2.1 Kast spectropolarimetry

Spectropolarimetry of SN 2019ein was obtained using the polarimetry mode of the Kast double spectrograph on the Shane 3 m telescope at Lick Observatory (Miller, Robinson & Goodrich 1988). In the polarimetry mode, the light beam incident on the spectrograph is passed through a rotatable, half-wave plate and then through a Wollaston prism. The prism splits the incident light into two perpendicularly polarized components, named the ordinary and the extraordinary beams, which appear on the detector as two parallel traces. Only the red channel of Kast was used for spectropolarimetry. A GG455 order-sorting filter was in place, blocking all first-order light below $\sim 4600 \text{ \AA}$ and all second-order light below $\sim 9800 \text{ \AA}$. The usable wavelength range of the set-up was $4600\text{--}9000 \text{ \AA}$. Observations were made with the $300 \text{ lines mm}^{-1}$ grating and the 3 arcsec wide slit, resulting in a resolution of $\Delta\lambda \approx 18 \text{ \AA}$ ($\sim 800 \text{ km s}^{-1}$) at the central wavelength $\sim 6800 \text{ \AA}$.

Flat-field and arc-lamp exposures were obtained at the beginning of the observation night. The flat-field spectra were produced by the reflection of the light from an incandescent lamp off the inner surface of the dome.

SN 2019ein and polarization standard stars were observed each night. Four exposures were carried out at retarder-plate angles of 0° , 45° , 22.5° , and 67.5° . Multiple sets of polarimetry exposures were obtained for SN 2019ein to achieve higher signal-to-noise (S/N) ratios. Since all observations were carried out at small airmasses (≤ 1.25 ; see Table 1), we aligned the slit to the position angle (PA) of 180° (north–south direction). Because Kast does not have an atmospheric dispersion compensator to atone for the differential loss of blue light (Filippenko 1982), the following sanity check was

¹The minimum resolution required to study a nearby SN, for instance 3 Mpc away with a photosphere 100 au wide, would be $\sim 10 \text{ \mu as}$. For comparison, the Event Horizon Telescope can achieve a resolution of $\sim 60 \text{ \mu as}$.

²<https://ned.ipac.caltech.edu>

Table 1. Journal of spectropolarimetric observations.

UT date (MM-DD-YYYY)	MJD	Phase ^a (d)	Airmass range	Avg. seeing (arcsec)	Wavelength range (Å)	Exposure time ^b (s)
05-05-2019	58607.3	−10.9	1.04–1.03	1.30	4570–9000	4 × 1080
05-06-2019	58608.3	−9.9	1.04–1.13	1.21	4570–9000	4 × 1080
05-12-2019	58614.3	−3.9	1.01–1.14	1.63	4570–9000	4 × 1080
05-13-2019	58615.4	−2.9	1.03–1.13	1.29	4570–9000	4 × 1080
05-26-2019	58628.4	+10.1	1.09–1.25	1.86	4570–9000	4 × 1080

^aRelative to *B*-band peak brightness at MJD 58618.2 (Kawabata et al. 2020).^bNumber of waveplate positions × exposure time at each position.

carried out. For each night, we compared the Stokes parameters measured for different sets of spectropolarimetry with the values derived for the set obtained at the smallest airmass, typically 1.05. The Stokes parameters for each set were consistent within the associated uncertainties, suggesting a negligible effect on the polarization measurement from the loss of blue light.

Our nightly observations of the unpolarized standard star HD 110897 confirmed the low instrumental polarization of the Kast spectrograph (Section 3.2). The polarization test to determine the instrumental response to 100 per cent linearly polarized light was done by observing the same unpolarized standard star through a polarizing filter. Each night we also conducted spectropolarimetry of two polarization standard stars chosen among HD 154445, HD 161056, and HD 155528 to determine the accuracy of polarimetric measurements (see Section 3.2 for more details). A polarization ‘probe star’ was also observed to estimate the Galactic interstellar polarization (ISP) (see Section 3.3).

Extraction of the ordinary and extraordinary beams was carried out following standard techniques for CCD processing and spectrum extraction within IRAF.³ The images were bias subtracted using an overscan region. Cosmic ray hits on the detector were removed with L.A.Cosmic (van Dokkum 2001). For each night, flat-field images were combined and normalized by a low-order spline function to fit the continuum before applying to the science images. Then, we used the prescription of Horne (1986) to optimally extract each spectrum independently from the science images with apertures typically set to width at ∼1–2 per cent of the maximum of the spectrum profile. The background apertures were usually 5–10 pixels wide and placed two to three times the value of the full width at half-maximum (FWHM) intensity away from the profile centre.

Wavelength calibration was conducted separately for the ordinary and extraordinary beams in each individual exposure (all four retarder-plate angles) using lamp exposures. Small wavelength adjustments determined from the night-sky lines in the object frames were also applied to fine-tune the wavelength calibration. A typical root-mean-square (RMS) accuracy of ∼0.2 Å was achieved. Flux calibration of the ordinary and extraordinary beams of the SN was applied using the corresponding beam of a flux standard star observed at a similar airmass. We fit splines to the continuum of the flux-standard spectrum to generate a ‘sensitivity function’ that maps CCD counts to the flux at each wavelength. This mapping function was then applied to the SN spectra. Correction for telluric absorption was carried out by interpolating over the atmospheric absorption regions of the flux-standard spectrum.

³IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation (NSF).

2.2 RINGO3 imaging polarimetry

In this work, we adopted the imaging polarimetry of SN 2019ein from Maund et al. (2021). Details of the RINGO3 observations and data reduction are provided by Maund et al. (2021). In brief, they obtained the data using the Liverpool Telescope located on the Canary Island of La Palma, with the RINGO3 polarimeter (Arnold et al. 2012). Observations were carried out by three cameras optimized to integrate over the following wavelength ranges: *b*, 3500–6400 Å; *g*, 6500–7600 Å; and *r*, 7700–10000 Å. ISP was subtracted from the Stokes parameters measured by Maund et al. (2021) (see Section 3.7 for details). The Stokes parameters, polarization, and its PA are shown in Table 2.

3 ANALYSIS AND RESULTS

3.1 Calculating the stokes *q* and *u*

We express the normalized Stokes parameters as $q = Q/I$ and $u = U/I$, where Q and U are the differences in flux with the E-vector oscillating in two perpendicular directions and I is the total flux. U measures polarization along angles that are rotated by 45° with respect to those measured by Q .

We calculate q and u from two sets of spectra obtained with the waveplate at [0°, 45°] and [22.5°, 67.5°], respectively. From the ordinary (o) and the extraordinary (e) flux beams (f), q can be expressed as

$$q_o = \frac{f_{o,0} - f_{o,45}}{f_{o,0} + f_{o,45}} \quad \text{and} \quad q_e = \frac{f_{e,0} - f_{e,45}}{f_{e,0} + f_{e,45}}, \quad (1)$$

respectively, which are then averaged. Similarly, we calculate u using the exposures at the other set of waveplate positions. The observed polarization is then given by

$$p_{\text{obs}} = \sqrt{q^2 + u^2} \quad (2)$$

and the polarization PA is

$$\text{PA}_{\text{obs}} = \frac{1}{2} \arctan \left(\frac{u}{q} \right). \quad (3)$$

The polarization defined this way is positive definite and biased towards higher polarization, especially in the low-S/N regime. The final derived polarization was achieved after a debiasing procedure following Wang, Wheeler & Höflich (1997):

$$p = \left(p_{\text{obs}} - \frac{\sigma_p^2}{p_{\text{obs}}} \right) \times h(p_{\text{obs}} - \sigma_p) \quad \text{and} \quad \text{PA} = \text{PA}_{\text{obs}}, \quad (4)$$

where σ_p denotes the 1σ uncertainty in p and h is the Heaviside step function. The flux spectrum is calculated by averaging all the spectra of o-rays and e-rays used in deriving q and u . Figs 1–5 show the measured Stokes q , u , p , PA, and total flux at each epoch.

Table 2. Summary of polarimetry results.

MJD	Phase ^a (d)	Instrument	q (per cent)	u (per cent)	p (per cent)	PA (°)	q (per cent)	u (per cent)	p (per cent)	PA (°)
				(Observed)				(ISP corrected)		
58607.3	−10.9	Kast	−0.23 (07)	0.05 (07)	0.21 (07)	83 (9)	0.01 (07)	−0.14 (07)	0.10 (07)	138 (15)
58608.3	−9.9	Kast	−0.18 (05)	0.09 (05)	0.19 (05)	77 (7)	0.06 (05)	−0.10 (05)	0.10 (05)	151 (12)
58612.9	−5.3	RINGO3: b	−0.50 (16)	0.43 (15)	0.64 (15)	70 (7)	−0.13 (25)	−0.04 (24)	<0.01	99 (>360)
58614.3	−3.9	Kast	−0.22 (02)	0.22 (02)	0.31 (02)	68 (2)	0.02 (02)	0.03 (02)	0.02 (02)	210 (18)
58615.4	−2.9	Kast	−0.23 (04)	0.23 (04)	0.32 (04)	68 (3)	0.01 (04)	0.03 (04)	0.00 (04)	215 (30)
58620.0	1.8	RINGO3: b	−0.33 (15)	0.64 (16)	0.70 (15)	59 (6)	0.04 (24)	0.17 (25)	<0.01	219 (>360)
58622.9	4.7	RINGO3: b	−0.28 (15)	0.34 (13)	0.42 (12)	65 (9)	0.09 (24)	−0.13 (22)	<0.01	152 (>360)
58628.4	10.1	Kast	−0.07 (06)	−0.05 (07)	0.03 (07)	108 (24)	0.17 (06)	−0.24 (07)	0.28 (07)	153 (7)
58630.9	12.7	RINGO3: b	−0.15 (24)	1.14 (26)	1.12 (25)	49 (6)	0.22 (33)	0.67 (35)	0.54 (34)	215 (17)
58639.0	20.8	RINGO3: b	−0.47 (23)	−0.91 (23)	0.99 (22)	122 (7)	−0.10 (32)	−1.38 (32)	1.31 (32)	133 (6)

^aRelative to B -band peak brightness at MJD 58618.2 (Kawabata et al. 2020).




Figure 1. Spectropolarimetry of SN 2019ein at day −10.9 relative to the B -band peak brightness at MJD 58618.2 (Kawabata et al. 2020). The cyan vertical bands represent the regions of telluric correction. The panels below the total-flux spectrum represent the polarimetry before (red) and after (black) the ISP correction. The grey-shaded area indicates the associated 1σ uncertainty. The PA panel shows only the polarization position angle after ISP correction. With the exception of the flux spectrum, we use 50 Å binning for the purpose of presentation.

3.2 Polarimetric calibration

The nightly measured Stokes q and u of the unpolarized standard star HD 110897 are consistent with a level of < 0.05 per cent, indicating a low instrumental polarization and a high stability of the Kast spectropolarimeter over time. In the polarizance test with the same standard star, we determined that the polarimetric response of the instrument is larger than 99.5 per cent across the entire wavelength range (4600–9000 Å) and therefore does not necessitate further correction. The polarization PA of SN 2019ein was corrected as follows:

$$q_{\text{corr}} = p_{\text{obs}} q_{\text{obs}} \cos 2(\text{PA}_{\text{obs}} - \text{PA}_i),$$

$$u_{\text{corr}} = p_{\text{obs}} u_{\text{obs}} \cos 2(\text{PA}_{\text{obs}} - \text{PA}_i),$$

where PA_i is the position angle of the instrumental polarization determined from the polarizance test.

The polarization and PA measurements of the two high-polarization standards observed on each night were respectively found to be within 0.1 per cent and 3° of the references (Schmidt, Elston & Lupie 1992; Wolff, Nordsieck & Nook 1996).

3.3 Interstellar polarization

Light passing through interstellar dust clouds is polarized through dichroic extinction by non-spherical paramagnetic dust grains present along the line of sight. The contribution to polarization by dust, namely the ISP, must be removed to determine the intrinsic polarization of the source. Although several approaches have been commonly used to estimate the ISP along the SN–Earth line of sight (see e.g. Stevance et al. 2019; Yang et al. 2020), the exact level of the ISP contribution to the observed polarization of SN 2019ein is generally uncertain.

Serkowski, Mathewson & Ford (1975) showed that the Galactic ISP can be constrained to $p_{\text{ISP}} < 9 \times E(B - V)$ per cent. The Milky Way colour excess $E(B - V)_{\text{MW}}$ along the line of sight of SN 2019ein is 0.011 mag (Schlafly & Finkbeiner 2011), constraining ISP_{MW} to < 0.1 per cent. This upper limit is commensurate with the measured polarization of an ISP ‘probe star’⁴ – an intrinsically unpolarized star within 1° of SN 2019ein that probes at least 150 pc scale height of the Galactic ISM. The Stokes q and u measured for the probe star were found to be < 0.05 per cent in the continuum wavelength range of SN 2019ein, indicating low contribution from Galactic reddening.

An upper limit of the ISP from the host galaxy of SN 2019ein (NGC 5353) can be estimated based on the host reddening $E(B - V)_{\text{host}} = 0.09 \pm 0.02$ mag (Kawabata et al. 2020). Accounting for both the Galactic and SN 2019ein-host dust, we place an upper limit of $p_{\text{ISP}} < 0.9$ per cent along the SN–Earth line of sight. Such a value of ISP is higher than the continuum polarization level seen in SN 2019ein; thus, interstellar dust could potentially account for all of the continuum polarization of SN 2019ein. The caveat, however, is that Serkowski’s law may not be applicable to all galaxies because different dust properties could lead to different efficiencies for ISP (Leonard et al. 2002). Therefore, we employ a more direct approach by following the method used by Yang et al. (2020) to estimate the ISP Stokes parameters q_{ISP} and u_{ISP} .

We consider the wavelength region 4800–5600 Å in the spectrum when the SN is near its peak brightness. This region is expected

⁴We observed the star *Gaia* DR2 1497177392672672128.




Figure 2. Similar to Fig. 1 but for day -9.9 . We use 40 \AA binning for the purpose of presentation.

to be intrinsically depolarized owing to multiple overlapping Fe absorption features, which create a ‘line blanketing’ effect whose opacity dominates over electron scattering (Howell et al. 2001; Maund et al. 2013). We set the level of Stokes q and u to 0 within this wavelength range on day -2.9 (see Fig. 4), giving us an estimate of $q_{\text{ISP}} \approx -0.24$ per cent and $u_{\text{ISP}} \approx 0.19$ per cent. We note that these estimates are consistent with the upper limit derived earlier using Serkowski’s rule. These ISP values were subtracted from the observed q and u on each night and the polarization and the PA were subsequently recalculated (see Table 2). We note that owing to the relatively low level of expected ISP, only a wavelength-independent ISP estimation is presented. We will discuss the ISP-corrected continuum and line polarization of SN 2019ein in Sections 3.4 and 3.5, respectively.

3.4 Continuum polarization

Aspherical distribution of electrons, for instance an ellipsoidal photosphere, will cause imperfect cancellation of polarization E-vectors, leading to a non-zero continuum polarization (Höflich 1991, 1995; Bulla, Sim & Kromer 2015; Stevanec et al. 2019). SNe Ia typically show low continuum polarization ($\lesssim 0.3$ per cent; see e.g. Wang & Wheeler 2008; Yang et al. 2020), indicating that SNe Ia tend to be remarkably close to being spherical.

The continuum polarization of SN 2019ein and the associated uncertainty were estimated by binning the Stokes parameters over a wavelength range of $6400\text{--}7150 \text{ \AA}$ following a procedure similar to that described by Yang et al. (2020). The selected spectral region is known to be free from strong absorption features (Patat et al. 2009). The uncertainty was correspondingly binned.

The continuum polarization on days -10.9 and -9.9 is low with values of $p_{\text{cont}, -10.9 \text{ d}} = 0.10 \pm 0.07$ per cent and $p_{\text{cont}, -9.9 \text{ d}} = 0.10 \pm 0.05$ per cent, respectively (Figs 1 and 2). The polarization is consistent with 0 as the SN approaches maximum brightness around day -3.9 and day -2.9 (Figs 3 and 4). After peak brightness, the polarization increases slightly, reaching $p_{\text{cont}, +10.1 \text{ d}} = 0.28 \pm$




Figure 3. Similar to Fig. 1 but for day -3.9 . We use 25 \AA binning for the purpose of presentation.




Figure 4. Similar to Fig. 1 but for day -2.9 . We use 25 \AA binning for the purpose of presentation. The wavelength region $4800\text{--}5600 \text{ \AA}$ was used to estimate the ISP as described in Section 3.3.

0.07 per cent (Fig. 5). We note that these values are consistent with infrared spectropolarimetry of SN 2019ein, which found a 3σ upper limit on polarization of 1.2 per cent around the SN peak brightness (Tinyanont et al. 2021).

The measured continuum polarization PA, which represents the position of global axisymmetry of the ejecta, remains fairly consistent before and after peak brightness, hovering around 150° . Even though the PA on days -3.9 and -2.9 is apparently larger ($\sim 210^\circ$), we note




Figure 5. Similar to Fig. 1 but for day +10.1. We use 30 Å binning for the purpose of presentation.

that the polarization is so close to zero that the PA is essentially undetermined on those days.

3.5 Line polarization

Polarization signal at specific spectral lines arises due to the presence of clumps of material above the photosphere. The absorbing material partially obscures the underlying Thomson-scattering photosphere, resulting in an excess of polarization superimposed on the continuum polarization over the range of the absorption wavelengths. Subsequently, we present the polarization of three absorption features as follows.

(i) Si II $\lambda 6355$: We observe weak line polarization (0.5 ± 0.4 per cent) on day -10.9 , whereas no significant line polarization was detected on day -9.9 . On days -3.9 and -2.9 , we see strong line polarization, which reaches its peak value of ~ 1 per cent at an expansion velocity of $\sim 17\,000\text{ km s}^{-1}$. The strong line polarization persists into day 10.1, at an expansion velocity of $\sim 13\,600\text{ km s}^{-1}$.

(ii) O I $\lambda 7774$: No significant line polarization was seen at any phase.

(iii) Ca II NIR3: Similarly to Si II, no significant line polarization was observed on days -10.9 and -9.9 . In contrast, on days -3.9 and -2.9 , we see strong line polarization, reaching its peak value of ~ 0.8 per cent at an expansion velocity of $\sim 18\,000\text{ km s}^{-1}$. The strong line polarization continues into day 10.1, reaching up to ~ 1.5 per cent at $17\,000\text{ km s}^{-1}$.

Starting on day -3.9 and thereafter, both Si II and Ca II show a complex structure in the polarization spectra, likely associated with high-velocity (HV) and normal-velocity (NV) components. For example, at day +10.1, the polarization across Ca II NIR3 reached two local maxima at $-17\,000$ and -5900 km s^{-1} . The emergence of the line polarization of both the HV and NV components over time may result from an increase of Si II and Ca II opacity from larger to smaller radii.

3.6 The q - u plane and the dominant axis

Plotting the Stokes parameters in the q - u plane allows us to examine the axisymmetry of the continuum and various spectral features. If the SN ejecta are smooth and axisymmetric, the data points should fall along a straight line called the 'dominant axis' (Wang et al. 2001, 2003). Deviations from the dominant axis in the perpendicular direction represent departures from axisymmetry and clumpiness of the ejecta. The dominant axis is determined by

$$u = \alpha q + \beta, \quad (5)$$

where α and β are the fitted parameters from an error-weighted orthogonal distance regression. In Fig. 6, we present the polarization in the q - u plane in the continuum as well as for Si II $\lambda 6355$ and Ca II NIR3. We omitted the Si II and Ca II lines when plotting the continuum q and u in the wavelength range 4700 – 8750 Å . The fitted parameters α and β that characterize the dominant axes are given in Table 3.

As suggested by the values of $\chi^2/\text{degrees of freedom (DoF)}$ in Fig. 6, the departure from the dominant axis fitted across the line profile indicates a significant clumpiness in the Si II-rich ejecta on days -10.9 and -9.9 . Considering the relatively large values of χ^2/DoF , it is ambiguous whether a dominant axis is present at early times. The absence of a clear dominant axis together with the measured low polarization suggests that any Si II-rich clumps are fairly uniformly distributed in the ejecta at early times, when the photosphere only intersects with the outermost part of the ejecta. Axial symmetry is evident on days -3.9 and -2.9 for Si II when the SN is near its peak brightness. However, the axial symmetry becomes weak and clumpiness increases again around day +10. Ca II shows an overall higher degree of clumpiness compared with Si II. Ca II also exhibits weak axial symmetry at early times. However, as time progresses, Ca II settles on to a more prominent symmetry axis. Unlike Si II, Ca II continues to exhibit high axial symmetry at day +10. The symmetry axes of both Si II and Ca II are roughly aligned with each other and remain fairly constant starting on day -10 and thereafter.

As seen in the left-hand panels of Fig. 6, no clear dominant axis can be identified in the continuum at any epoch. The data points form a cloud centred near the origin. This strengthens the case that even though the overall ejecta geometry is spherical, the polarization in Si II and Ca II is due to clumps of explosively synthesized material.

3.7 Polarization time series

We build a temporal series of polarization measurements of SN 2019ein by combining the Kast spectropolarimetry and the RINGO3 imaging polarimetry. In order to compare the polarization measured by the two instruments, we binned the Kast spectropolarimetry over the RINGO3 b , g , and r filter passbands. This process estimates the equivalent imaging polarimetry data points in RINGO3 filters. In this way, we built the polarimetric data set with a time baseline from -11 to 21 d relative to the B -band light-curve peak of SN 2019ein. The broad-band polarization from Kast was derived by integrating over the wavelength of the filter-transmission-weighted polarized flux.

The combined polarization time series is presented in Fig. 7. The top panel displays two light curves obtained in the Landolt I and B bands with the Katzman Automatic Imaging Telescope (KAIT; Filippenko et al. 2001) at Lick Observatory. The middle panel shows polarization over time. The red, blue, and green circles represent the synthesized Kast polarization in wavelength ranges that roughly match the three channels of RINGO3. The black squares present the





Figure 6. Polarization in the Stokes q - u plane. The Si II $\lambda 6355$ and Ca II NIR3 lines were omitted from the left-hand panels labelled 'continuum'. The dashed lines in the 'Si II $\lambda 6355$ ' and 'Ca II NIR3' panels represent the dominant axes for the labelled absorption features.

Table 3. Fitted parameters for the dominant axes.

Phase (d)	α_{cont}	β_{cont}	$\alpha_{\text{Si II}}$	$\beta_{\text{Si II}}$	Velocity range (-10^3 km s^{-1})	$\alpha_{\text{Ca II NIR3}}$	$\beta_{\text{Ca II NIR3}}$	Velocity range (-10^3 km s^{-1})
-10.9	0.81 (21)	-0.15 (04)	-2.23 (1.35)	-0.12 (23)	28.6–9.7	-0.74 (24)	-0.11 (07)	34.8–6.7
-9.9	4.68 (2.62)	-0.47 (24)	0.99 (49)	-0.10 (09)	28.5–7.3	0.87 (28)	-0.08 (06)	34.8–6.7
-3.9	0.33 (09)	-0.04 (01)	1.95 (27)	-0.06 (05)	23.9–5.0	2.24 (52)	0.00 (08)	26.1–3.2
-2.9	-0.94 (14)	0.03 (02)	1.37 (24)	-0.09 (06)	23.8–3.9	1.33 (23)	-0.01 (07)	26.1–3.2
+10.1	0.02 (13)	-0.20 (04)	0.90 (31)	-0.23 (08)	21.5–2.6	0.62 (11)	-0.28 (06)	22.5–0.3

**Figure 7.** Polarization versus time. The polarization and PA have been corrected for the ISP. The black squares represent the continuum polarization and PA from the Kast spectropolarimeter. The red, blue, and green circles show the polarization measured by Kast in different wavelength regions: blue (4580–6400 Å), green (6500–7600 Å), and red (7700–9000 Å). The blue inverted triangles show the polarization and PA measured by RINGO3 in the blue channel with wavelength range 3500–6400 Å. The light curves in the top panel were obtained by KAIT at Lick Observatory.

Kast continuum polarization in the wavelength range 6400–7150 Å. We note that the *b* and *r* bins include the polarization of the Si II λ 6355 and Ca II NIR3 features, respectively. The blue inverted triangles show the polarization measured by the blue channel of RINGO3. The bottom panel provides the measured PA in the continuum region (6400–7150 Å) by Kast and by the blue channel of RINGO3. The polarimetry presented here has been ISP corrected. Since we do not know the exact magnitude of any systematic bias (e.g. instrumental polarization) in RINGO3 measurements, we employed a different strategy to account for ISP and any systematic bias: We calculated

the mean *q* and *u* from the three epochs within ~ 5 d of the *B*-band peak brightness (days -5.3, +1.8, and +4.7; see Table 2), which gives $q_{\text{ISP+sys}} \approx -0.37$ per cent and $u_{\text{ISP+sys}} \approx 0.47$ per cent. We then subtracted the averaged *q* and *u* from the observed Stokes parameters of all RINGO3 data under the assumption that SNe Ia exhibit effectively zero continuum polarization near peak brightness. This assumption is validated independently by the Kast spectropolarimetry of SN 2019ein on days -2.9 and -3.9. We also propagated the uncertainty of ISP subtraction into the final calculations of *p* and PA.

4 DISCUSSION

A non-zero continuum polarization may result from either an overall inhomogeneous electron density distribution or a non-spherical heating source. The latter case was seen in models of Bulla et al. (2016a), Bulla et al. (2016b) for SNe Ia, and has also been used to explain the observed increase in polarization during the plateau phase of SNe IIP, in which an aspherical ionization front of ^{56}Ni is typically present (Höflich et al. 1996).

Overall, the polarization properties of SN 2019ein before its peak luminosity are consistent with the typical behaviour of SNe Ia. For example, the continuum polarization is <0.2 per cent. Distinct line polarization, which is typically of the order of 1 per cent, can also be identified across some prominent spectral lines, including Si II, Fe II, and Ca II (Wang & Wheeler 2008). The asymmetric distribution of the intermediate-mass elements (IMEs; $9 \leq Z \leq 20$, including Si, Ca, S, and Mg) inferred from the line polarization is indicative of sufficient outward mixing. IMEs generated in the nuclear burning can also be produced at higher velocities compared with the SN ejecta. In thermonuclear SNe, below the production zone of the IMEs, the inner burning region is surrounded by C and O from the progenitor WD. The energy input is given by the radioactive decay chain of $^{56}\text{Ni} \rightarrow ^{56}\text{Co} \rightarrow ^{56}\text{Fe}$, which is initiated by the nucleosynthesis of ^{56}Ni as the main product of the silicon-burning process. An asymmetric ^{56}Ni distribution hence results in an asymmetric energy source.

The low line and continuum polarization at early times (day ~ -11) do not support the idea put forward by Pellegrino et al. (2020) that the blueshifted emission peaks in early-time spectra are due to an aspherical explosion enhancing abundance of material at high velocities. Furthermore, the lack of a clear dominant axis in the $q-u$ plane at early times indicates that even if clumps of high-velocity material are present, they must be fairly uniformly distributed in the outer ejecta. We cannot, however, rule out the possibility that the apparent blueshift is due to optical-depth effects arising from a steep density profile in the ejecta (Pellegrino et al. 2020).

As shown in the bottom panel of Fig. 7, we do not identify significant variation in PA at different epochs. We note that near peak brightness, the q and u intrinsic to SN 2019ein are very close to zero after correcting for the ISP, as discussed in Section 3.7. Such small values of Stokes parameters lead to effectively random values of the PA around the SN light-curve peak. Furthermore, we remark that the PA calculated before ISP removal also tends to be consistent from day -11 to day $+21$ (see Table 2). A relatively low level of continuum polarization together with a roughly constant direction of the dominant axis suggests a common axial symmetry from the outer electron-scattering zone to the inner region near the energy source.

We remark that RINGO3 uses two dichroics to separate the three wavelength channels and a depolarizing Lyot prism, resulting in an induced systematic uncertainty in polarization of up to ~ 0.5 per cent (Jermak 2017). As shown in the third and fourth panels of Fig. 7, after day $+5$, the continuum level of polarization estimated from RINGO3 observations is mostly from u , while q is consistent with zero. A moderate degree of asphericity is suggested by Kast spectropolarimetry at day $+10$. Unfortunately, we were not able to conduct Kast spectropolarimetry of SN 2019ein after day $+10$ owing to technical issues. Therefore, the ‘rise’ in polarization after maximum brightness is anchored by just one RINGO3 measurement from day $+21$. For this reason, out of an abundance of caution, we refrain from claiming that a definitive rise in polarization was observed in SN 2019ein post maximum brightness. However, if the post-peak rise of polarization is real and intrinsic to SN 2019ein, we cautiously provide our interpretations subsequently, hoping to invite more sophisticated theoretical investigations.

We suggest that the secondary maximum in the NIR luminosity could be the key for understanding the post-peak rise of the continuum polarization. The formation of the secondary maximum in the NIR can be understood as an opacity effect. To first order, the NIR luminosity is given by $L_{\text{NIR}}(t) \propto T(t)^4 \times R(t)^2$, where $T(t)$ and $R(t)$ represent the temperature and the radius of the photosphere as a function of time, respectively. In normal SNe Ia, although the photospheric radius increases with time because the opacity remains high for ~ 2 – 3 weeks after the SN explosion, it recedes gradually in mass coordinate (see e.g. fig. 3 of Höflich 2017) until the SN light curve approaches its secondary maximum in the NIR. This may explain the relatively little evolution of polarization seen in SN 2019ein until peak brightness.

After peak brightness, the recession of the SN photosphere is governed by the geometrical dilution of the homologously expanding envelope. However, $R(t)$ decreases rapidly owing to the significant drop in opacity when Fe-group elements begin recombining from ionization states III to II (Höflich, Mueller & Khokhlov 1993; Höflich, Wheeler & Thielemann 1998; Kasen, Thomas & Nugent 2006). Therefore, when the SN reaches its secondary NIR maximum ~ 40 – 50 d after the explosion, most of the energy input emerges above the photosphere. In the presence of an asymmetric energy source, the flux at the photosphere will be direction dependent. Consequently, an inhomogeneous photosphere could develop, leading to rise in polarization of the SN after maximum brightness. The interpretation of any rise in late-time polarization may be complicated by optical-depth effects that still remain poorly understood (Höflich 1991) in SNe photospheres. As such, detailed theoretical investigations into how polarization behaves over time for various explosion mechanisms are pressingly warranted.

On the other hand, if the rise in late-time polarization is shown to be not true and the inner energy source is found to be spherical, we also arrive at an interesting implication. In that case, models with burning starting on the surface might be better at explaining the observed polarimetry. We speculate that if the burning starts on the surface, the detonation will propagate through the central zone supersonically and preserve the spherical nature of the WD.

Spectropolarimetric observations of SNe Ia beyond 20 d after maximum light are very rare. The handful of SNe Ia for which such measurements exist display low continuum polarization. For example, SN 2012fr was polarized to ~ 0.2 per cent on day $+24$ (Maund et al. 2013), and SN 2001el and SN 2006X showed < 0.1 per cent polarization on days $+38$ and $+39$, respectively (Wang et al. 2003; Patat et al. 2009). These observations challenge the apparent late-time increase in polarization of SN 2019ein. However, owing to a small sample and the fact that these SNe display a diverse set of properties (expansion velocities, decline rates, etc.), the question remains whether any SNe Ia display late-time increases in polarization. More polarimetric observations of SNe Ia at similar phases are required to answer that question.

4.1 Implications for explosion scenarios

The time-invariant PA observed from early times to the phase just prior to the secondary maximum in the NIR light curves of SN 2019ein brings us to the implications for explosion scenarios and the characteristics of p . Based on current understanding, SNe Ia might be triggered through the following mechanisms.

(i) Deflagration: Compressional heating in a slow accretion near the WD centre triggers subsonic burning when the WD approaches the Chandrasekhar mass M_{Ch} (Nomoto, Thielemann & Yokoi

1984; Gamezo, Khokhlov & Oran 2004; Röpke 2007; Ma et al. 2014).

(ii) Delayed detonation: Explosion begins at the end of a deflagration phase near or at the centre of the WD. When a critical density of $\sim 10^7 \text{ g cm}^{-3}$ is reached, a transition to detonation occurs (Khokhlov 1991).

(iii) Colliding/inspiralling WDs: Heat released on dynamical time-scales triggers a detonation of a double-degenerate system (Iben & Tutukov 1984; Webbink 1984; Benz et al. 1990; Nugent et al. 1997; Pakmor et al. 2010; Kushnir et al. 2013; García-Berro & Lorén-Aguilar 2017).

(iv) Double/helium detonation: A sub- M_{Ch} C–O WD may explode by detonating a thin surface He layer, which triggers a detonation front in the WD (Woosley, Weaver & Taam 1980; Nomoto 1982a,b; Livne 1990; Woosley & Weaver 1994; Höflich & Khokhlov 1996; Kromer et al. 2010).

In the framework of off-centre delayed-detonation models, a slightly asymmetric excitation will lead to an off-centre distribution of iron-group elements. The axis of symmetry will be defined by the centroid of the density distribution and the point of off-centre delayed-detonation transition (Livne 1999; Höflich et al. 2006; Fesen et al. 2007). If the ^{56}Ni region is above the photosphere, the time-invariant PA observed in SN 2019ein would indicate a moderate asphericity of the central energy source. Otherwise, a change of PA may be seen as the SN approaches its secondary peak. The low continuum polarization observed in SN 2019ein challenges any model that predicts significant asymmetry of the photosphere.

Dynamical or head-on collisions of WDs are expected to show larger asymmetries (Benz et al. 1990; Pakmor et al. 2011; Katz et al. 2016; Sato et al. 2016; García-Berro et al. 2017; García-Berro & Lorén-Aguilar 2017). The dynamical models are not favoured since they predict high polarization levels at early phases (Pakmor et al. 2012; Bulla et al. 2016a) and larger asymmetries in the inner layers or off-centred energy sources, which are incompatible with our observations of SN 2019ein.

For sub- M_{Ch} explosions through a double/helium detonation, outer asymmetry is expected owing to the He-ignition process. Classical He-detonation models require a significant He surface mass of the order of $0.01\text{--}0.1 M_{\odot}$ (e.g. Nomoto 1982a,b; Woosley & Weaver 1994; Höflich & Khokhlov 1996; Bildsten et al. 2007; Shen & Bildsten 2009). Starting from the SN explosion, the photosphere recedes and will first cross the burning product of the outermost He layer. The O I $\lambda 7774$ feature is prominent at day -10.9 in the flux spectrum. However, we see no polarization signal at the corresponding wavelength, suggesting that oxygen was present in the outer layer and maintained a spherically symmetric distribution in the expanding envelope. This is in contrast with the ~ 0.4 per cent O I $\lambda 7774$ line polarization predicted by Bulla et al. (2016b) for the double/helium-detonation models. Therefore, we infer that the outermost layer is dominated by the spherical pre-explosion C and O from the WD, rather than a He shell, since the latter is likely to produce an abundance jump in O in the line-forming region (see e.g. Yang et al. 2020). In fact, SNe 2006X and 2004dt, both of which are high-velocity SNe Ia, also show no polarization across O I, putting strong constraints on the distribution of oxygen in the explosion ejecta. We note that the polarimetric properties of more modern models of sub- M_{Ch} double/helium detonation (e.g. Shen et al. 2018) are currently theoretically unexamined.

4.2 Comparison with a detonating failed deflagration model

Kasen & Plewa (2007) made theoretical predictions for spectropolarimetric observations of a M_{Ch} WD using the detonating failed deflagration (DFD) model. They studied one particular model in detail, named Y12, in which the WD’s ignition starts within a small spherical region, 50 km in diameter and offset 12.5 km from the centre. Here, we compare the observed spectropolarimetric properties of SN 2019ein with the predictions of Kasen & Plewa (2007).

According to Kasen & Plewa (2007), if an SN is observed from the deflagration side – the side where ignition began – high ejecta velocities are expected. Since SN 2019ein exhibited very high expansion velocities, we may be observing the explosion from the ignition side (viewing angles $\theta \approx 0^\circ$). From this orientation, the projected surface of the intrinsically ‘egg-shaped’ density structure in the observer’s direction would be fairly circular, leading to low continuum polarization. In fact, Kasen & Plewa (2007) argue that low continuum polarization is expected from all viewing angles at peak brightness. Therefore, continuum polarization is not informative for constraining the viewing angle of the SN.

The Y12 model predicts substantial line polarization (1–2 per cent) depending on the viewing angle (fig. 13 of Kasen & Plewa 2007). Indeed, we observe significant line polarization across both Si II $\lambda 6355$ and Ca II NIR3 features in SN 2019ein, which suggests a viewing angle of $\theta \approx 0^\circ$ or 90° . A viewing angle of $\sim 180^\circ$ (opposite the ignition side) is disfavoured because we observe high polarization across both Si II and Ca II, whereas in the Y12 model only Si II polarization is seen for angles $\sim 180^\circ$. Together with the high expansion velocity of SN 2019ein, $\theta \approx 0^\circ$ is favoured over other orientations, strengthening the case that we may be viewing the SN from the ignition side. According to Kasen & Plewa (2007), such events are rare and expected to constitute roughly 10 per cent of all SNe Ia. Spectropolarimetry of more SNe Ia is needed to test this prediction.

As described in Section 3.6, both Si II and Ca II display a higher degree of clumpiness at early epochs than near and after peak brightness. This is also expected in the DFD model, which can produce a clumpy outer layer of IMEs but maintains a relatively smoother IME distribution in the inner layers.

4.3 The Si II $\lambda 6355$ polarization compared with a larger sample

We notice that the continuum polarization of SN 2019ein on the Stokes q – u diagram can be fitted with straight lines (e.g. see the left-hand panels of Fig. 6). A dominant axis is present in Kast spectropolarimetry between days ~ -11 and $+10$. Except for the first epoch, the direction of the dominant axis appears to be unchanged, suggesting that different layers of the ejecta share a roughly fixed axial symmetry. These properties indicate that SN 2019ein belongs to the spectropolarimetric type D1 (Wang & Wheeler 2008), in which a dominant axis is identifiable but with significant scatter.

We compared the polarimetric properties of SN 2019ein with those of SNe 2004dt and 2006X, both of which display high expansion velocities at early phases. As inferred from the absorption minimum of the Si II $\lambda 6355$ line, SN 2004dt shows an expansion velocity of $\sim 17\,000 \text{ km s}^{-1}$ $\sim 6\text{--}8$ d before the optical maximum (Wang et al. 2006), and SN 2006X exhibits an expansion velocity of $20\,700 \text{ km s}^{-1}$ at day -11.3 (Wang et al. 2008). A linear correlation between the maximum polarization measured across Si II $\lambda 6355$,

$p_{\text{Si II}}^{\text{max}5}$, and the SN expansion velocity traced by the same line at day -5 , $v_{\text{Si II}@-5\text{d}}$, has been found by Cikota et al. (2019) based on the analysis of a sample of 35 SNe Ia. The velocity–polarization relation connects the kinematics with the ejecta asymmetry and indicates that a higher departure from spherical symmetry for Si is produced at higher velocities.

For comparison with the Si II velocity–polarization relation presented in fig. 13 of Cikota et al. (2019), we estimated $v_{\text{Si II}@-5\text{d}} = 15\,100 \pm 300 \text{ km s}^{-1}$ for SN 2019ein. The peak polarization of SN 2019ein across the Si II $\lambda 6355$ line, i.e. $p_{\text{Si II}}^{\text{max}}$ derived based on 100 and 50 Å binnings on day -4 , is 0.76 ± 0.10 per cent and 0.82 ± 0.16 per cent, respectively. These values place SN 2019ein slightly above the predicted Si II $\lambda 6355$ polarization. We remark that SN 2019ein is still broadly consistent with the Si II velocity–polarization relation, corroborating the trend that higher velocity SNe Ia tend to exhibit higher polarization. SN 2019ein shows significantly lower Si II $\lambda 6355$ polarization compared with SN 2004dt ($14\,870 \pm 140 \text{ km s}^{-1}$, 1.34 ± 0.14 per cent), which exhibits an exceptionally high peak polarization across the Si II line and was considered an outlier by Cikota et al. (2019). On the other hand, SN 2006X ($17\,040 \pm 90 \text{ km s}^{-1}$, 0.63 ± 0.05 per cent) is in good agreement with the Si II velocity–polarization relation.

Wang, Baade & Patat (2007) also derived a correlation between the maximum line polarization of Si II $\lambda 6355$ and $\Delta m_{15}(B)$. For the former parameter, the observations are often converted to the level at 5 d before the B -band maximum, i.e. $p_{\text{Si II}}^{\text{corr}-5}$. Owing to the sparsely sampled spectropolarimetry, we adopt the peak Si II $\lambda 6355$ polarization measured at day -4 for SN 2019ein. The B -band light-curve decline rate of SN 2019ein has been determined as $\Delta m_{15}(B) = 1.36 \pm 0.02 \text{ mag}$ (Kawabata et al. 2020) and $\Delta m_{15}(B) = 1.40 \pm 0.004 \text{ mag}$ (Pellegrino et al. 2020). We infer that SN 2019ein is consistent with the $\Delta m_{15}(B)$ – $p_{\text{Si II}}^{\text{corr}-5}$ relation as presented by Wang et al. (2007) and Cikota et al. (2019). This relation can be interpreted such that at a given epoch, higher line polarization is expected for less luminous SNe, which indicates a higher chemical non-uniformity. This can be understood as an indication that less material is burnt in fainter events, and such incomplete burning may not be sufficient to wipe out lumpy chemical configurations.

Therefore, we conclude that SN 2019ein is consistent with both the Si II velocity–polarization relationship and the light-curve decline rate–Si II polarization relation. Such behaviour may be explained with the off-centre delayed-detonation model (e.g. Höflich et al. 2006; Cikota et al. 2019).

5 CONCLUDING SUMMARY

We have presented spectropolarimetry of SN 2019ein, a high-velocity SN Ia in NGC 5353. Our observations range from day -10.9 to $+10.1$ from the B -band light-curve peak. We found that the continuum polarization in SN 2019ein is low, staying < 0.25 per cent until about a month after the explosion. This indicates that the photosphere is quite close to being spherical.

The blueshifted emission peaks observed in early-time spectra of SN 2019ein cannot be due to a highly asymmetric explosion, as evidenced by low continuum polarization at early times. However, our observations do not preclude the possibility that optical-depth effects in steep-density ejecta lead to the apparent blueshift of the emission peaks.

⁵The peak of the Si II $\lambda 6355$ polarization is measured between roughly days -11 and $+1$ (Cikota et al. 2019).

The RINGO3 imaging polarimetry shows an apparent increase in polarization (~ 1 per cent) around day $+21$. However, owing to significant systematic uncertainties found in previous RINGO3 measurements, we are cautious of the observed rise in polarization. If the post-peak increase in polarization is real and intrinsic to SN 2019ein, we note that it coincides with the beginning of the transition from Fe III to Fe II ionization states. The recombination decreases the opacity, providing us a deeper view into the SN ejecta. We speculate that the possible post-peak rise of the polarization, therefore, could indicate the presence of an aspherical central energy source.

The polarization PA does not change drastically over the observed epochs. We also observe high line polarization (~ 1 per cent) across the Si II $\lambda 6355$ and the Ca II NIR3 features around peak brightness of SN 2019ein. The polarization signatures of SN 2019ein are consistent with models predicting SNe Ia explosions that produce a modest amount of asphericity. To summarize,

- (i) a low amount of asphericity in the high-velocity layers is detected, as in previous observations of other SNe Ia.
- (ii) significant departures from global spherical symmetry can be ruled out throughout the ejecta. A common symmetry axis persists from the outer to the inner layers.
- (iii) after day $+21$, the possibility of a small amount of polarization caused by an asymmetric distribution of ^{56}Ni , which may arise from many different models of SN explosions (Höflich 1991; Leonard et al. 2005; Kasen, Röpke & Woosley 2009; Pakmor et al. 2010; Seitenzahl et al. 2013; Moll et al. 2014; Raskin et al. 2014; Bulla et al. 2015, 2016a,b; Höflich et al. 2017), cannot be eliminated. Spectropolarimetry of more SNe Ia at post-peak epochs is needed to confirm whether the polarization rises beyond day $+20$.

The spectropolarimetric observations of SN 2019ein strengthen existing evidence that the explosions of SNe Ia are largely spherical, especially when considering that SN 2019ein is an event with one of the highest expansion velocities ever observed.

Finally, we compared the results with the detonating failed deflagration model of Kasen & Plewa (2007) and found that the low continuum polarization but high line polarization is consistent with the model. A viewing angle of $\theta \approx 0^\circ$ is favoured, which means we may be viewing SN 2019ein from the ignition side.

We recommend high-quality spectropolarimetric observations of bright, nearby future SNe Ia to be carried out covering both their rising and falling phases. Such polarimetric tomography is essential for building a robust picture of how polarization signatures vary over time and the consequences of various explosion mechanisms and progenitor scenarios.

ACKNOWLEDGEMENTS

KCP is thankful to Sergiy Vasylyev and Matthew Chu for helpful discussions. AVF's group at U.C. Berkeley acknowledges generous support from the Miller Institute for Basic Research in Science, Sunil Nagaraj, Landon Noll, Gary and Cynthia Bengier, Clark and Sharon Winslow, Sanford Robertson, and many additional donors. The U.C. Santa Cruz team is supported in part by NASA grants NNG17PX03C, 80NSSC19K1386, and 80NSSC20K0953; National Science Foundation (NSF) grant AST-1815935; the Gordon & Betty Moore Foundation; the Heising-Simons Foundation; and by a fellowship from the David and Lucile Packard Foundation to RJF. The research of YY is supported through the Bengier-Winslow-Robertson Fellowship and the Benozio Prize Postdoctoral Fellowship. PH acknowledges the support by the NSF project 'Signatures of Type

Ia Supernovae, New Physics and Cosmology’, grant AST-1715133. The supernova research by LW is supported by the NSF award AST-1817099. JCW is supported by the NSF grant AST-1813825. The research of JM is supported through a Royal Society University Research Fellowship. MB acknowledges support from the Swedish Research Council (reg. no. 2020-03330).

A major upgrade of the Kast spectrograph on the Shane 3 m telescope at Lick Observatory, led by Brad Holden, was made possible through generous gifts from the Heising-Simons Foundation, William and Marina Kast, and the University of California Observatories. The Katzman Automatic Imaging Telescope (KAIT) and its ongoing operation were made possible by donations from Sun Microsystems, Inc., the Hewlett-Packard Company, AutoScope Corporation, Lick Observatory, the US NSF, the University of California, the Sylvia & Jim Katzman Foundation, and the Tabasco Foundation. Research at Lick Observatory is partially supported by a generous gift from Google. We appreciate the excellent assistance of the staff at Lick Observatory.

PYRAF, PYFITS, and STSCI-PYTHON are products of the Space Telescope Science Institute (STScI), which is operated by the Association of Universities for Research in Astronomy, Inc., (AURA) under NASA contract NAS5-26555. This research has made use of NASA’s Astrophysics Data System Bibliographic Services; the Set of Identifications, Measurements and Bibliography for Astronomical Data (SIMBAD); and the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA.

DATA AVAILABILITY

The raw data used in this work may be shared upon request to Kishore C. Patra (kcpatra@berkeley.edu).

REFERENCES

- Arnett W. D., 1969, *AP&SS*, 5, 180
- Arnold D. M., Steele I. A., Bates S. D., Mottram C. J., Smith R. J., 2012, in McLean I. S., Ramsay S. K., Takami H., eds, *SPIE Conf. Ser. Vol. 8446, Ground-based and Airborne Instrumentation for Astronomy IV*. SPIE, Bellingham, p. 84462J
- Benz W., Bowers R. L., Cameron A. G. W., Press W. H., 1990, *ApJ*, 348, 647
- Bildsten L., Shen K. J., Weinberg N. N., Nelemans G., 2007, *ApJ*, 662, L95
- Bulla M., Sim S. A., Kromer M., 2015, *MNRAS*, 450, 967
- Bulla M., Sim S. A., Pakmor R., Kromer M., Taubenberger S., Röpke F. K., Hillebrandt W., Seitenzahl I. R., 2016a, *MNRAS*, 455, 1060
- Bulla M. et al., 2016b, *MNRAS*, 462, 1039
- Cikota A. et al., 2019, *MNRAS*, 490, 578
- Ciotti L., D’Ercole A., Pellegrini S., Renzini A., 1991, *ApJ*, 376, 380
- Fesen R. A., Höflich P. A., Hamilton A. J. S., Hammell M. C., Gerardy C. L., Khokhlov A. M., Wheeler J. C., 2007, *ApJ*, 658, 396
- Filippenko A. V., 1982, *PASP*, 94, 715
- Filippenko A. V., 1997, *ARA&A*, 35, 309
- Filippenko A. V., Li W. D., Treffers R. R., Modjaz M., 2001, in Paczynski B., Chen W.-P., Lemme C., eds, *ASP Conf. Ser. Vol. 246, IAU Colloq. 183: Small Telescope Astronomy on Global Scales*. Astron. Soc. Pac., San Francisco, p. 121
- Fink M., Röpke F. K., Hillebrandt W., Seitenzahl I. R., Sim S. A., Kromer M., 2010, *A&A*, 514, A53
- Gal-Yam A., 2017, *Observational and Physical Classification of Supernovae*. Springer, New York, p. 195
- Gal-Yam A. et al., 2014, *Nature*, 509, 471
- Gamezo V. N., Khokhlov A. M., Oran E. S., 2004, *Phys. Rev. Lett.*, 92, 211102
- García-Berro E., Lorén-Aguilar P., 2017, in Alsabti A., Murdin P., eds, *Handbook of Supernovae*. Springer, Cham, p. 1237
- García-Berro E., Badenes C., Aznar-Siguán G., Lorén-Aguilar P., 2017, *MNRAS*, 468, 4815
- Hayden B. T. et al., 2010, *ApJ*, 722, 1691
- Hillebrandt W., Kromer M., Röpke F. K., Ruiter A. J., 2013, *Front. Phys.*, 8, 116
- Höflich P., 1991, *A&A*, 246, 481
- Höflich P., 1995, *ApJ*, 443, 89
- Höflich P., 2017, *Explosion Physics of Thermonuclear Supernovae and Their Signatures*. Handbook of Supernovae, Springer, Cham, p. 1151
- Höflich P., Khokhlov A., 1996, *ApJ*, 457, 500
- Höflich P., Mueller E., Khokhlov A., 1993, *A&A*, 268, 570
- Höflich P., Wheeler J. C., Hines D. C., Trammell S. R., 1996, *ApJ*, 459, 307
- Höflich P., Wheeler J. C., Thielemann F. K., 1998, *ApJ*, 495, 617
- Höflich P., Gerardy C. L., Marion H., Quimby R., 2006, *New Astron. Rev.*, 50, 470
- Höflich P. et al., 2017, *ApJ*, 846, 58
- Horne K., 1986, *PASP*, 98, 609
- Howell D. A., 2011, *Nat. Commun.*, 2, 350
- Howell D. A., Höflich P., Wang L., Wheeler J. C., 2001, *ApJ*, 556, 302
- Hoyle F., Fowler W. A., 1960, *ApJ*, 132, 565
- Iben Jr. I., Tutukov A. V., 1984, *ApJS*, 54, 335
- Jermak H., 2017, PhD thesis, Liverpool John Moores University
- Kasen D., Plewa T., 2007, *ApJ*, 662, 459
- Kasen D., Thomas R. C., Nugent P., 2006, *ApJ*, 651, 366
- Kasen D., Röpke F. K., Woosley S. E., 2009, *Nature*, 460, 869
- Katz M. P., Zingale M., Calder A. C., Swesty F. D., Almgren A. S., Zhang W., 2016, *ApJ*, 819, 94
- Kawabata M. et al., 2020, *ApJ*, 893, 143
- Khokhlov A. M., 1991, *A&A*, 245, 114
- Kromer M., Sim S. A., Fink M., Röpke F. K., Seitenzahl I. R., Hillebrandt W., 2010, *ApJ*, 719, 1067
- Kushnir D., Katz B., Dong S., Livne E., Fernández R., 2013, *ApJ*, 778, L37
- Leonard D. C., Filippenko A. V., Chornock R., Li W., 2002, *AJ*, 124, 2506
- Leonard D. C., Li W., Filippenko A. V., Foley R. J., Chornock R., 2005, *ApJ*, 632, 450
- Livne E., 1990, *ApJ*, 354, L53
- Livne E., 1999, *ApJ*, 527, L97
- Luminet J. P., Pichon B., 1989, *A&A*, 209, 103
- Ma B., Wei P., Shang Z., Wang L., Wang X., 2014, *Astron. Telegram*, 5794
- Maoz D., Mannucci F., Nelemans G., 2014, *ARA&A*, 52, 107
- Maund J. R. et al., 2013, *MNRAS*, 433, L20
- Maund J. R. et al., 2021, *MNRAS*, 503, 312
- Miller J. S., Robinson L. B., Goodrich R. W., 1988, in Robinson L. B., ed., *Instrumentation for Ground-Based Optical Astronomy, Present and Future. The Ninth Santa Cruz Summer Workshop in Astronomy and Astrophysics*, July 13–24, 1987, Lick Observatory. Springer-Verlag, New York, p. 157
- Moll R., Raskin C., Kasen D., Woosley S. E., 2014, *ApJ*, 785, 105
- Niemeyer J. C., Hillebrandt W., Woosley S. E., 1996, *ApJ*, 471, 903
- Nomoto K., 1982a, *ApJ*, 253, 798
- Nomoto K., 1982b, *ApJ*, 257, 780
- Nomoto K., Sugimoto D., Neo S., 1976, *Ap&SS*, 39, L37
- Nomoto K., Thielemann F.-K., Yokoi K., 1984, *ApJ*, 286, 644
- Nugent P., Baron E., Branch D., Fisher A., Hauschildt P. H., 1997, *ApJ*, 485, 812
- Nugent P. E. et al., 2011, *Nature*, 480, 344
- Pakmor R., Kromer M., Röpke F. K., Sim S. A., Ruiter A. J., Hillebrandt W., 2010, *Nature*, 463, 61
- Pakmor R., Hachinger S., Röpke F. K., Hillebrandt W., 2011, *A&A*, 528, A117
- Pakmor R., Kromer M., Taubenberger S., Sim S. A., Röpke F. K., Hillebrandt W., 2012, *ApJ*, 747, L10
- Patat F., Baade D., Höflich P., Maund J. R., Wang L., Wheeler J. C., 2009, *A&A*, 508, 229
- Pellegrino C. et al., 2020, *ApJ*, 897, 159
- Perlmutter S. et al., 1999, *ApJ*, 517, 565

- Phillips M. M., 1993, *ApJ*, 413, L105
- Plewa T., Calder A. C., Lamb D. Q., 2004, *ApJ*, 612, L37
- Raskin C., Kasen D., Moll R., Schwab J., Woosley S., 2014, *ApJ*, 788, 75
- Reinecke M., Hillebrandt W., Niemeyer J. C., 2002, *A&A*, 391, 1167
- Renzini A., 1999, in Walsh J. R., Rosa M. R., eds, Proc. ESO Workshop held at Garching. Chemical Evolution from Zero to High Redshift. Springer-Verlag, New York, p. 185
- Riess A. G., 2020, *Nat. Rev. Phys.*, 2, 10
- Riess A. G. et al., 1998, *AJ*, 116, 1009
- Röpke F. K., 2007, *ApJ*, 668, 1103
- Rosswog S., Ramirez-Ruiz E., Hix W. R., 2009, *ApJ*, 695, 404
- Sato Y., Nakasato N., Tanikawa A., Nomoto K., Maeda K., Hachisu I., 2016, *ApJ*, 821, 67
- Schlaflly E. F., Finkbeiner D. P., 2011, *ApJ*, 737, 103
- Schmidt G. D., Elston R., Lupie O. L., 1992, *AJ*, 104, 1563
- Seitzzahl I. R. et al., 2013, *MNRAS*, 429, 1156
- Serkowski K., Mathewson D. S., Ford V. L., 1975, *ApJ*, 196, 261
- Shen K. J., Bildsten L., 2009, *ApJ*, 699, 1365
- Shen K. J., Kasen D., Weinberg N. N., Bildsten L., Scannapieco E., 2010, *ApJ*, 715, 767
- Shen K. J., Kasen D., Miles B. J., Townsley D. M., 2018, *ApJ*, 854, 52
- Soker N., 2019, *New Astron. Rev.*, 87, 101535
- Stevance H. F. et al., 2019, *MNRAS*, 485, 102
- Taam R. E., 1980, *ApJ*, 242, 749
- Tinyanont S. et al., 2021, *Nat. Astron.*, 5, 544
- Tonry J. et al., 2019, Transient Name Server Discovery Report, 2019-3, 1
- van Dokkum P. G., 2001, *PASP*, 113, 1420
- Wang L., Wheeler J. C., 2008, *ARA&A*, 46, 433
- Wang L., Wheeler J. C., Höflich P., 1997, *ApJ*, 476, L27
- Wang L., Howell D. A., Höflich P., Wheeler J. C., 2001, *ApJ*, 550, 1030
- Wang L. et al., 2003, *ApJ*, 591, 1110
- Wang L., Baade D., Höflich P., Wheeler J. C., Kawabata K., Khokhlov A., Nomoto K., Patat F., 2006, *ApJ*, 653, 490
- Wang L., Baade D., Patat F., 2007, *Science*, 315, 212
- Wang X., Li W., Filippenko A. V., Foley R. J., Smith N., Wang L., 2008, *ApJ*, 677, 1060
- Webbink R. F., 1984, *ApJ*, 277, 355
- Whelan J., Iben Jr. I., 1973, *ApJ*, 186, 1007
- Wolff M. J., Nordsieck K. H., Nook M. A., 1996, *AJ*, 111, 856
- Woosley S. E., Weaver T. A., 1994, *ApJ*, 423, 371
- Woosley S. E., Weaver T. A., Taam R. E., 1980, in Wheeler J. C., ed., Proc. Texas Workshop on Type I Supernovae. University of Texas, Austin, TX, p. 96
- Yang Y. et al., 2020, *ApJ*, 902, 46

This paper has been typeset from a \LaTeX file prepared by the author.