## **PERSPECTIVES**

**EARTH SCIENCES** 

# Interdisciplinary approaches based on imaging radar enable cutting-edge cultural heritage applications

Fulong Chen<sup>1,2</sup>, Huadong Guo<sup>1,2,\*</sup>, Deodato Tapete<sup>3</sup>, Nicola Masini<sup>4</sup>, Francesca Cigna<sup>5</sup>, Rosa Lasaponara<sup>6</sup>, Salvatore Piro<sup>4</sup>, Hui Lin<sup>7</sup> and Peifeng Ma<sup>8</sup>


Heritage assets reflecting memories of the past are increasingly sustained by societies as a necessary condition for the continuing evolution of people's culture. The call for 'strengthening efforts to protect and safeguard the world's cultural and natural heritage' in United Nations Sustainable Development Goal (SDG) 11 [1] is a clear acknowledgement of the role of heritage in sustaining resilient societies and lifestyles.

To understand and safeguard irreplaceable cultural assets, archaeological prospection and heritage preservation are two interconnected and essential research domains. Prospection aims at archaeological discovery and documentation, to bring back to light buried and hitherto unknown structures, antiquities and even extensive paleo-landscapes. Preservation orients us to prevent further deterioration and to design conservation measures to enhance heritage sustainability by monitoring the condition of heritage assets through time. Both archaeological investigation and heritage preservation include the use of all possible invasive and non-invasive means.

Over the last few decades, an increasing number of technologies have been applied to facilitate sustainability goals of cultural heritage [2,3], and among them remote sensing is recognized as a viable tool [4] owing to its capacity for noninvasive, multi-scale observations and measurements over large territories. Optical methods have been widely used because the data are intuitive and usually

easy to interpret. As a complementary tool, imaging radar has emerged as an effective means for systematic and continuous long-term remote observations of cultural heritage to improve documentation, monitoring, preservation and management.

Imaging radar includes airborne/ spaceborne Synthetic Aperture Radar (SAR), Ground Penetrating Radar (GPR) and Ground-Based SAR (GB-SAR), whose operation characteristics are generally diverse but complementary, depending on their peculiar observation capabilities coupled with investigation needs and scales (Fig. 1). Following a holistic approach, the usage of these different instrumentations



**Figure 1.** Framework of multi-platform imaging radar in cultural heritage applications with complementary observation views and monitoring scopes.

<sup>©</sup> The Author(s) 2021. Published by Oxford University Press on behalf of China Science Publishing & Media Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

should be conceived according to a multi-sensor/data and multi-scale perspective, although constraints due to data and instrumentation accessibility (e.g. lack of funds) may prevail in some circumstances.

Across large landscapes, satellite SAR imagery is capable of revealing ancient drainage patterns [5], palaeo-channels and other human traces. Confidence in the observed archaeological traces improves as imaging resolution from satellite or aircraft increases. Then archaeological remains can be further detailed using geo-archaeology tools and close-range remote sensing methods, including GPR (in use for over three decades) to explore anthropogenic layers and determine their depths [6]. Hence, such a multi-scale SAR-GPR integrated approach could be effective for archaeological site detection across palaeolandscapes (Fig. 1). In this domain, SAR investigations proved successful, especially for analysing cause-effect mechanisms between backscattering signals and archaeological proxy indicators, linked to micro-topographical variations and changes in moisture content.

The benefits brought by technology integration are particularly relevant in the desert and semi-arid regions, where optical remote sensing is limited, whereas SAR can offer large-scale imaging, subsurface penetration and effective feature discrimination. Moreover, today a huge amount of satellite SAR data can be promptly processed using cloud tools (also available as open-access tools such as Google Earth Engine) to automatically identify archaeological traces, using artificial intelligence as machine and deep learning approaches [7]. After that, indepth archaeological investigations need to be implemented at the site level, or even in specific subzones of archaeological interest, using high-resolution SAR products and GPR subsurface imaging. The performance of GPR prospection in a dry-arid environment is maximized when soil moisture is low and the penetration of radar signatures is enhanced.

Moreover, being sensitive to subtle motions, spaceborne Multitemporal SAR Interferometry (MTInSAR) is recommended for monitoring purposes.

MTInSAR detects millimetric deformation anomalies at the cultural-monument scale [8], providing early-warning signals for identifying zones of structural instability at individual monuments which could then be monitored using the 2D-static (motion) measurements of GB-SAR [9]. Details on issues affecting monument components can be further clarified using the health diagnosis model of portable GPR, which helps identify structural fissures or cavities [10]. The integrated use of MTInSAR-GB-SAR-GPR does improve the resolution and provide multi-scale/level information ranging from the identification of cracks and deformations of masonry structures and artefacts, to the detection of fresco detachment or the identification of the diverse construction phases/techniques of monuments and works of art (Fig. 1).

A dedicated literature review based on the Web of Science database reveals a clear but uneven increase in the use of imaging radar technology in cultural heritage management over the last 30 years; from the first publication in 1992 to  $\sim$ 40 papers/yr between 2015 and 2020, with >80% publications focusing on GPR-based case studies. Surprisingly, despite the intrinsic complementarities, the integrated use of GPR, GB-SAR and airborne-satellite SAR is rare. It is also revealed that collaboration had become a necessary condition for interdisciplinary heritage studies, particularly in complex landscapes and situational contexts that call for knowledge and expertise from scientists, engineers and other stakeholders. Institution-level cooperation has become a mainstream trend in scientific and industrial communities in order to achieve benefits through in-depth theoretical investigations and technical applications.

A viable pathway could be the design of pilot projects enhancing trans- and interdisciplinary collaborations at the international level, as already experienced in the past in the framework of the United Nations Educational, Scientific and Cultural Organization (UNESCO) and European Space Agency (ESA) Open Initiative, which focused on 'the Use of Space Technologies to Support the World Heritage Convention'.

Launch and enlargement of such initiatives would be welcome, to provide an optimum platform on a worldwide scale for building case studies on digital heritage applications using integrated imaging technologies (with particular emphasis on radar). A partnership with international organizations, including UNESCO's Category 1 and 2 centres and institutes, is proposed. It contributes to the execution of UNESCO's programme and addresses the gap of Tier III SDG indicators (i.e. 11.4.1) by way of the International Centre on Space Technologies for Natural and Cultural Heritage (HIST), whose mission is the protection and safeguarding of cultural and natural heritage.

### **FUNDING**

This work was jointly supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDA19030502), the National Key Research and Development Program of China (2017YFE0134400), the National Natural Science Foundation of China (41771489) and CNR FOE Project 'China-Italy Innovation Forum'.

#### Conflict of interest statement. None declared.

Fulong Chen<sup>1,2</sup>, Huadong Guo<sup>1,2,\*</sup>, Deodato Tapete<sup>3</sup>, Nicola Masini<sup>4</sup>, Francesca Cigna<sup>5</sup>, Rosa Lasaponara<sup>6</sup>, Salvatore Piro<sup>4</sup>, Hui Lin<sup>7</sup> and Peifeng Ma<sup>8</sup> <sup>1</sup>Key Laboratory of Digital Earth Science, Aerospace Information Research Institute, Chinese Academy of Sciences, China; <sup>2</sup>International Centre on Space Technologies for Natural and Cultural Heritage, China; <sup>3</sup>Italian Space Agency, Italy; <sup>4</sup>Institute of Science of Cultural Heritage, National Research Council, Italy; <sup>5</sup>Institute of Atmospheric Sciences and Climate, National Research Council, Italy; <sup>6</sup>Institute of Methodologies for Environmental Analysis, National Research Council, Italy; 7School of Geography and Environment, Jiangxi Normal University, China and <sup>8</sup>Institute of Space and Earth Information Science, The Chinese University of Hong Kong, China

#### \*Corresponding author.

E-mail: hdguo@radi.ac.cn

### **REFERENCES**

United Nations. Transforming Our World:
 The 2030 Agenda for Sustainable Development. https://sustainabledevelopment. un.org/post2015/transformingourworld (29 June 2021, date last accessed).

- 2. Reimann L, Vafeidis AT and Brown S *et al. Nat Commun* 2018: **9**: 4161.
- 3. Inomata T, Triadan D and Vázquez López VA *et al. Nature* 2020; **582**: 530–3.
- 4. Aminzadeh B and Samani F. *Remote Sens Environ* 2006; **102**: 52–62.
- Ghoneim E, Benedetti M and El-Baz F. Geomorphology 2012; 139-40: 242–57.
- Gizzi FB and Leucci G. Surv Geophys 2018; 39: 1039–68.
- Orengo HA, Conesa FC and Garcia-Molsosa A et al. Proc Natl Acad Sci USA 2020; 117: 18240– 50
- Chen F, Guo H and Ma P et al. Sci Adv 2017; 3: e1601284.
- Tapete D, Casagli N and Luzi G. J Archaeol Sci 2013; 40: 176–89.
- Ludeno G, Cavalagli N and Ubertini F et al. Surv Geophys 2020; 41: 647–67.

National Science Review 8: nwab123, 2021 https://doi.org/10.1093/nsr/nwab123 Advance access publication 7 July 2021

#### EARTH SCIENCES

# Super pathogens from environmental biotechnologies threaten global health

Yong Xiao <sup>1</sup>, Feng Zhao , Josep Peñuelas<sup>2,3</sup>, Qiansheng Huang and Yong-Guan Zhu<sup>4,5,\*</sup>

Microbial pathogens are viruses, bacteria, fungi and protozoa infecting humans, animals or plants. Super pathogens (also known as superbugs) able to resist multiple antimicrobials and cause fatal infections have been increasingly considered an emerging threat to global health [1]. The COVID-19 pandemic has caused more than 3.8 million deaths as of 15 June 2021, reminding us that we should always be alert to the next pandemic caused by super pathogens. We know little, however, about the origin and transmission of super pathogens in the environment. Among the various potential sources of super pathogens, special attention should be paid to environmental biotechnologies (EBTs) for treating wastewater.

Human activities generate wastewater, with a global production of  $\sim$ 330 km³/year [2], that contains heavy metals, carbohydrates, antimicrobials and pathogens polluting the environment and spreading diseases. Various methods have been developed to purify wastewater, but EBTs are the most common ones because they use microbes to degrade pollutants economically, i.e. at a cost of <0.1 US\$/m³ municipal wastewater in China.

EBTs are valuable for purifying wastewater, but they carry a risk of incubating and releasing super pathogens. Millions of microbial species exist in

EBTs, many of which are pathogens playing important roles in degrading pollutants [3]. EBTs, e.g. municipal wastewater-treatment plants, have been widely reported as hotspots of pathogens and antimicrobial resistance genes (ARGs) [4]. During long-term use of EBTs, pathogens can evolve to resist multiple pollutants, which unfortunately enhances their resistance to antimicrobials, because microbes use similar strategies to resist antimicrobials and toxic pollutants such as heavy metals [5]. Microbial genes can also be transferred between microbes [6], enabling pathogens to acquire new ARGs from others to become super pathogens. Furthermore, traditional disinfection can also increase the spread of ARGs between different microbes.

Fecal coliforms or *Escherichia coli* are widely used microbial indexes for the discharge of treated wastewater, e.g. 100–10 000 colony-forming units/L in some countries. The abundance of fecal coliforms in EBTs is low, e.g. <0.22% in most wastewater-treatment plants [3], suggesting that a substantial volume of other microbes, including super pathogens, is released to the natural environment and then spread around the world by human activities [7]. For example, super pathogens that produce New Delhi metallo-beta-lactamase 1

(NDM-1) conferring resistance to multiple antimicrobials are spread to rivers via effluent from wastewater-treatment plants lacking proper sanitation [8]. Super pathogens from EBTs can be transferred to humans mainly via accidental contact with reclaimed water and bioaerosols from EBTs and foods contaminated by EBT effluent. We therefore call for close cooperation between the research community, industry and government to reduce the threat to global health caused by super pathogens from EBTs (Fig. 1).

'Know yourself and know your enemy, and you will never be defeated', as said in Sunzi Bingfa (The Art of War). First, we must better understand the evolution of super pathogens in EBTs on a global scale. Many studies have investigated the dynamics of ARGs in the ecosystem, but little is known about super pathogens. Monitoring the emission of super pathogens from EBTs and tracking their fates in various environments is important for evaluating their risks to human health. Such research can inform the public with regard to avoiding environments and products contaminated by EBT effluent, and most importantly, can support governments to appropriately supervise the EBT industry. Highthroughput sequencing of nucleic acids provides community-level information,