

Molecular Phylogenetic and Expression Analysis of the Complete WRKY Transcription Factor Family in Maize

KAI-FA Wei^{1,*}, JUAN Chen¹, YAN-FENG Chen¹, LING-JUAN Wu¹, and DAO-XIN Xie²

Department of Biological Sciences and Biotechnology, Zhangzhou Normal University, 36 Xian Qian Zhi Street, Zhangzhou 363000, Fujian, China¹ and School of Life Sciences, Tsinghua University, Beijing 10084, China²

*To whom correspondence should be addressed. Tel. +86 13960150031 Fax. +86 596-2528735.
E-mail: kaifa-wei@163.com

Edited by Kazuo Shinozaki
(Received 11 October 2011; accepted 28 December 2011)

Abstract

The WRKY transcription factors function in plant growth and development, and response to the biotic and abiotic stresses. Although many studies have focused on the functional identification of the WRKY transcription factors, much less is known about molecular phylogenetic and global expression analysis of the complete WRKY family in maize. In this study, we identified 136 WRKY proteins coded by 119 genes in the B73 inbred line from the complete genome and named them in an orderly manner. Then, a comprehensive phylogenetic analysis of five species was performed to explore the origin and evolutionary patterns of these WRKY genes, and the result showed that gene duplication is the major driving force for the origin of new groups and subgroups and functional divergence during evolution. Chromosomal location analysis of maize WRKY genes indicated that 20 gene clusters are distributed unevenly in the genome. Microarray-based expression analysis has revealed that 131 WRKY transcripts encoded by 116 genes may participate in the regulation of maize growth and development. Among them, 102 transcripts are stably expressed with a coefficient of variation (CV) value of <15%. The remaining 29 transcripts produced by 25 WRKY genes with the CV value of >15% are further analysed to discover new organ- or tissue-specific genes. In addition, microarray analyses of transcriptional responses to drought stress and fungal infection showed that maize WRKY proteins are involved in stress responses. All these results contribute to a deep probing into the roles of WRKY transcription factors in maize growth and development and stress tolerance.

Key words: WRKY transcription factor; maize; phylogenetic analysis; expression profile; development

1. Introduction

The WRKY family is one of the 10 largest transcription factor families in higher plants and extends all over the green lineage.¹ WRKY proteins possess one or two unique DNA-binding domain with nearly 60 amino acids composed of the absolutely conserved signature WRKYGQK followed by a novel zinc-finger-like motif C₂H₂ (C-X₄₋₅-C-X₂₂₋₂₃-H-X-H) or C₂HC (C-X₇-C-X₂₃-H-X-C). As a transcription factor, WRKY factors act in concert with other

components of the transcriptional machinery and direct the temporal- and spatial-specific expression of the designated genes, thereby ensuring proper cellular responses to both internal and external stimuli.¹⁻³ The stimuli may be from abiotic, such as cold, heat, UV-B, drought, wound, etc., or may be from biotic, such as bacteria, fungi and viruses. As well, WRKY proteins have been implicated in the regulation of developmental processes such as trichome,⁴ seed development,⁵ embryogenesis,⁶ leaf senescence,^{7,8} dormancy,⁹ plant growth¹⁰ and metabolic

pathways.^{4,11} Since the first WRKY cDNA, *SPF1*, was cloned from sweet potato (*Ipomoea batatas*),¹² more and more WRKY genes in various plant species have been experimentally identified, including wild oat (*Avena fatua*; *ABF1*),¹³ parsley (*Petroselinum crispum*; *PcWRKY1*, 2, 3),^{14,15} *Arabidopsis thaliana* (*ZAP1*),¹⁶ orchardgrass (*Dactylis glomerata*),¹⁷ tobacco (*Nicotiana tabacum*),^{18–21} chamomile (*Matricaria chamomilla*),²² rice (*Oryza sativa*),²³ sugarcane (*Saccharum* hybrid cultivar),²⁴ bittersweet nightshade (*Solanum dulcamara*),²⁵ potato (*Solanum tuberosum*),^{26,27} wheat (*Triticum aestivum*)²⁸ and grape (*Vitis vinifera*).²⁹ *Zea mays* (hereafter called maize) is not only a vital crop, but also serves as an important model system for basic biological research.^{30,31} Due to its wide ecological potential and excellent characters, maize roots all over the world under a wide range of environmental conditions, occupying 167 million hectares worldwide and producing 872 million tons of grains in July, 2011 according to the USDA (<http://www.fas.usda.gov/psdonline/>). Recently, the maize B73 genome sequence is now available to public,³² providing a good opportunity to study WRKY transcription factor family extensively.

To date, researchers are still active in this field to reveal the underlying molecular mechanism of WRKY proteins in plant growth and development, as well as to elucidate a huge interaction web involving in response to stresses which is mediated by this transcription factor family.^{33–38} The WRKY proteins in model plant *A. thaliana* (hereafter called *Arabidopsis*) have set a good example for later researchers to study the transcription factor family. In *Arabidopsis*, 72 characterized genes can be categorized into three groups based on the number of WRKY domains and the pattern of the zinc finger motif according to Somssich and colleagues.³⁹ The first group contains two WRKY domains (N-terminal and C-terminal), whereas the other two groups have only one domain. Group III differs in the pattern of the zinc finger motif because it has the zinc finger motif of C₂HC rather than C₂H₂ in the other groups. Group II proteins can be further subdivided into five subgroups according to the amino acid motifs outside the WRKY domain. All these preliminary knowledge might contribute to our understanding of maize WRKY gene family.

In this study, we identified 136 WRKY proteins encoded by 119 WRKY genes from the maize B73 genome and named them from ZmWRKY1 to ZmWRKY119 based on their loci on corresponding chromosomes. Considering of the different transcripts produced by the same gene, totally 136 putative proteins are all named. These 136 WRKY proteins in maize can be classified into three groups in terms of the phylogenetic tree which is constructed using the

WRKY domain peptide sequences. The phylogenetic comparisons of maize, *Arabidopsis*, *O. sativa* (hereafter called rice), *Hordeum vulgare* (hereafter called barley), *Physcomitrella patens* (hereafter called *P. patens*) and ancestral eukaryotic organisms reveal the origin and evolutionary expansion of the WRKY family. The maize chromosome sequence information derived from databases is applied to map the 136 WRKY transcripts to their corresponding gene locus in the 10 haploid chromosomes in order to investigate the duplication events occurred on maize chromosomes.

Nowadays, a large number of maize microarray experiments are accessible via different public resources such as GEO or Plexdb, and allow for *in silico* expression analyses of maize genes. The microarray data of genome-wide gene expression atlas of the maize inbred line B73 offer the chance to understand the distinct expression patterns of the WRKY family under stresses response and growth and development. All these favourable conditions enable us to further our research and get a deep understanding of the roles of WRKY family members in maize.

2. Materials and methods

2.1. Identification and annotation of WRKY family in maize

Since the amended maize B73 genome sequence is now publicly available, we can download the genome sequence (*ZmB73_5b_FGS_genes.fasta.gz*) and the proteome sequence (*ZmB73_5b_FGS_translations.fasta.gz*) from the Maize Genome Sequence Project (<http://ftp.maizesequence.org/current/filtered-set/>). Then, local BLASTP program was run to search the complete WRKY members. The proteome sequences are used as a database for local BLASTP search. The *Arabidopsis* WRKY proteins are used as query sequences. The *e*-value for BLASTP was set at $1e-10$ to obtain the final data set of WRKY proteins. After removing the overlapping genes manually, a total of 136 protein sequences are eventually identified as WRKY proteins and named based on their exact loci on chromosomes. The putative orthologues were assigned from *Arabidopsis*, rice and *H. vulgare* with significance under $1e-20$ from National Center of Biotechnology Information (NCBI).

2.2. Phylogenetic analysis of WRKY family

The WRKY family protein sequences of rice (*O. sativa japonica*) and *Arabidopsis* were downloaded from TIGR rice database (<http://www.tigr.org/>) and TAIR (The *Arabidopsis* Information Resource; <http://www.arabidopsis.org/>), respectively. Sequences of barley and *P. patens* were extracted from NCBI. Maize sequences were identified by local BLASTP program

as described above. Phylogenetic and molecular evolutionary analyses were conducted using MEGA version 5⁴⁰ with pairwise distance and Neighbor-Joining algorithm. The evolutionary distances were computed using the *p*-distance method and were used to estimate the number of amino acid substitutions per site. The reliability of each tree was established by conducting 1000 bootstrap sampling steps.

2.3. Mapping WRKY genes on maize chromosomes

The maize databases were used in a BLAST-based search of the entire maize genomic sequence to confirm the physical locations of all WRKY genes. The Genome Pixelizer software was used for a graphical display of the WRKY loci in each pair of corresponding maize chromosomes (<http://atgc.org/GenomePixelizer/>⁴¹).

2.4. Microarray-based expression analysis

Transcriptome data of genome-wide gene expression atlas of the maize inbred line B73 made by NimbleGen microarray technology, together with the normalized expression data using RMA (robust multi-chip average) algorithm, were downloaded from the publicly available databases such as MaizePLEX (<http://www.plexdb.org/plex.php?database=Corn>) and GEO (<http://www.ncbi.nlm.nih.gov/geo/>). The RMA data which were log₂-transformed beforehand were loaded into R and Bioconductor for expression analysis (<http://www.bioconductor.org/>⁴²). The package Limma^{43,44} was used to model the systematic parts of data by fitting a linear model in the function lmFit. The model was specified by a design matrix. The empirical Bayes approach, in the function eBayes, was used to moderate the standard errors of the estimated log-fold changes and shrink the standard errors to a common value. Then, the gplots⁴⁵ package was used to make the heat map.

3. Results and discussion

3.1. Identification and nomenclature of 136 maize WRKY proteins

A keyword search against the NCBI and UniProt protein sequence databases returned 34 and 31 previously annotated maize WRKY protein sequences, respectively, by the year 2011; among which, 28 proteins are derived from the maize B73 inbred line (Supplementary Table S1). We greatly appreciate the foregoing efforts made by Alexandrov *et al.*⁴⁶ and other scientists, who have named these sequences which were derived from a large-scale cDNA sequencing or from experiments individually. However, the annotations are so disordered and redundant, for they are named neither in chromosome order nor in

the group order and the same numbers occur repeatedly, which make research work difficult to go ahead. A uniform nomenclature with consecutive numbering may help avoid confusion and should help communicate in the scientific community. Thus, we renamed the 136 protein sequences which we have identified through an overall search of the complete genome sequence from ZmWRKY1 to ZmWRKY119 based on the exact position of their corresponding genes on chromosomes 1–10 and from top to bottom, while the variant proteins produced from the same locus are recognized by the same name added 1, 2 or 3 behind (Supplementary Table S2). We use the similar numbering system as proposed for bZIP⁴⁷ and R2R3-MYB,⁴⁸ provides a unique identifier for each WRKY gene. Genes in different genomes that evolved from a common ancestral gene by speciation can be considered as orthologous genes,⁴⁹ which retain the similar functions during evolution.⁵⁰ As a good plant for experiments, *Arabidopsis*, together with monocotyledonous plants rice and barley, is used to assign orthology for maize, which is valuable for functional analysis of the WRKY family members in maize. The putative orthologues are identified according to their *e*-value (under $1e-20$) and the topology of phylogenetic tree.

3.2. Evolutionary analysis of the WRKY transcription factor family

Exposed to a wide variety of biotic and abiotic stresses connected with plant sessile, autotrophic lifestyle, the WRKY family appears to have been under positive selection pressure during evolution. To study the evolutionary origin of maize WRKY family and classify the newly discovered members, multiple sequence alignment is performed using the complete domain sequences derived from *Arabidopsis* (71),³⁹ maize (136), rice (82),^{51,52} barley (36)⁵³ and *P. patens* (36). Because of the difference between the C- and the N-terminal WRKY domain sequences of Group I, which will be described later, we separate them as two independent domains. Then the alignment result is used to generate a phylogenetic tree (Fig. 1). The result shows a well-organized classification according to that in *Arabidopsis*.³⁹ Here, we call them Group I, IIa–e and III.

The result of phylogenetic analysis implies that domain gain and loss is a divergent force for expansion of the WRKY gene family. It has been reported that WRKY transcription factors have their evolutionary origin in ancient eukaryotes with the most basal WRKY genes identified in two non-photosynthetic organisms, the unicellular protist *Giardia lamblia* and the slime mold *Dictyostelium discoideum*.⁵⁴ Each ancestral eukaryotic organism has single WRKY




Figure 1. Unrooted Neighbor-Joining phylogenetic tree of WRKY domains in five plants. The tree is reconstructed using WRKY domain sequences from the complete WRKY gene families in *A. thaliana* (yellow), *Z. mays* (B73, brown), *O. sativa* (blue), *H. vulgare* (green) and *P. patens* (khaki). The evolutionary distances are computed using the *p*-distance method and are in the units of the number of amino acid substitutions per site.

protein which belongs to Group I because of the two WRKY domains, suggesting that the proteins of this group evolved early and represent the ancestral form. Except AtWRKY10, all WRKY proteins of Group I in *Arabidopsis* have two WRKY domains. But in monocotyledons, the loss of WRKY domain seems to be more common. For example, Supplementary Fig. S1 shows that Group I contains 16 proteins having only one domain, among which 9 proteins possess the C-terminal domain and the rest with the N-terminal domain only. In addition, some proteins with two domains are clustered into Group III according to the phylogenetic tree, which are OsWRKY41, OsWRKY61, OsWRKY63, OsWRKY81 and HvWRKY25. As the phenomenon occurs in monocotyledons exclusively, the result shows that there exists a divergence between monocot, and demonstrates that Group III proteins emerged later during evolution and maybe evolved from Group I.

The early researches have reported that only the C-terminal domain in Group I has the sequence-specific DNA-binding activity with the W-box, and N-terminal WRKY domain shows weaker binding activity.^{3,16,55} The first solution structure of the WRKY domain was AtWRKY4_C (PDB: 1WJ2), obtained by NMR spectroscopy, which confirmed the

reports described above.⁵⁶ Further, DNA titration experiment proved that the region corresponding to the conserved sequence WRKYGQK in WRKY4-C is directly involved in DNA binding. In contrast, the N-terminal WRKY domain might participate in the binding process or alternatively provide an interface for protein–protein interactions that coincide with the function of some zinc-finger-like domains.⁵⁷ In this scenario, we can reasonably assume that the N-terminal WRKY domain seems to be more variable during evolution, so it may be deleted from the sequence or evolve into another pattern to achieve additional functions. Meanwhile, it can be of no question that the single WRKY domains of Groups II and III family members are more similar to the C-terminal WRKY domain of Group I proteins for their common function of DNA binding.

A large number of WRKY proteins exist in the five species, suggesting that they play crucial roles in plant developmental and physiological processes (Table 1).⁵⁸ The evidence shows that the rapid duplication of WRKY genes occurred before the divergence of monocots and dicots.⁵⁴ It might be the environmental pressures that impelled WRKY family to enlarge in order to adapt to new or changing environments. In *G. lamblia* and *D. discoideum*, only a single


Table 1. Number of WRKY domains and proteins (numbers in parenthesis) of *Arabidopsis*, rice, maize, barley and *P. patens* in the WRKY Groups I–III

WRKY group	AtWRKY	OsWRKY	ZmWRKY	HvWRKY	PpWRKY
I_N	13	9	22	4	3
I_C	14	11	21	5	3
IIa	3	4	7	4	0
IIb	8	7	11	1	6
IIc	18	19	29	10	18
IId	7	6	14	5	4
IIe	8	8	17	2	0
III	14	26	31	12	5
Total	85 (71)	90 (78)	152 (136)	43 (39)	40 (37)

WRKY protein has been discovered. And then, in bryophyte, the family member of *P. patens* has increased to 37 by duplication. There is no related report about the members of Group III in *P. patens* ever before. However, on the basis of the complete genome sequencing and the topology of the phylogenetic tree, five proteins are found to group into Group III. To angiosperm, monocotyledonous plants develop a larger family of WRKY proteins than dicotyledonous plants. One hundred and thirty-six maize WRKY proteins form the largest WRKY family, while those of rice ranks second, only 71 WRKY members in *Arabidopsis*. With plants from low to higher and from aquatic environment to terrestrial environment, plants suffering severer stresses shut down basic growth functions, while WRKY proteins may act synergistically with other family members to promote plant development and reinforce resistance to environment stressors by modulating the expression of genes. As an important transcription factors, the rapid duplication of WRKY family members may contribute to the increase in adaptability and the establishment of signal transduction webs to go through the adversity.

3.3. Chromosome mapping and genetic analysis

There are 10 chromosomes in maize genome and the length of each chromosome varies from each other. The first chromosome is the longest with 301.0 Mb, and the last is the shortest which is only half of the longest. According to the physical and genetic framework of the maize B73 genome, the 10 chromosomes of the maize genome are structurally diverse and have undergone dynamic changes in chromatin composition.^{30,32} Mapping genes to individual chromosome can facilitate the elimination of redundant data, and provide a powerful analytical method to conduct genetic research. In order to map 119 WRKY genes to maize chromosomes, the physical location of each gene was required. Many

**Figure 2.** Histogram showing the number and distribution of WRKY genes of three groups on 10 chromosomes.

searches are performed in the interest of finding gene orientation and genetic distance of corresponding WRKY genes against maize-related databases. Finally, the software GenomePixelizer was used to visualize the distribution of all WRKY genes on maize 10 chromosomes. The genes are presented by small colour boxes, and the different colours mean different groups and subgroups of the WRKY gene family.

The physical distribution of WRKY genes in the maize B73 genome illustrates the genetic events which result in the diversity and complexity of this gene family. The study of chromosome map (Supplementary Fig. S2) along with the histogram (Fig. 2) comes to the conclusion that the WRKY genes are dispersedly distributed across all the chromosomes in maize. The highest number of WRKY genes is located on Chromosome 8 (24 genes), accounting for 20.17% of the total. The chromosome having least number is Chromosome 9, with only four genes belonging to three different groups, accounting for just 3.36% of the total. The members of three groups are distributed on all chromosomes except Chromosome 5 on which WRKY genes of Group II are located exclusively. Without regard to Chromosome 5, the distribution of Group I genes in each chromosome is more even than that of Groups II and III, with the number of genes ranging from 2 to 3. The formation of this distribution pattern implies that some genetic events of this gene family have taken place during evolution.

The analysis of genetics and genomics has offered insights into the generation and the evolution of WRKY genes. Evidences have shown that the dramatic variations in size and distribution of the WRKY gene family are influenced by many processes, including gene duplication owing to large-scale genome events such as polyploidy, segmental duplications and tandem duplication.^{59,60} An important fact is that maize genome has undergone several rounds of genome duplication, including an ancient paleopolyploid event ~70 million years ago (mya)⁶¹ and an additional whole-genome duplication event ~11

mya.^{31,62,63} Furthermore, many studies have revealed that transcriptional factors are more likely to retain after duplication.^{64–66} All these suggest that genome duplication might be a major mechanism for expansion of WRKY family in maize. Our inference can be confirmed by another gramineous plant, rice, in which the WRKY family comprises ~77.7% duplicated regions.⁶⁷ Most WRKY genes show clustering in maize genome. Here, we adopted the standpoint proposed by Meyers *et al.*⁶⁸ to define a cluster that the numbers or sizes of clusters changed little when the maximum number of intervening open-reading frames (ORFs) increased to 20 or even 50. Therefore, two or more WRKY genes that occurred within a maximum of 40 ORFs were identified as a cluster in this study. Totally, 47 WRKY genes forming 20 gene clusters distribute dispersedly on the overall chromosomes except Chromosomes 5 and 9 (Supplementary Table S3). The remaining 72 genes are singletons. Strikingly, Chromosome 8 not only owns the most of WRKY genes but also possesses nearly half of gene clusters. In general, we can classify the clusters into three kinds. In the first kind, eight gene clusters form monophyletic tandem duplications with non-WRKY genes intervening, including seven clusters in Group III and one in subgroup IIa. In the second kind, seven clusters are made up of four subgroups except IIa, forming three types of gene pairs. Among them, three are II-b, c pair type; three II-c, e pair and one II-d, e pair. The third kind is mixed by two groups. Beyond that, the rest singletons maybe occurred by random events.

According to Xu *et al.*,⁶⁹ if two closed related genes which are located within the same chromosomal region apart from each other fewer than 20 genes can be reckoned as tandem duplication. In this scenario, 16 tandem duplications in our study are inside the gene cluster regions. Interestingly, *ZmWRKY78* and *ZmWRKY91* are the same gene distributed on Chromosomes 7 and 8, respectively, which may suggest the occurrence of segmental duplication event in maize WRKY family during evolution. The increased number of WRKY genes correlates closely with the frequency of duplication events, and these duplications may be responsible for the maintenance of transcriptional regulation activity. More copies of one gene will contribute to offset the effects of mutations and help the plants to survive adversity.

3.4. Expression analysis of WRKY factors in global transcriptome at different developmental stages and in specific organs

Researches have revealed a multiple roles of WRKY factors in response to abiotic stresses, including drought and salt, which is regarded as an ancestral

role of WRKY proteins,⁴ and biotic stresses such as bacteria and fungi. Elucidating the functions of WRKY family members require great endeavour. Microarray is one of the useful global transcriptome analysis technologies which provides an opportunity to understand the patterns of gene expression. Many expression data generated by microarrays are publicly available in microarray databases, such as GEO and ArrayExpress, which provide valuable resources for gene discovery and functional characterization. We mine the gene expression data under drought stress shown in Supplementary Fig. S3 and during *Ustilago maydis* infection shown in Supplementary Fig. S4.

The gene expression profiling of 31 *ZmWRKYs* was compared between the drought-tolerant line Han21 and the drought-sensitive line Ye478 (Supplementary Fig. S3). Interestingly, we can notice that the gene expression levels of the drought-tolerant line Han21 in generally change less than that of the drought-sensitive line Ye478 and recovered more quickly when the seedlings were re-watered. We can see that the expression of *ZmWRKY115* is apparently decreased under drought stress, which may be a response to the increased ABA level resulting from drought. Simultaneously, some genes show a cumulative trend under drought, such as *ZmWRKY 15.1*, *-46*, *-52* and *-77*. Interestingly, *ZmWRKY 90* in the drought-sensitive line Ye478 tend to be expressed at higher mRNA abundances under any conditions. Meanwhile, many WRKY proteins are recognized as pathogen-induced transcription factors and active in immune response such as *AtWRKY18*, *AtWRKY22*, *AtWRKY29*, *AtWRKY53*, *AtWRKY54* and *AtWRKY70*. Supplementary Figure S4 displays the expression levels of maize WRKY genes during infection with *U. maydis*. Compared with the mock stimulus, most of the *ZmWRKYs* exhibit a gradual increase in expression levels with the prolonging of infection time. For example, the *ZmWRKY78* and *91* are two orthologues of *AtWRKY53* which are activated by the pathogen-triggered SA signalling pathway, showing an obvious increasing trend during the infection with *U. maydis*. *AtWRKY70* functions as an activator of SA-dependent defence genes and a repressor of JA-regulated genes, and the expression levels of its orthologue, *ZmWRKY115*, are also apparently increased upon the fungal infection, implying a parallel function in defence response.

It is of great significance to study the expression patterns and regulative mechanisms of WRKYs in stress responses. However, in the work described here, we mainly focus on the spatial and temporal specific expression patterns of maize WRKY genes in order to identify their roles in developmental regulation, as the expression of WRKY genes was detected in a wide variety of plant species and is involved in plant




Figure 3. Line graph showing the CV of the signal intensities of 131 WRKY transcripts across 60 developmental stages. Normalized signal intensities of 131 ZmWRKY transcripts are displayed (according to MaizePlex), and corresponding ZmWRKY transcripts are presented on the x-axis with the order according to the groups. The transcripts with $CV \geq 15\%$ are represented by a tiny hollow circle. The reference gene which encodes E2 enzyme is shown as a small square.

growth and development.^{5,70–71} In this case, we mined microarray data that record the gene expression levels of 60 distinct tissues representing 11 major organ systems and varying developmental stages of the maize (Supplementary Table S4). The organ systems included the germinating seed, primary root, whole seedling, stem and shoot apical meristem, internodes, cob, tassel and anthers, silk, leaf, husk and seed.

A total number of 131 probes detected in R could be assigned to 131 corresponding ZmWRKY transcripts (Supplementary Table S5). The remaining five transcripts with no detectable expression signal are *ZmWRKY25.1*, *ZmWRKY25.2*, *ZmWRKY25.3*, *ZmWRKY32* and *ZmWRKY102*. There are three major reasons for why their signals cannot be detected. First, the different versions of the official maize genome sequence are used in the study. The version of B73 maize genome that are used to map probe set design sequences on the microarray is 5a (RefGen_v2), whereas the version we used to identify the complete WRKY family is 5b (RefGen_v2). The second plausible explanation is that these genes are very specific to organs or developmental stages which are not covered in these experiments so that no signal intensities are detected on the microarray. Last, it is also possible that these genes could be stress-inducible or pseudogenes. In order to investigate the temporal and spatial transcription patterns of the WRKY genes in maize life cycle, a hierarchy cluster was performed to visualize global transcription profile of the WRKY genes across the 60 developmental stages, and the results were illustrated as a heat map (Supplementary Fig. S5).

It can be seen from the heat map that all of the 131 detected transcripts produced by 116 genes are involved in many biological processes, and expressed in all tissues, but their expression levels are distinct. Most of the genes appear to be invariable and lowly expressed among all tissues. Interestingly, the

reference gene encoding E2 enzyme is the most stably and highly expressed across maize life, for which is known as a component in the ubiquitin-dependent protein degradation pathway. The stable gene expression across all tissues can be regarded as constitutive expression, and we can infer that many maize WRKY genes were expressed at low level, which may work synergistically with other family proteins during plant growth and development.

In order to investigate the differential expression patterns, especially fluctuating patterns, in different developmental stages and diverse tissues, we calculate the coefficient of variation (CV value; $CV = sd/mean$, where *sd* and *mean* represent the standard deviation and mean expression level of a gene across all the tissues, respectively) of each gene in 60 tissues accurately (Fig. 3), and the results show that the CV values of these genes range from 2 to 28% (Supplementary Table S6). Those with a CV value of $<15\%$ are considered to be the least expression variability.⁷³ Consequently, total 102 transcripts are stably expressed across all tissues. In contrast, the CV values of the rest 29 transcripts are $>15\%$, which indicate the existence of stage-specific gene expression during development. As we know that WRKY factors can act as either negative or positive regulators in gene expressions thus we are more concerned about the dynamic changes in gene expression during the development of major maize organs. Therefore, we highlight the genes with a CV value of $>15\%$ and explore their expression levels among the distinct organs in order to find the organ-specific genes in maize (data see Supplementary Table S5).

The 11 organ systems can be further subdivided into single tissues which represent the gradual developmental process to some extent. Figure 4 shows the expression levels of the 29 maize WRKY genes in 60 tissues with a CV value of $>15\%$. We ordered the transcripts based on their groups in order to find differential expression between groups. And as expected,




Figure 4. Heat map showing the signal intensities of 29 ZmWRKY transcripts with CV $\geq 15\%$ in 11 specific organs. The alphabet represents specific developmental stages: A, germinating seed; B, primary root; C, whole seedling; D, stem and SAM; E, internode; F, cob; G, tassel and anthers; H, leaves; I, husk; J, seed.




Figure 5. Line graph of the signal intensities of six interest ZmWRKY transcripts with CV $\geq 20\%$ in 11 specific organs, as well as three reference transcripts.

genes in the same group appear to share the similar expression patterns as a whole, while about half of 29 WRKY gene expressions show that variation could be generated by different organ types. Expression of *ZmWRKY92* appears to be accumulated in the late-development leaves stage. Both *ZmWRKY116.2* and *ZmWRKY116.3* are highly expressed in the anther and silk, and also highly increased in early whole seed after pollination and before the formation of the endosperm and embryo. Notably, *ZmWRKY28* in IIc is remarkably expressed in the stem and SAM, internode, and cob and tassel, and its expression level is gradually changed in consistent with the development of the relevant organs. *ZmWRKY68* is highly expressed in late-development leaf and husk. It is notable that there is no gene detected in IIId with a CV value of $>15\%$. *ZmWRKY39* in subgroup IIc is highly expressed in the primary root, and late leaf and husk. However, the WRKY genes in Group III display a leaf-specific gene expression. Besides, the expression of *ZmWRKY78* and *ZmWRKY91* may also

be specific in the pericarp while *ZmWRKY115* perhaps plays a special role in early whole seed. In addition to the highly expressed genes in specific organs, the genes with significant low expression levels may also have effects on the development of given organs. Most of the genes are likely to be stably expressed at low levels in the development of the endosperm and embryo, suggesting that they are not stage-specific and tissue-specific expression.

At the same time, we also care about the most variable gene expressions with a CV value of $>20\%$. The line graph in Fig. 5 will do a great help to observe the interest genes and analyse their dynamic changing processes clearly. We use *E2 enzyme* as an external reference and *ZmWRKY23* and *ZmWRKY105* as internal references to show the constitutively high expression and constitutively low expression, respectively. It is very interesting that *ZmWRKY12* and *ZmWRKY28* form an arc line, respectively, revealing their important roles in the development of the stem and shoot apical meristems, and the internode

and cob, and *ZmWRKY12* is also highly expressed in the pericarp. *ZmWRKY56* has the highest expression level in the germinating seed, while *ZmWRKY113* shows a fluctuating pattern of expression during the development of the leaf.

4. Conclusions

Data acquisition of WRKY transcription factor family is the initial step to go on our analysis of the gene family. Meanwhile, numbering and annotating the identified genes are also of great importance for classification and function characterization. However, a big block stand on our way is the existing identifiers in databases. In database like NCBI, the data of maize WRKY are submitted from large-scale sequencing or experiments individually. The numbering of these data is disorder and redundant ineluctably, and also the data are from several different maize lines, so the numbering system may be diverse and lineage-specific. As the maize B73 genome has been sequenced, we propose to number from chromosomes 1 to 10 and from top to bottom, which can help avoid confusion and promote communication. On this occasion, we give the 136 putative proteins uniform identifier.

Among the 136 WRKY proteins we have identified, 27 proteins belong to Group I, 78 proteins are contained in Group II and 31 proteins are members of Group III. The classification system is based on the features of WRKY domains which can be regarded as evolutionary unit. The sequence divergence usually goes with functional difference in the course of evolution. And it is the divergence of both expression and protein sequence that makes the lineage-specific expansion. Thus, we can conclude that proteins from different groups may accomplish diverse functions.

The comparative analysis of orthologues may help us obtain the information of the evolutionary relationships among the gene family members and help us predict the potential functions of putative proteins. The high-throughput screening method identified that *AtWRKY53* is expressed at a very early time point of leaf senescence.⁸ Not only *AtWRKY53*, *AtWRKY4*, -6, -7 and -11 in *Arabidopsis* play a role in leaf development.³⁹ Among the WRKY genes with the CV value of >15%, *ZmWRKY116.2*, -116.3 are orthologues of *AtWRKY4*, and *ZmWRKY77*, -78, -91 are orthologues of *AtWRKY53*. However, their functions seem to diverge from each other. As seen in the heat map, *ZmWRKY77*, -78 -91 are highly expressed in the leaf and present a gradual cumulative trend, implying a regulatory role of these genes in leaf development, especially in the early stage of leaf senescence. While the specific expression of

ZmWRKY116.2 and -116.3 emerge mainly in reproductive-related tissues, such as the anther and silk. In addition, *ZmWRKY116.2* and -116.3 are also orthologues of *AtWRKY44* which are associated with the tannin and mucilage production in the seed coat.^{4,73} Thus, both genes in maize are expressed at a high level in the early development of the whole seed but not in the development of the endosperm and embryo. Endogenesis abscisic acid (ABA) is crucial in adversity responses as well as growth and development. During the late maturity, the seed enters into the dormant phase due to the increase of ABA. While after imbibition, ABA content usually rapidly decreases and the seed begins to germinate. This ABA-dependent progress of the seed germination and post-germination growth was mediated by *AtWRKY2*.⁷² In maize, *ZmWRKY56* is the orthologue of *AtWRKY2*. Coincidentally, *ZmWRKY56* is highly expressed in the germinating seed after 24 h imbibition, suggesting a similar function of these two orthologues. It has been proved that *AtWRKY40* functions as a central negative regulator of ABA signalling in seed germination and post-germination,³⁶ while its orthologues in maize seem insensitive to drought stress in our study.

By means of the microarray-based data mining and homologous analysis, we can obtain much useful information about the putative functions of the WRKYs in maize. It is of great importance to elucidate the biological functions of these transcription factors and provide us deeper understanding in molecular mechanisms of corn growth development and adversity resistance.

Supplementary data: Supplementary data are available at www.dnaresearch.oxfordjournals.org.

Acknowledgements: We are grateful to the providers who submitted the microarray data to the public expression databases which can be applied freely.

Funding

The project was supported by the Natural Science Foundation of Fujian Province, China (Grant No. B0810040).

References

1. Ulker, B. and Somssich, I.E. 2004, WRKY transcription factors: from DNA binding towards biological function, *Curr. Opin. Plant Biol.*, **7**, 491–8.
2. Ciolkowski, I., Wanke, D., Birkenbihl, R.P. and Somssich, I.E. 2008, Studies on DNA-binding selectivity

- of WRKY transcription factors lend structural clues into WRKY-domain function, *Plant Mol. Biol.*, **68**, 81–92.
3. Eulgem, T., Rushton, P.J., Schmelzer, E., Hahlbrock, K. and Somssich, I.E. 1999, Early nuclear events in plant defence signalling: rapid gene activation by WRKY transcription factors, *EMBO J.*, **18**, 4689–99.
 4. Johnson, C.S., Kolevski, B. and Smyth, D.R. 2002, *TRANSPARENT TESTA GLABRA2*, a trichome and seed coat development gene of *Arabidopsis*, encodes a WRKY transcription factor, *Plant Cell*, **14**, 1359–75.
 5. Luo, M., Dennis, E.S., Berger, F., Peacock, W.J. and Chaudhury, A. 2005, *MINISEED3 (MINI3)*, a WRKY family gene, and *HAIKU2 (IKU2)*, a leucine-rich repeat (LRR) KINASE gene, are regulators of seed size in *Arabidopsis*, *Proc. Natl Acad. Sci. USA*, **102**, 17531–6.
 6. Lagacé, M. and Matton, D.P. 2004, Characterization of a WRKY transcription factor expressed in late torpedo-stage embryos of *Solanum chacoense*, *Planta*, **219**, 185–9.
 7. Robatzek, S. and Somssich, I.E. 2002, Targets of AtWRKY6 regulation during plant senescence and pathogen defense, *Genes Dev.*, **16**, 1139–49.
 8. Hinderhofer, K. and Zentgraf, U. 2001, Identification of a transcription factor specifically expressed at the onset of leaf senescence, *Planta*, **213**, 469–73.
 9. Pnueli, L., Hallak-Herr, E., Rozenberg, M., et al. 2002, Molecular and biochemical mechanisms associated with dormancy and drought tolerance in the desert legume *Retama raetam*, *Plant J*, **31**, 319–30.
 10. Chen, C. and Chen, Z. 2002, Potentiation of developmentally regulated plant defense response by AtWRKY18, a pathogen-induced *Arabidopsis* transcription factor, *Plant Physiol.*, **129**, 706–16.
 11. Willmott, R.L., Rushton, P.J., Hooley, R. and Lazarus, C.M. 1998, DNase1 footprints suggest the involvement of at least three types of transcription factors in the regulation of alpha-Amy2/A by gibberellin, *Plant Mol. Biol.*, **38**, 817–25.
 12. Ishiguro, S. and Nakamura, K. 1994, Characterization of a cDNA encoding a novel DNA-binding protein, SPF1, that recognizes SP8 sequences in the 5' upstream regions of genes coding for sporamin and beta-amylase from sweet potato, *Mol. Gen. Genet.*, **244**, 563–71.
 13. Rushton, P.J., Macdonald, H., Huttly, A.K., Lazarus, C.M. and Hooley, R. 1995, Members of a new family of DNA-binding proteins bind to a conserved cis-element in the promoters of alpha-Amy2 genes, *Plant Mol. Biol.*, **29**, 691–702.
 14. Rushton, P.J., Torres, J.T., Parniske, M., Wernert, P., Hahlbrock, K. and Somssich, I.E. 1996, Interaction of elicitor-induced DNA-binding proteins with elicitor response elements in the promoters of parsley PR1 genes, *EMBO J.*, **15**, 5690–700.
 15. Cormack, R.S., Eulgem, T., Rushton, P.J., Kochner, P., Hahlbrock, K. and Somssich, I.E. 2002, Leucine zipper-containing WRKY proteins widen the spectrum of immediate early elicitor-induced WRKY transcription factors in parsley, *Biochim. Biophys. Acta*, **1576**, 92–100.
 16. De Pater, S., Greco, V., Pham, K., Memelink, J. and Kijne, J. 1996, Characterization of a zinc-dependent transcriptional activator from *Arabidopsis*, *Nucleic Acids Res.*, **24**, 4624–31.
 17. Alexandrova, K.S. and Conger, B.V. 2002, Isolation of two somatic embryogenesis-related genes from orchard-grass (*Dactylis glomerata*), *Plant Sci.*, **162**, 301–7.
 18. Chen, C. and Chen, Z. 2000, Isolation and characterization of two pathogen- and salicylic acid-induced genes encoding WRKY DNA-binding proteins from tobacco, *Plant Mol. Biol.*, **42**, 387–96.
 19. Hara, K., Yagi, M., Kusano, T. and Sano, H. 2000, Rapid systemic accumulation of transcripts encoding a tobacco WRKY transcription factor upon wounding, *Mol. Gen. Genet.*, **263**, 30–7.
 20. Wang, Z., Yang, P., Fan, B. and Chen, Z. 1998, An oligo selection procedure for identification of sequence-specific DNA-binding activities associated with the plant defence response, *Plant J.*, **16**, 515–22.
 21. Yoda, H., Ogawa, M., Yamaguchi, Y., Koizumi, N., Kusano, T. and Sano, H. 2002, Identification of early-responsive genes associated with the hypersensitive response to tobacco mosaic virus and characterization of a WRKY-type transcription factor in tobacco plants, *Mol. Genet. Genomics*, **267**, 154–61.
 22. Ashida, Y., Nishimoto, M., Matsushima, A., Watanabe, J. and Hirata, T. 2002, Molecular cloning and mRNA expression of geraniol-inducible genes in cultured shoot primordia of *Matricaria chamomilla*, *Biosci. Biotechnol. Biochem.*, **66**, 2511–4.
 23. Kim, C.Y., Lee, S.H., Park, H.C., et al. 2000, Identification of rice blast fungal elicitor-responsive genes by differential display analysis, *Mol. Plant Microbe Interact.*, **13**, 470–4.
 24. MR, L. 2001, In silico differential display of defense-related expressed sequence tags from sugarcane tissues infected with diazotrophic endophytes, *Genet. Mol. Biol.*, **24**, 103–11.
 25. Huang, T. and Duman, J.G. 2002, Cloning and characterization of a thermal hysteresis (antifreeze) protein with DNA-binding activity from winter bittersweet nightshade, *Solanum dulcamara*, *Plant Mol. Biol.*, **48**, 339–50.
 26. Beyer, K., Binder, A., Boller, T. and Collinge, M. 2001, Identification of potato genes induced during colonization by *Phytophthora infestans*, *Mol. Plant Pathol.*, **2**, 125–34.
 27. Dellagi, A., Helibronn, J., Avrova, A.O., et al. 2000, A potato gene encoding a WRKY-like transcription factor is induced in interactions with *Erwinia carotovora* subsp. *atroseptica* and *Phytophthora infestans* and is coregulated with class I endochitinase expression, *Mol. Plant Microbe Interact.*, **13**, 1092–101.
 28. Sun, C., Palmqvist, S., Olsson, H., Boren, M., Ahlandsberg, S. and Jansson, C. 2003, A novel WRKY transcription factor, SUSIBA2, participates in sugar signaling in barley by binding to the sugar-responsive elements of the *iso1* promoter, *Plant Cell*, **15**, 2076–92.
 29. Marchive, C., Mzid, R., Deluc, L., et al. 2007, Isolation and characterization of a *Vitis vinifera* transcription factor, VvWRKY1, and its effect on responses to fungal pathogens in transgenic tobacco plants, *J. Exp. Bot.*, **58**, 1999–2010.

30. Wei, F., Zhang, J., Zhou, S., et al. 2009, The physical and genetic framework of the maize B73 genome, *PLoS Genet.*, **5**, e1000715.
31. Gaut, B.S., Le Thierry d'Ennequin, M., Peek, A.S. and Sawkins, M.C. 2000, Maize as a model for the evolution of plant nuclear genomes, *Proc. Natl Acad. Sci. USA*, **97**, 7008–15.
32. Schnable, P.S., Ware, D., Fulton, R.S., et al. 2009, The B73 maize genome: complexity, diversity, and dynamics, *Science*, **326**, 1112–5.
33. Shen, Y.Y., Wang, X.F., Wu, F.Q., et al. 2006, The Mg-chelatase H subunit is an abscisic acid receptor, *Nature*, **443**, 823–6.
34. Rushton, P.J., Somssich, I.E., Ringler, P. and Shen, Q.J. 2010, WRKY transcription factors, *Trends Plant Sci.*, **15**, 247–58.
35. Eulgem, T. and Somssich, I.E. 2007, Networks of WRKY transcription factors in defense signaling, *Curr. Opin. Plant Biol.*, **10**, 366–71.
36. Shang, Y., Yan, L., Liu, Z.Q., et al. 2010, The Mg-chelatase H subunit of *Arabidopsis* antagonizes a group of WRKY transcription repressors to relieve ABA-responsive genes of inhibition, *Plant Cell*, **22**, 1909–35.
37. Ryu, H.S., Han, M., Lee, S.K., et al. 2006, A comprehensive expression analysis of the WRKY gene superfamily in rice plants during defense response, *Plant Cell Rep.*, **25**, 836–47.
38. Pandey, S.P. and Somssich, I.E. 2009, The role of WRKY transcription factors in plant immunity, *Plant Physiol.*, **150**, 1648–55.
39. Eulgem, T., Rushton, P.J., Robatzek, S. and Somssich, I.E. 2000, The WRKY superfamily of plant transcription factors, *Trends Plant Sci.*, **5**, 199–206.
40. Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. 2011, MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods, *Mol. Biol. Evol.*, **28**, 2731–9.
41. Kozik, A., Kochetkova, E. and Michelmores, R. 2002, GenomePixelizer—a visualization program for comparative genomics within and between species, *Bioinformatics*, **18**, 335–6.
42. Team, R.D.C. 2011, *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing: Vienna, Austria.
43. Smyth, G.K. 2004, Linear models and empirical Bayes methods for assessing differential expression in microarray experiments, *Stat. Appl. Genet. Mol. Biol.*, **3**, Article 3.
44. Smyth, G.K. 2005, *Bioinformatics and Computational Biology Solutions using R and Bioconductor*. Springer: New York.
45. Warnes, G.R. 2010, Includes R source code and/or documentation contributed by: Ben Bolker (in alphabetical order), L. B., Robert Gentleman, Wolfgang Huber Andy Liaw, Thomas Lumley, Martin Maechler, Arni Magnusson, Steffen Moeller, Marc Schwartz and Bill Venables, gplots: Various R programming tools for plotting data. R package version 2.8.0.
46. Alexandrov, N.N., Brover, V.V., Freidin, S., et al. 2009, Insights into corn genes derived from large-scale cDNA sequencing, *Plant Mol. Biol.*, **69**, 179–94.
47. Nijhawan, A., Jain, M., Tyagi, A.K. and Khurana, J.P. 2008, Genomic survey and gene expression analysis of the basic leucine zipper transcription factor family in rice, *Plant Physiol.*, **146**, 333–50.
48. Stracke, R., Werber, M. and Weisshaar, B. 2001, The R2R3-MYB gene family in *Arabidopsis thaliana*, *Curr. Opin. Plant Biol.*, **4**, 447–56.
49. Fitch, W.M. 1970, Distinguishing homologous from analogous proteins, *Syst. Zool.*, **19**, 99–113.
50. Li, W.H., Yang, J. and Gu, X. 2005, Expression divergence between duplicate genes, *Trends Genet.*, **21**, 602–7.
51. Ross, C.A., Liu, Y. and Shen, Q.J. (2007), The WRKY Gene Family in Rice (*Oryza sativa*), *J. Integr. Plant Biol.*, **49**, 827–42.
52. Xie, Z., Zhang, Z.L., Zou, X., et al. 2005, Annotations and functional analyses of the rice WRKY gene superfamily reveal positive and negative regulators of abscisic acid signaling in aleurone cells, *Plant Physiol.*, **137**, 176–89.
53. Mangelsen, E., Kilian, J., Berendzen, K.W., et al. 2008, Phylogenetic and comparative gene expression analysis of barley (*Hordeum vulgare*) WRKY transcription factor family reveals putatively retained functions between monocots and dicots, *BMC Genomics*, **9**, 194.
54. Wu, K.L., Guo, Z.J., Wang, H.H. and Li, J. 2005, The WRKY family of transcription factors in rice and *Arabidopsis* and their origins, *DNA Res.*, **12**, 9–26.
55. Maeo, K., Hayashi, S., Kojima-Suzuki, H., Morikami, A. and Nakamura, K. 2001, Role of conserved residues of the WRKY domain in the DNA-binding of tobacco WRKY family proteins, *Biosci. Biotechnol. Biochem.*, **65**, 2428–36.
56. Yamasaki, K., Kigawa, T., Inoue, M., et al. 2005, Solution structure of an *Arabidopsis* WRKY DNA binding domain, *Plant Cell*, **17**, 944–56.
57. Mackay, J.P. and Crossley, M. 1998, Zinc fingers are sticking together, *Trends Biochem. Sci.*, **23**, 1–4.
58. Babu, M.M., Iyer, L.M., Balaji, S. and Aravind, L. 2006, The natural history of the WRKY-GCM1 zinc fingers and the relationship between transcription factors and transposons, *Nucleic Acids Res.*, **34**, 6505–20.
59. Lynch, M. and Conery, J.S. 2000, The evolutionary fate and consequences of duplicate genes, *Science*, **290**, 1151–5.
60. Paterson, A.H., Bowers, J.E. and Chapman, B.A. 2004, Ancient polyploidization predating divergence of the cereals, and its consequences for comparative genomics, *Proc. Natl Acad. Sci. USA*, **101**, 9903–8.
61. Blanc, G. and Wolfe, K.H. 2004, Widespread paleopolyploidy in model plant species inferred from age distributions of duplicate genes, *Plant Cell*, **16**, 1667–78.
62. Swigonova, Z., Lai, J., Ma, J., et al. 2004, Close split of sorghum and maize genome progenitors, *Genome Res.*, **14**, 1916–23.
63. Blanc, G. and Wolfe, K.H. 2004, Functional divergence of duplicated genes formed by polyploidy during *Arabidopsis* evolution, *Plant Cell*, **16**, 1679–91.

64. Thomas, B.C., Pedersen, B. and Freeling, M. 2006, Following tetraploidy in an *Arabidopsis* ancestor, genes were removed preferentially from one homeolog leaving clusters enriched in dose-sensitive genes, *Genome Res.*, **16**, 934–46.
65. Wu, Y., Zhu, Z., Ma, L. and Chen, M. 2008, The preferential retention of starch synthesis genes reveals the impact of whole-genome duplication on grass evolution, *Mol. Biol. Evol.*, **25**, 1003–6.
66. Ramamoorthy, R., Jiang, S.Y., Kumar, N., Venkatesh, P.N. and Ramachandran, S. 2008, A comprehensive transcriptional profiling of the WRKY gene family in rice under various abiotic and phytohormone treatments, *Plant Cell Physiol.*, **49**, 865–79.
67. Meyers, B.C., Kozik, A., Griego, A., Kuang, H. and Michelmore, R.W. 2003, Genome-wide analysis of NBS-LRR-encoding genes in *Arabidopsis*, *Plant Cell*, **15**, 809–34.
68. Xu, G., Ma, H., Nei, M. and Kong, H. 2009, Evolution of F-box genes in plants: different modes of sequence divergence and their relationships with functional diversification, *Proc. Natl Acad. Sci. USA*, **106**, 835–40.
69. Ohashi, Y., Oka, A., Ruberti, I., Morelli, G. and Aoyama, T. 2002, Entopically additive expression of *GLABRA2* alters the frequency and spacing of trichome initiation, *Plant J.*, **29**, 359–69.
70. Xing, Y., Jia, W. and Zhang, J. 2008, AtMKK1 mediates ABA-induced *CAT1* expression and H₂O₂ production via AtMPK6-coupled signaling in *Arabidopsis*, *Plant J.*, **54**, 440–51.
71. Jiang, W. and Yu, D. 2009, *Arabidopsis* WRKY2 transcription factor mediates seed germination and postgermination arrest of development by abscisic acid, *BMC Plant Biol.*, **9**, 96.
72. Sekhon, R.S., Lin, H., Childs, K.L., et al. 2011, Genome-wide atlas of transcription during maize development, *Plant J.*, **66**, 553–63.
73. Ishida, T., Hattori, S., Sano, R., et al. 2007, *Arabidopsis* *TRANSPARENT TESTA GLABRA2* is directly regulated by R2R3 MYB transcription factors and is involved in regulation of *GLABRA2* transcription in epidermal differentiation, *Plant Cell*, **19**, 2531–43.