

Should we individualize lipid profiling in women with polycystic ovary syndrome?

Inger Sundström Poromaa^{1,*}, Jan Roar Mellembakken²,
Laure Morin-Papunen^{3,4}, Terhi Piltonen^{3,4}, Johanna Puurenen^{3,4},
Juha S. Tapanainen^{3,4,5}, Elisabet Stener-Victorin⁶,
Angelica Lindén Hirschberg⁷, Eszter Vanky^{8,9}, Pernille Ravn¹⁰,
Dorte Glintborg¹¹, and Marianne Andersen¹¹

¹Department of Women's and Children's Health, Uppsala University, Uppsala, Sweden ²Section for Reproductive Medicine, Department of Gynaecology, Women's Division, Oslo University Hospital, Oslo, Norway ³Department of Obstetrics and Gynaecology, Oulu University Hospital, University of Oulu, Oulu, Finland ⁴PEDEGO Research Unit, Medical Research Centre, Oulu, Finland ⁵Department of Obstetrics and Gynaecology, Helsinki University Hospital and University of Helsinki, Helsinki, Finland ⁶Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden ⁷Department of Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden ⁸Department of Obstetrics and Gynaecology, St. Olav's Hospital, Trondheim University Hospital, Trondheim, Norway ⁹Department of Laboratory Medicine, Children's and Women's Health, Norwegian University of Science and Technology, Trondheim, Norway ¹⁰Department of Gynaecology and Obstetrics, Odense University Hospital, Odense, Denmark ¹¹Department of Endocrinology, Odense University Hospital, Odense, Denmark

*Correspondence address. Department of Women's and Children's Health, Uppsala University, Uppsala, Sweden.
E-mail: Inger.sundstrom@kbh.uu.se

Hum Reprod 2016;**31**:2791–2795

The authors would like to apologize for a typing error in the surname of one of the authors of this article. Laure M Papunen should be listed as Laure Morin-Papunen.

The authors would like to assure readers that this does not affect any content of the article.